

Product Datasheet

Nucleolin Antibody (364-5) - Azide and BSA Free NBP2-47861-0.1mg

Unit Size: 0.1 mg

Store at -20 to -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-47861

Updated 10/23/2024 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-47861



NBP2-47861-0.1mg


Nucleolin Antibody (364-5) - Azide and BSA Free

Product Information	
Unit Size	0.1 mg
Concentration	1.0 mg/ml
Storage	Store at -20 to -80C. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	364-5
Preservative	No Preservative
Isotype	IgG1 Kappa
Purity	Protein A or G purified
Buffer	10 mM PBS
Target Molecular Weight	76 kDa
Product Description	
Description	1.0 mg/ml of antibody purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WITH BSA & azide (NBP2-44612). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C.
Host	Mouse
Gene ID	4691
Gene Symbol	NCL
Species	Human, Bovine (Negative), Mouse (Negative), Rat (Negative)
Reactivity Notes	Does not react with Mouse, Rat or Bovine.
Marker	Marker of Human Cells
Specificity/Sensitivity	Recognizes a protein of ~76kDa, which is identified as Nucleolin (NCL). It is the major nucleolar phosphoprotein of growing eukaryotic cells. NCL is located mainly in dense fibrillar regions of the nucleolus. It is found associated with intranucleolar chromatin and pre-ribosomal particles. Human NCL gene consists of 14 exons with 13 introns and spans approximately 11kb. It induces chromatin decondensation by binding to histone H1. It is thought to play a role in pre-rRNA transcription and ribosome assembly. This monoclonal antibody can be used to stain the nucleoli in cell or tissue preparations and can be used as a marker of the nucleoli in subcellular fractions. It produces a speckled pattern in the nuclei of cells of normal and malignant cells and may be used to stain the nucleoli of cells in fixed or frozen tissue sections. It can be used with paraformaldehyde fixed frozen tissue or cell preparations and formalin fixed, paraffin-embedded tissue sections.
Immunogen	lysate of SU-DHL-1 Nuclei (Uniprot: P19338)
Product Application Details	
Applications	Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence


Recommended Dilutions	Western Blot 0.5 - 1.0 ug/ml, Flow Cytometry 0.5 - 1 ug/million cells in 0.1 ml, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin 0.25 - 0.5 ug/ml, Immunofluorescence 0.5 - 1.0 ug/ml, CyTOF-ready
Application Notes	Immunohistochemistry (Formalin-fixed): 0.25-0.5ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes. Optimal dilution for a specific application should be determined.

Images


Immunocytochemistry/Immunofluorescence: Nucleolin Antibody (364-5) - Azide and BSA Free [NBP2-47861] - Immunofluorescence Analysis of PFA-fixed MCF-7 cells stained with Nucleolin Antibody (364-5) followed by Goat anti-Mouse IgG-CF488 (Green). Membrane is stained with Phalloidin-CF640.




Immunohistochemistry-Paraffin: Nucleolin Antibody (364-5) - Azide and BSA Free [NBP2-47861] - Human Skin stained with Nucleolin Monoclonal Antibody (364-5).




Flow Cytometry: Nucleolin Antibody (364-5) - Azide and BSA Free [NBP2-47861] - Flow Cytometry of Human Nucleolin Ag on 293T cells. Black: cells alone; Grey: Isotype Control; Green: AF488-labeled Nucleolin Antibody (364-5).




Immunocytochemistry/Immunofluorescence: Nucleolin Antibody (364-5) - Azide and BSA Free [NBP2-47861] - Immunofluorescence Analysis of PFA-fixed K562 cells stained with Nucleolin Antibody (364-5) followed by Goat anti-Mouse IgG-CF488 (Green). Membrane is stained with Phalloidin-CF640.




Immunocytochemistry/Immunofluorescence: Nucleolin Antibody (364-5) - Azide and BSA Free [NBP2-47861] - Immunofluorescence Analysis HeLa cells labeling Nucleolin with Nucleolin Antibody (364-5) followed by Goat anti-Mouse IgG-CF488 (Green). The nuclear counterstain is DAPI (Blue).




Immunohistochemistry-Paraffin: Nucleolin Antibody (364-5) - Azide and BSA Free [NBP2-47861] - Formalin-fixed paraffin-embedded human Colon Carcinoma stained with AF488 conjugate of Nucleolin Monoclonal Antibody (364-5).



Immunohistochemistry-Paraffin: Nucleolin Antibody (364-5) - Azide and BSA Free [NBP2-47861] - Human Tonsil stained with Nucleolin Monoclonal Antibody (364-5).



Flow Cytometry: Nucleolin Antibody (364-5) - Azide and BSA Free [NBP2-47861] - Flow Cytometry of Human Nucleolin Ag on 293T cells. Black: cells alone; Grey: Isotype Control; Green: AF488-labeled Nucleolin Antibody (364-5).





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-47861-0.1mg

HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB720-B	Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]
NBP1-43319-0.5mg	Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1)
NBP3-21345PEP	Nucleolin Recombinant Protein Antigen

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-47861

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

