Product Datasheet

CD43/Sialophorin Antibody (SPN/839) [Biotin] NBP2-47879B

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-47879B

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-47879B



NBP2-47879B

CD43/Sialophorin Antibody (SPN/839) [Biotin]

Unit Size0.1 mlConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonallyMonoclonalCloneSPN/839Preservative0.6% Sodium AzideIsotypeIgG1 KappaConjugateBiotinPurityProtein A or G purifiedBufferPotein A or G purifiedBufferThis conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size ated on the datasheet.HostMouseGene ID6933Gene BymbolSPNSpecificity/SensitivityIt recognizes a cell surface glycoprotein of 95/115/135KDa (depending upon the extent of glycosylation). Identified as CD43. No reactivity has been observed with reactive B-cell population. When CD43 antibody is used upons, expresidel you hour, especially a low-grade lymphoma, expresidel you shorma, especially a low-grade lymphoma, expresidel you shorma, especially a low-grade lymphoma, especially	Product Information	
services. Storage Store at 4C in the dark. Clonality Monoclonal Clone SPN/839 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate Biotin Purity Protein A or G purified Buffer Bo Poduct Description This conjugate is made on demand. Actual recovery may vary from the stated valume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 6693 Gene Symbol SPN Specificity/Sensitivity Recombinant full recovery context or express cD43. No reactivity has been observed with reactive B-cell syndo and trom 22-37% of B-cell lymphomas express CD43. No reactivity has been observed with reactive B-cells of a theody of the oret hymphomas and from 22-37% of B-cell lymphomas express CD43. No reactivity has been observed with reactive B-cells of a theody is used in combination with anti-CD20, effective immunophenotyping of the lymphomas and from 22-37% of B-cell lymphomas express CD43. No reactivity has been observed with reactive B-cells on a theodats No reactivity has been observed with reactive B-cells on a botatined. Co-expresses CD43 antioody is used in combination with anti-CD20, effective immunophenotyping of the lymphomas and from abotation with anti-CD20, effective immunophenotyping of the lymphomas and from abotation co-expresses CD43. No reactive B-cel	Unit Size	0.1 ml
ClonalityMonoclonalCloneSPN/839Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateBiotinPurityProtein A or G purifiedBufferPBSProduct DescriptionThis conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.HostMouseGene ID6693Gene BD6693SpeciesHumanMarkerT-Cell MarkerSpecificity/SensitivityIt recognizes a cell surface glycoprotein of 95/115/135kDa (depending upon the extent of glycosylation), identified as CD43. 70-90% of T-cell lymphomas and fromali-fixed tissues can be obtained. Co-staining of a lymphoid infiltrate with anti-CD20 and anti-CD43 argues against a reactive process and favors a diagnosis of lymphoma.ImmunogenRecombinant full-length human CD43/Sialophorin protein (Uniprot: P16150)Product Application DetailsApplicationsApplicationsWestern Biot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Parafin, CyTOF-ready, Immunofluorescence, Immunohistochemistry-Parafin, Lymportic mility of the secence, Immunohistochemistry-Parafin, Immunofluorescence, Immunohistoche	Concentration	
Clone SPN/839 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate Biotin Purity Protein A or G purified Buffer PBS Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 6693 Gene Symbol SPN Species Human Marker T-Cell Marker Specificity/Sensitivity It recognizes a cell surface glycoprotein of 95/115/135kDa (depending upon the extent of glycosylation), identified as CD43. 70-90% of T-cell lymphomas and from 22-37% of B-cell lymphomas express CD43. No reactivity has been observed with reactive B-cells. So a B-lineage population that co-expresses CD43 is highly likely to be an alignant lymphoma, especially a low-grade lymphoma, rather than a reactive B-cell population. When CD43 antibody is used in combination with anti-CD20, effective immunophenotyping of the lymphomas in formalin-fixed tissues can be obtained. Co-staining of a lymphoid infiltrate with anti-CD20 and anti-CD20, effective immunophenotyping of the lymphomas in formalin-fixed tissues can be obtained. Co-staining of a lymphoid infiltrate with anti-CD20 and anti-CD20, effective immunophenotyping of the lymphomas in formalin-fixed tissues can be obtained. Co-staining of a lymphoid infiltrate with anti-CD20 and anti-CD	Storage	Store at 4C in the dark.
Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate Biotin Purity Protein A or G purified Buffer PBS Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 6693 Gene Symbol SPN Species Human Marker T-Cell Marker Specificity/Sensitivity It recognizes a cell surface glycoprotein of 95/115/136kDa (depending upon the extent of glycosylation), identified as CD43. 70-90% of T-cell lymphomas and from 22-37% of B-cell lymphomas express CD43. No reactivity has been observed with reactive B-cells. So a B-lineage population that co-expresses cD43 as highly likely to be a malignant lymphoma, especially a low-grade lymphoma, rather than a reactive B-cell population. When CD43 antibody is used in combination with anti-CD20, effective immunophenotyping of the lymphomas in formalin-fixed tissues can be obtained. Co-staining of a lymphoid infiltrate with anti-CD20 and anti-CD20 and anti-CD20 and anti-CD20 and sing infirmation. When CD43 antibody is used in formalin-fixed tissues can be obtained. Co-staining of a lymphoid infiltrate with anti-CD20 and anti-CD20 and anti-CD20 and sing is obtained. Co-staining of a lymphoid infiltrate with anti-CD20 and anti-CD20 and anti-CD20 and guy anti-CD20 and anti-CD20 and anti-CD3 areactive process and favor	Clonality	Monoclonal
IsotypeIgG1 KappaConjugateBiotinPurityProtein A or G purifiedBufferPBSProduct DescriptionThis conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.HostMouseGene ID6693Gene SymbolSPNSpecificity/SensitivityH recognizes a cell surface glycoprotein of 95/115/135kDa (depending upon the extent of glycosylation), identified as CD43. 70-90% of T-cell lymphomas and from 22-37% of B-cell Symbol as D-lineage population that co-expresses CD43 is highly likely to be a malignant lymphoma, especially a low-grade lymphoma, rather than a reactive B-cell population. When CD43 aritbody is used in combination with anti-CD20, effective immunophenotyping of the lymphomas in formalin-fixed tissues can be obtained. Co-staining of a lymphoid infiltrate with anti-CD20 and anti-CD43 argues against a reactive process and favors a diagnosis of lymphoma.ImmunogenRecombinant full-length human CD43/Sialophorin protein (Uniprot: P16150)Product Application Details ApplicationsWestern Blot, Flow Cytometry, Immunohistochemistry/Immunofluorescence, LimmunofluorescenceRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry/Immunofluorescence, CyTOF-ready	Clone	SPN/839
ConjugateBiotinPurityProtein A or G purifiedBufferPBSProduct DescriptionThis conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.HostMouseGene ID6693Gene SymbolSPNSpeciesHumanMarkerT-Cell MarkerSpecificity/SensitivityIt recognizes a cell surface glycoprotein of 95/115/135kDa (depending upon the extent of glycosylation), identified as CD43. 70-90% of T-cell lymphomas and from 22-37% of B-cell lymphomas express CD43. No reactivity has been observed with reactive B-cells. So a B-lineage population. When CD43 antibody is used in combination with anti-CD20, effective immunophenotyping of the lymphomas in formalin-fixed tissues can be obtained. Co-staining of a lymphoid anti-CD20 and anti-CD43 argues against a reactive process and favors a diagnosis of lymphoma.ImmunogenWestern Blot, Flow Cytometry, Immunocytochemistry/Immunofluorescence, ImmunofluorescenceWestern Blot, Flow Cytometry, Immunohistochemistry-Paraffin, CyTOF-ready, Ummunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence, CyTOF-ready	Preservative	0.05% Sodium Azide
Purity Protein A or G purified Buffer PBS Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 6693 Gene Symbol SPN Species Human Marker T-Cell Marker Specificity/Sensitivity It recognizes a cell surface glycoprotein of 95/115/135kDa (depending upon the extent of glycosylation), identified as CD43. No reactivity has been observed with reactive B-cell. So a B-lineage population that co-expresses CD43 is highly likely to be a malignant lymphoma, sepsecially a low-grade lymphomas, rather than a reactive B-cell population. When CD43 antibody is used in combination with anti-CD20, effective immunophenotyping of the lymphomas in formalin-fixed tissues can be obtained. Co-staining of a lymphodi infiltrate with anti-CD20 and anti-CD43 argues against a reactive process and favors a diagnosis of lymphoma. Immunogen Recombinant full-length human CD43/Sialophorin protein (Uniprot: P16150) Product Application Details Western Blot, Flow Cytometry, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofiltorescence, CyTOF-ready.	Isotype	IgG1 Kappa
Buffer PBS Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 6693 Gene Symbol SPN Species Human Marker T-Cell Marker Specificity/Sensitivity It recognizes a cell surface glycoprotein of 95/115/135kDa (depending upon the extent of glycosylation), identified as CD43. 70-90% of T-cell lymphomas and from 22-37% of B-cell lymphomas express CD43. No reactivity has been observed with reactive B-cells. So a B-lineage population that co-expresses CD43 is highly likely to be a malignant lymphoma, especially a low-grade lymphoma, rather than a reactive B-cell supplemotyping of the lymphomas in formalin-fixed tissues can be obtained. Co-staining of a lymphodi infiltrate with anti-CD20 and anti-CD20 arguing arguing a lawnors a diagnosis of lymphoma. Immunogen Recombinant full-length human CD43/Sialophorin protein (Uniprot: P16150) Product Application Details Western Blot, Flow Cytometry, Immunofucerscence, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence, CyTOF-ready. Recommended Dilutions Western Blot, Flow Cytometry, Immunohistochemistry-Paraffin, Immunofluorescence, CyTOF-ready.	Conjugate	Biotin
Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 6693 Gene Symbol SPN Species Human Marker T-Cell Marker Specificity/Sensitivity It recognizes a cell surface glycoprotein of 95/115/135kDa (depending upon the extent of glycosylation), identified as CD43. 70-90% of T-cell lymphomas and from 22-37% of B-cell lymphomas express CD43. No reactivity has been observed with reactive B-cells. So a B-lineage population that co-expresses CD43 is highly likely to be a malignant lymphoma, especially a low-grade lymphoma, rather than a reactive B-cell surfaciency of a lymphoma in formalin-fixed tissues can be obtained. Co-staining of a lymphoid infiltrate with anti-CD20 and anti-CD20, effective immunophenotyping of the lymphomas in formalin-fixed tissues can be obtained. Co-staining of a lymphoid infiltrate with anti-CD20 and anti-CD20 and anti-CD20 and anti-CD20 and anti-CD20. effective incomponentyping of the lymphomas in formalin-fixed tissues can be obtained. Co-staining of a lymphoid infiltrate with anti-CD20 and anti-CD20 and anti-CD20 and anti-CD20. effective incomponentyping of the lymphomas and form set of lymphoma. Immunogen Recombinant full-length human CD43/Sialophorin protein (Uniprot: P16150) Product Application Details Vestern Blot, Flow Cytometry, Immunocytochemistry-Paraffin, CyTOF-ready, Immunofiluorescence, Immunohistochemistry-Paraffin, Immunofluorescence, CyTOF-ready. Recom	Purity	Protein A or G purified
DescriptionThis conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.HostMouseGene ID6693Gene SymbolSPNSpeciesHumanMarkerT-Cell MarkerSpecificity/SensitivityIt recognizes a cell surface glycoprotein of 95/115/135kDa (depending upon the extent of glycosylation), identified as CD43. 70-90% of T-cell lymphomas and from 22-37% of B-cell lymphomas express CD43. No reactivity has been observed with reactive B-cells. So a B-lineage population that co-expresses CD43 is highly likely to be a malignant lymphoma, especially a low-grade lymphoma, rather than a reactive B-cell population. When CD43 antibody is used in combination with anti-CD20, effective immunophenotyping of the lymphomas and informalin-fixed tissues can be obtained. Co-staining of a lymphoid infiltrate with anti-CD20 and anti-CD43 argues against a reactive process and favors a diagnosis of lymphoma.Product Application DetailsWestern Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, ImmunofluorescenceRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence, CyTOF-ready	Buffer	PBS
Volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.HostMouseGene ID6693Gene SymbolSPNSpeciesHumanMarkerT-Cell MarkerSpecificity/SensitivityIt recognizes a cell surface glycoprotein of 95/115/135kDa (depending upon the extent of glycosylation), identified as CD43. 70-90% of T-cell lymphomas and from 22-37% of B-cell Jymphomas express CD43. No reactivity has been observed with reactive B-cells. So a B-lineage population that co-expresses CD43 is highly likely to be a malignant lymphoma, especially a low-grade lymphoma, rather than a reactive B-cell population. When CD43 antibody is used in combination with anti-CD20, effective immunophenotyping of the lymphomas a diagnosis of lymphoma.ImmunogenRecombinant full-length human CD43/Sialophorin protein (Uniprot: P16150)Product Application DetailsWestern Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, ImmunofluorescenceWestern Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, CyTOF-ready	Product Description	
Gene ID6693Gene SymbolSPNSpeciesHumanMarkerT-Cell MarkerSpecificity/SensitivityIt recognizes a cell surface glycoprotein of 95/115/135kDa (depending upon the extent of glycosylation), identified as CD43. 70-90% of T-cell lymphomas and from 22-37% of B-cell lymphomas express CD43. No reactivity has been observed with reactive B-cells. So a B-lineage population that co-expresses CD43 is highly likely to be a malignant lymphoma, especially a low-grade lymphoma, rather than a reactive B-cell population. When CD43 antibody is used in combination with anti-CD20, effective immunophenotyping of the lymphomas in formalin-fixed tissues can be obtained. Co-staining of a lymphoid infiltrate with anti-CD20 and anti-CD43 argues against a reactive process and favors a diagnosis of lymphoma.ImmunogenRecombinant full-length human CD43/Sialophorin protein (Uniprot: P16150)Product Application Details ApplicationsWestern Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, ImmunofluorescenceRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry. Paraffin, CyTOF-ready, Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence, CyTOF-ready	Description	volume of this product. The volume will be greater than or equal to the unit size
Gene SymbolSPNSpeciesHumanMarkerT-Cell MarkerSpecificity/SensitivityIt recognizes a cell surface glycoprotein of 95/115/135kDa (depending upon the extent of glycosylation), identified as CD43. 70-90% of T-cell lymphomas and from 22-37% of B-cell lymphomas express CD43. No reactivity has been observed with reactive B-cells. So a B-lineage population that co-expresses CD43 is highly likely to be a malignant lymphoma, especially a low-grade lymphoma, rather than a reactive B-cell population. When CD43 antibody is used in combination with anti-CD20, effective immunophenotyping of the lymphomas in formalin-fixed tissues can be obtained. Co-staining of a lymphoid infiltrate with anti-CD20 and anti-CD43 argues against a reactive process and favors a diagnosis of lymphoma.ImmunogenRecombinant full-length human CD43/Sialophorin protein (Uniprot: P16150)Product Application DetailsWestern Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, ImmunofluorescenceApplicationsWestern Blot, Flow Cytometry, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence, CyTOF-ready	Host	Mouse
SpeciesHumanMarkerT-Cell MarkerSpecificity/SensitivityIt recognizes a cell surface glycoprotein of 95/115/135kDa (depending upon the extent of glycosylation), identified as CD43. 70-90% of T-cell lymphomas and from 22-37% of B-cell lymphomas express CD43. No reactivity has been observed with reactive B-cells. So a B-lineage population that co-expresses CD43 is highly likely to be a malignant lymphoma, especially a low-grade lymphoma, rather than a reactive B-cell population. When CD43 antibody is used in combination with anti-CD20, effective immunophenotyping of the lymphomas in formalin-fixed tissues can be obtained. Co-staining of a lymphoid infiltrate with anti-CD20 and anti-CD43 argues against a reactive process and favors a diagnosis of lymphoma.ImmunogenRecombinant full-length human CD43/Sialophorin protein (Uniprot: P16150)Product Application DetailsWestern Blot, Flow Cytometry, Immunohistochemistry-Paraffin, CyTOF-ready, ImmunofluorescenceRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, CyTOF-ready	Gene ID	6693
MarkerT-Cell MarkerSpecificity/SensitivityIt recognizes a cell surface glycoprotein of 95/115/135kDa (depending upon the extent of glycosylation), identified as CD43. 70-90% of T-cell lymphomas and from 22-37% of B-cell lymphomas express CD43. No reactivity has been observed with reactive B-cells. So a B-lineage population that co-expresses CD43 is highly likely to be a malignant lymphoma, especially a low-grade lymphoma, rather than a reactive B-cell population. When CD43 antibody is used in combination with anti-CD20, effective immunophenotyping of the lymphomas in formalin-fixed tissues can be obtained. Co-staining of a lymphoid infiltrate with anti-CD20 and anti-CD43 argues against a reactive process and favors a diagnosis of lymphoma.ImmunogenRecombinant full-length human CD43/Sialophorin protein (Uniprot: P16150)Product Application Details ApplicationsWestern Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, ImmunofluorescenceRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence, CyTOF-ready	Gene Symbol	SPN
Specificity/SensitivityIt recognizes a cell surface glycoprotein of 95/115/135kDa (depending upon the extent of glycosylation), identified as CD43. 70-90% of T-cell lymphomas and from 22-37% of B-cell lymphomas express CD43. No reactivity has been observed with reactive B-cells. So a B-lineage population that co-expresses CD43 is highly likely to be a malignant lymphoma, especially a low-grade lymphoma, rather than a reactive B-cell population. When CD43 antibody is used in combination with anti-CD20, effective immunophenotyping of the lymphomas in formalin-fixed tissues can be obtained. Co-staining of a lymphoid infiltrate with anti-CD20 and anti-CD43 argues against a reactive process and favors a diagnosis of lymphoma.ImmunogenRecombinant full-length human CD43/Sialophorin protein (Uniprot: P16150)Product Application DetailsWestern Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, ImmunofluorescenceRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence, CyTOF-ready	Species	Human
extent of glycosylation), identified as CD43. 70-90% of T-cell lymphomas and from 22-37% of B-cell lymphomas express CD43. No reactivity has been observed with reactive B-cells. So a B-lineage population that co-expresses CD43 is highly likely to be a malignant lymphoma, especially a low-grade lymphoma, rather than a reactive B-cell population. When CD43 antibody is used in combination with anti-CD20, effective immunophenotyping of the lymphomas in formalin-fixed tissues can be obtained. Co-staining of a lymphoid infiltrate with anti-CD20 and anti-CD43 argues against a reactive process and favors a diagnosis of lymphoma.ImmunogenRecombinant full-length human CD43/Sialophorin protein (Uniprot: P16150)Product Application Details ApplicationsWestern Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, ImmunofluorescenceRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunofluorescence, Immunofluorescence, CyTOF-ready	Marker	T-Cell Marker
Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence Recommended Dilutions Western Blot, Flow Cytometry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunofluorescence, CyTOF-ready	Specificity/Sensitivity	extent of glycosylation), identified as CD43. 70-90% of T-cell lymphomas and from 22-37% of B-cell lymphomas express CD43. No reactivity has been observed with reactive B-cells. So a B-lineage population that co-expresses CD43 is highly likely to be a malignant lymphoma, especially a low-grade lymphoma, rather than a reactive B-cell population. When CD43 antibody is used in combination with anti-CD20, effective immunophenotyping of the lymphomas in formalin-fixed tissues can be obtained. Co-staining of a lymphoid infiltrate with anti-CD20 and anti-CD43 argues against a reactive process and favors a
ApplicationsWestern Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, ImmunofluorescenceRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence, CyTOF-ready	Immunogen	Recombinant full-length human CD43/Sialophorin protein (Uniprot: P16150)
Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence Recommended Dilutions Western Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence, CyTOF-ready	Product Application Details	
Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence, CyTOF-ready	Applications	Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready,
Application NotesOptimal dilution of this antibody should be experimentally determined.	Recommended Dilutions	Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence,
	Application Notes	Optimal dilution of this antibody should be experimentally determined.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-47879B

NBP2-29370	Streptavidin Native Protein
IC002B	Mouse IgG1 Isotype Control (11711) [Biotin]
NBP2-23472	Recombinant Human CD43/Sialophorin His Protein
210-TA-005	TNF-alpha [Unconjugated]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-47879B

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

