Product Datasheet

CD30/TNFRSF8 Antibody (Ki-1/779) [FITC] NBP2-47906F

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-47906F

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-47906F



NBP2-47906F

CD30/TNFRSF8 Antibody (Ki-1/779) [FITC]

services. Storage Store at 4C in the dark. Clonality Monoclonal Clone Ki-1/779 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate FITC Purity Protein A or G purified Buffer PBS Product Description Description Description This conjugate is made on demand. Actual recovery may vary from the state volume of this product. The volume will be greater than or equal to the unit is stated on the datasheet. Host Mouse Gene ID 943 Gene Symbol TNFRSF8 Species Human Marker Hodgkin & Reed-Stemberg Cell Marker Specificity/Sensitivity Recognizes a single chain glycoprotein of 105/120kDa, identified as CD30// Complex into a membrane-bound phosphorylated mature 105/120kDa glycoprotein. In Hodgkins disease, CD30/Ki-1 antiggn is expressed by a glycoprotein of activated T and B cells. This monoclonal antibody distinguishes large cell lymphomas derived from activated lymphom cells from histiocytic malignancies and lymphomas derived from resting and precursor in motoclear-Hodgkin and multinucleated T and B cells. This monoclonal antibody distinguishes large cell lymphomas derived from resting and precursor in motoclear-Hodgkin and multinucleated T and B cells. This monoclonal antibody distinguishes large cell lymphomas derive	Product Information	
services. Storage Store at 4C in the dark. Clonality Monoclonal Clone Ki-1/779 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate FITC Purity Protein A or G purified Buffer PBS Product Description This conjugate is made on demand. Actual recovery may vary from the stativolume of this product. The volume will be greater than or equal to the unit a stated on the datasheet. Host Mouse Gene ID 943 Gene Symbol TNFRSF8 Species Human Marker Hodgkin & Reed-Sternberg Cell Marker Specificity/Sensitivity Recognizes a single chain glycoprotein of 105/120kDa, identified as CD30//CD30 is synthesized as a 90kDa precursor, which is processed by mononuclear-Hodgkin and multinucleated Reed-Sternborg cells. It is also expressed by the tumor cells of a majority of anaplastic large cell lymphoma well as by a varying proportion of activated T and B cells. This monoclonal artibody distinguishes large cell lymphomas derived from resting and tobd xi situal contenses. Journal actibody distinguishes large cell lymphomas derived from resting and tobd y distinguishes large cell lymphomas derived from resting and tobd y distinguishes large cell lymphomas derived from resting and tobd y distinguishes large cell lymphomas derived	Unit Size	0.1 ml
Clonality Monoclonal Clone Ki-1/779 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate FITC Purity Protein A or G purified Buffer PBS Product Description Description Description This conjugate is made on demand. Actual recovery may vary from the state volume of this product. The volume will be greater than or equal to the unit is stated on the datasheet. Host Mouse Gene ID 943 Gene Symbol TNFRSF8 Species Human Marker Hodgkin & Reed-Stemberg Cell Marker Specificity/Sensitivity Recognizes a single chain glycoprotein of 105/120kDa, identified as CD30// CD30 is synthesized as a 90kDa precursor, which is processed in the Golgi complex into a membrane-bound phosphorylated mature 105/120kDa glycoprotein. In Hodgkins disease, CD30/Ki-1 antigen is expressed by mononuclear-Hodgkin and multinucleated Reed-Sternberg cells. It is also expressed by the tumor cells of a majority of anaplastic large cell lymphome well as by a varying proportion of activated T and B cells. This monoclonal antibody distinguishes large cell lymphomas derived from resting and cells from histicocytic malignancies and lymphomas. About on third of Ki-1 positive lymphorid cells of from anaplastic carcinomas. About on thid of Ki-1 positive lymphores large of from instructed l	Concentration	Please see the vial label for concentration. If unlisted please contact technical services.
Clone Ki-1/779 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate FITC Purity Protein A or G purified Buffer PBS Product Description Description Description This conjugate is made on demand. Actual recovery may vary from the state volume of this product. The volume will be greater than or equal to the unit is stated on the datasheet. Host Mouse Gene ID 943 Gene Symbol TNFRSF8 Species Human Marker Hodgkin & Reed-Stemberg Cell Marker Specificity/Sensitivity Recognizes a single chain glycoprotein of 105/120kDa, identified as CD30// CD30 is synthesized as a 90kDa precursor, which is processed in the Golgi complex into a membrane-bound phosphorylated mature 105/120kDa glycoprotein. In Hodgkins disease, CD30/Ki-1 antigen is expressed by mononuclear-Hodgkin and multinucleated Reed-Sternberg cells. It is also expressed by the tumor cells of a majority of anaplastic large cell lymphome well as by a varying proportion of activated T and B cells. This monoclonal antibody distinguishes large cell lymphomas derived from resting and cells from histicocytic malignancies and lymphomas. About one third of Ki-1 positive lymphoid cells of from anaplastic carcinomas. About one third of Ki-1 positive lymphomas lack the leukocyte common antigen (CD45). Immunogen	Storage	Store at 4C in the dark.
Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate FITC Purity Protein A or G purified Buffer PBS Product Description This conjugate is made on demand. Actual recovery may vary from the state volume of this product. The volume will be greater than or equal to the unit is stated on the datasheet. Host Mouse Gene ID 943 Gene Symbol TNFRSF8 Species Human Marker Hodgkin & Reed-Sternberg Cell Marker Specificity/Sensitivity Recognizes a single chain glycoprotein of 105/120kDa, identified as CD30// CD30 is synthesized as a 90kDa precursor, which is processed in the Golg complex into a membrane-bound phosphorylated mature 105/120kDa glycoprotein. In Hodgkins disease, CD30/Ki-1 antigen is expressed by mononuclear-Hodgkin and multinucleated Reed-Sternberg cells. It is also expressed by the tumor cells of a majority of anaplastic large cell lymphoma well as by a varying proportion of activated T and B cells. This monoclonal antibody distinguishes large cell glymphomas derived from activated lymphoma precursor lymphoid cells or from anaplastic carcinomas. About one third of Ki-1 positive lymphomas lack the leukocyte common antigen (CD45). Immunogen Recombinant human CD30/TNFRSF8 protein (Uniprot: P28908) Product Application Details Flow Cytometry, Immunohistochemistry/ Immunofluorescence, Immunohis	Clonality	Monoclonal
Isotype IgG1 Kappa Conjugate FITC Purity Protein A or G purified Buffer PBS Product Description This conjugate is made on demand. Actual recovery may vary from the state volume of this product. The volume will be greater than or equal to the unit is stated on the datasheet. Host Mouse Gene ID 943 Gene Symbol TNFRSF8 Species Human Marker Hodgkin & Reed-Stemberg Cell Marker Specificity/Sensitivity Recognizes a single chain glycoprotein of 105/120kDa, identified as CD30// CD30 is synthesized as a 90kDa precursor, which is processed in the Golgi complex into a membrane-bound phosphorylated mature 105/120kDa glycoprotein. In Hodgkins disease, CD30/Ki-1 antigen is expressed by mononuclear-Hodgkin and multinucleated Reed-Stemberg cells. It is also expressed by the tumor cells of a majority of anaplastic large cell lymphome well as by a varying proportion of activated and the dells. This monoclonal antibody distinguishes large cell lymphomas derived from activated lympho cells from histiocytic malignancies and lymphomas derived from nesting and recursor lymphoid cells or from anaplastic carcinomas. About one third of Ki-1 positive lymphomas lack the leukocyte common antigen (CD45). Immunogen Recombinant human CD30/TNFRSF8 protein (Uniprot: P28908) Product Application Details Flow Cytometry, Immunohistochemistry/Immunofluorescence, Immunofluorescence, Immunohistochemistry-Paraffin, CyTOF-ready, Imm	Clone	Ki-1/779
Conjugate FITC Purity Protein A or G purified Buffer PBS Product Description This conjugate is made on demand. Actual recovery may vary from the stativolume of this product. The volume will be greater than or equal to the unit stated on the datasheet. Host Mouse Gene ID 943 Gene Symbol TNFRSF8 Species Human Marker Hodgkin & Reed-Sternberg Cell Marker Specificity/Sensitivity Recognizes a single chain glycoprotein of 105/120kDa, identified as CD30// CD30 is synthesized as a 90kDa precursor, which is processed in the Goigi complex into a membrane-bound phosphorylated mature 105/120kDa glycoprotein. In Hodgkins disease, CD30/Ki-1 antigen is expressed by mononuclear-Hodgkin and multinucleated Reed-Sternberg cells. This monoclonal antibody distinguishes large cell lymphomas derived from resting and precursor lymphoid cells or from anaplastic large cell lymphom kell as by a varying proportion of activated T and B cells. This monoclonal antibody distinguishes large cell lymphomas derived from resting and precursor lymphoid cells or from anaplastic carcinomas. About one third of Ki-1 positive lymphomas lack the leukocyte common antigen (CD45). Immunogen Recombinant human CD30/TNFRSF8 protein (Uniprot: P28908) Product Application Details Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunofluorescence Applications Flow Cytometry, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluoresce	Preservative	0.05% Sodium Azide
Purity Protein A or G purified Buffer PBS Product Description This conjugate is made on demand. Actual recovery may vary from the state volume of this product. The volume will be greater than or equal to the unit is stated on the datasheet. Host Mouse Gene ID 943 Gene Symbol TNFRSF8 Species Human Marker Hodgkin & Reed-Sternberg Cell Marker Specificity/Sensitivity Recognizes a single chain glycoprotein of 105/120kDa, identified as CD30// CD30 is synthesized as a 90kDa precursor, which is processed in the Golgi complex into a membrane-bound phosphorylated mature 105/120kDa glycoprotein. In Hodgkins disease, CD30/ki-1 antigen is expressed by mononuclear-Hodgkin and multinucleated Reed-Sternberg cells. It is also expressed by the tumor cells of a majority of anaplastic large cell lymphomas derived from activated lymphomas derived from resting and precursor lymphoid cells or from anaplastic carcinomas. About one third of Ki-1 positive lymphomas lack the leukocyte common antigen (CD45). Immunogen Recombinant human CD30/TNFRSF8 protein (Uniprot: P28908) Product Application Details Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunofluorescence, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence	Isotype	IgG1 Kappa
Buffer PBS Product Description This conjugate is made on demand. Actual recovery may vary from the stat volume of this product. The volume will be greater than or equal to the unit stated on the datasheet. Host Mouse Gene ID 943 Gene Symbol TNFRSF8 Species Human Marker Hodgkin & Reed-Sternberg Cell Marker Specificity/Sensitivity Recognizes a single chain glycoprotein of 105/120kDa, identified as CD30// CD30 is synthesized as a 90kDa precursor, which is processed in the Golgi complex into a membrane-bound phosphorylated mature 105/120kDa glycoprotein. In Hodgkins disease, CD30/Ki-1 antigen is expressed by mononuclear-Hodgkin and multinucleated Reed-Sternberg cells. It is also expressed by the tumor cells of a majority of anaplastic large cell lymphoma well as by a varying proportion of activated T and B cells. This monoclonal antibody distinguishes large cell lymphomas derived from resting and precursor lymphoid cells or from anaplastic carcinomas. About one third of Ki-1 positive lymphomas lack the leukocyte common antigen (CD45). Immunogen Recombinant human CD30/TNFRSF8 protein (Uniprot: P28908) Product Application Details Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunofluorescence Recommended Dilutions Flow Cytometry, Immunohistochemistry-Paraffin, Immunofluorescence	Conjugate	FITC
Product Description Description This conjugate is made on demand. Actual recovery may vary from the state volume of this product. The volume will be greater than or equal to the unit is stated on the datasheet. Host Mouse Gene ID 943 Gene Symbol TNFRSF8 Species Human Marker Hodgkin & Reed-Sternberg Cell Marker Specificity/Sensitivity Recognizes a single chain glycoprotein of 105/120kDa, identified as CD30// CD30 is synthesized as a 90kDa precursor, which is processed in the Golgi complex into a membrane-bound phosphorylated mature 105/120kDa glycoprotein. In Hodgkins disease, CD30/Ki-1 antigen is expressed by mononuclear-Hodgkin and multinucleated Reed-Sternberg cells. It is also expressed by the tumor cells of a majority of anaplastic large cell lymphoma well as by a varying proportion of activated T and B cells. This monoclonal antibody distinguishes large cell lymphomas derived from resting and precursor lymphoid cells or from anaplastic carcinomas. About one third of it Ki-1 positive lymphomas lack the leukocyte common antigen (CD45). Immunogen Recombinant human CD30/TNFRSF8 protein (Uniprot: P28908) Product Application Details Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunofluorescence, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence, Immunohistochemistry-Paraffin, Turnunofluorescence, Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence	Purity	Protein A or G purified
DescriptionThis conjugate is made on demand. Actual recovery may vary from the stativolume of this product. The volume will be greater than or equal to the unit stated on the datasheet.HostMouseGene ID943Gene SymbolTNFRSF8SpeciesHumanMarkerHodgkin & Reed-Sternberg Cell MarkerSpecificity/SensitivityRecognizes a single chain glycoprotein of 105/120kDa, identified as CD30// CD30 is synthesized as a 90kDa precursor, which is processed in the Golgi complex into a membrane-bound phosphorylated mature 105/120kDa glycoprotein. In Hodgkins disease, CD30/Ki-1 antigen is expressed by mononuclear-Hodgkin and multinucleated Reed-Sternberg cells. It is also expressed by the tumor cells of a majority of anaplastic large cell lymphoma well as by a varying proportion of activated T and B cells. This monoclonal antibody distinguishes large cell lymphomas derived from activated lympho cells form histiocytic malignancies and lymphomas. About one third of I Ki-1 positive lymphomas lack the leukocyte common antigen (CD45).ImmunogenRecombinant human CD30/TNFRSF8 protein (Uniprot: P28908)Product Application DetailsFlow Cytometry, Immunohistochemistry/Immunofluorescence, ImmunofluorescenceRecommended DilutionsFlow Cytometry, Immunohistochemistry, Immunofluorescence, Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence	Buffer	PBS
volume of this product. The volume will be greater than or equal to the unit is stated on the datasheet.HostMouseGene ID943Gene SymbolTNFRSF8SpeciesHumanMarkerHodgkin & Reed-Sternberg Cell MarkerSpecificity/SensitivityRecognizes a single chain glycoprotein of 105/120kDa, identified as CD30/ CD30 is synthesized as a 90kDa precursor, which is processed in the Golgi complex into a membrane-bound phosphorylated mature 105/120kDa glycoprotein. In Hodgkins disease, CD30/Ki-1 antigen is expressed by mononuclear-Hodgkin and multinucleated Reed-Sternberg cells. It is also expressed by the tumor cells of a majority of anaplastic large cell lymphoma well as by a varying proportion of activated T and B cells. This monoclonal antibody distinguishes large cell lymphomas derived from activated lympho cells from histiocytic malignancies and lymphomas. About one third of Ki-1 positive lymphomas lack the leukocyte common antigen (CD45).ImmunogenRecombinant human CD30/TNFRSF8 protein (Uniprot: P28908)Product Application DetailsFlow Cytometry, Immunohistochemistry/Immunofluorescence, ImmunofluorescenceRecommended DilutionsFlow Cytometry, Immunohistochemistry, Immunofluorescence, Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence	Product Description	
Gene ID 943 Gene Symbol TNFRSF8 Species Human Marker Hodgkin & Reed-Sternberg Cell Marker Specificity/Sensitivity Recognizes a single chain glycoprotein of 105/120kDa, identified as CD30/I CD30 is synthesized as a 90kDa precursor, which is processed in the Golgi complex into a membrane-bound phosphorylated mature 105/120kDa glycoprotein. In Hodgkins disease, CD30/Ki-1 antigen is expressed by mononuclear-Hodgkin and multinucleated Reed-Sternberg cells. It is also expressed by the tumor cells of a majority of anaplastic large cell lymphoma well as by a varying proportion of activated T and B cells. This monoclonal antibody distinguishes large cell lymphomas derived from activated lymphoic cells from histiocytic malignancies and lymphomas derived from resting and precursor lymphoid cells or from anaplastic carcinomas. About one third of the Ki-1 positive lymphomas lack the leukocyte common antigen (CD45). Immunogen Recombinant human CD30/TNFRSF8 protein (Uniprot: P28908) Product Application Details Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunofluorescence Applications Flow Cytometry, Immunohistochemistry.Paraffin, CyTOF-ready, Immunofluorescence Recommended Dilutions Flow Cytometry, Immunohistochemistry.Paraffin, Immunofluorescence	Description	This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.
Gene Symbol TNFRSF8 Species Human Marker Hodgkin & Reed-Sternberg Cell Marker Specificity/Sensitivity Recognizes a single chain glycoprotein of 105/120kDa, identified as CD30/I CD30 is synthesized as a 90kDa precursor, which is processed in the Golgi complex into a membrane-bound phosphorylated mature 105/120kDa glycoprotein. In Hodgkins disease, CD30/Ki-1 antigen is expressed by mononuclear-Hodgkin and multinucleated Reed-Sternberg cells. It is also expressed by the tumor cells of a majority of anaplastic large cell lymphoma well as by a varying proportion of activated T and B cells. This monoclonal antibody distinguishes large cell lymphomas derived from activated lymphoi cells from histiocytic malignancies and lymphomas derived from resting and precursor lymphoid cells or from anaplastic carcinomas. About one third of Ki-1 positive lymphomas lack the leukocyte common antigen (CD45). Immunogen Recombinant human CD30/TNFRSF8 protein (Uniprot: P28908) Product Application Details Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence Recommended Dilutions Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence	Host	Mouse
SpeciesHumanMarkerHodgkin & Reed-Sternberg Cell MarkerSpecificity/SensitivityRecognizes a single chain glycoprotein of 105/120kDa, identified as CD30/l CD30 is synthesized as a 90kDa precursor, which is processed in the Golgi complex into a membrane-bound phosphorylated mature 105/120kDa glycoprotein. In Hodgkins disease, CD30/Ki-1 antigen is expressed by mononuclear-Hodgkin and multinucleated Reed-Sternberg cells. It is also expressed by the tumor cells of a majority of anaplastic large cell lymphoma well as by a varying proportion of activated T and B cells. This monoclonal antibody distinguishes large cell lymphomas derived from activated lymphoi cells from histiocytic malignancies and lymphomas derived from resting and precursor lymphoid cells or from anaplastic carcinomas. About one third of tic-1 positive lymphomas lack the leukocyte common antigen (CD45).ImmunogenRecombinant human CD30/TNFRSF8 protein (Uniprot: P28908)Product Application DetailsFlow Cytometry, Immunocytochemistry/ Immunofluorescence, ImmunofluorescenceRecommended DilutionsFlow Cytometry, Immunohistochemistry. Immunofluorescence, Immunofluorescence, Immunohistochemistry. Paraffin, LyTOF-ready, Immunofluorescence, Immunohistochemistry. Paraffin, Immunofluorescence	Gene ID	943
MarkerHodgkin & Reed-Sternberg Cell MarkerSpecificity/SensitivityRecognizes a single chain glycoprotein of 105/120kDa, identified as CD30/I CD30 is synthesized as a 90kDa precursor, which is processed in the Golgi complex into a membrane-bound phosphorylated mature 105/120kDa glycoprotein. In Hodgkins disease, CD30/Ki-1 antigen is expressed by mononuclear-Hodgkin and multinucleated Reed-Sternberg cells. It is also expressed by the tumor cells of a majority of anaplastic large cell lymphoma well as by a varying proportion of activated T and B cells. This monoclonal antibody distinguishes large cell lymphomas derived from activated lymphoi cells from histiocytic malignancies and lymphomas derived from resting and precursor lymphoid cells or from anaplastic carcinomas. About one third of tki-1 positive lymphomas lack the leukocyte common antigen (CD45).ImmunogenRecombinant human CD30/TNFRSF8 protein (Uniprot: P28908)Product Application DetailsFlow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, ImmunofluorescenceRecommended DilutionsFlow Cytometry, Immunohistochemistry, Immunofluorescence, Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence	Gene Symbol	TNFRSF8
Specificity/SensitivityRecognizes a single chain glycoprotein of 105/120kDa, identified as CD30/l CD30 is synthesized as a 90kDa precursor, which is processed in the Golgi complex into a membrane-bound phosphorylated mature 105/120kDa glycoprotein. In Hodgkins disease, CD30/Ki-1 antigen is expressed by mononuclear-Hodgkin and multinucleated Reed-Sternberg cells. It is also expressed by the tumor cells of a majority of anaplastic large cell lymphoma well as by a varying proportion of activated T and B cells. This monoclonal antibody distinguishes large cell lymphomas derived from resting and precursor lymphoid cells or from anaplastic carcinomas. About one third of t Ki-1 positive lymphomas lack the leukocyte common antigen (CD45).ImmunogenRecombinant human CD30/TNFRSF8 protein (Uniprot: P28908)Product Application DetailsFlow Cytometry, Immunocytochemistry/ Immunofluorescence, ImmunofluorescenceRecommended DilutionsFlow Cytometry, Immunohistochemistry, Immunofuorescence, Immunofluorescence, Immunofluorescence, Immunofluorescence	Species	Human
CD30 is synthesized as a 90kDa precursor, which is processed in the Golgi complex into a membrane-bound phosphorylated mature 105/120kDa glycoprotein. In Hodgkins disease, CD30/Ki-1 antigen is expressed by mononuclear-Hodgkin and multinucleated Reed-Sternberg cells. It is also expressed by the tumor cells of a majority of anaplastic large cell lymphoma well as by a varying proportion of activated T and B cells. This monoclonal antibody distinguishes large cell lymphomas derived from activated lympho cells from histiocytic malignancies and lymphomas derived from resting and precursor lymphoid cells or from anaplastic carcinomas. About one third of the Ki-1 positive lymphomas lack the leukocyte common antigen (CD45).ImmunogenRecombinant human CD30/TNFRSF8 protein (Uniprot: P28908)Product Application DetailsFlow Cytometry, Immunocytochemistry/ Immunofluorescence, ImmunofluorescenceRecommended DilutionsFlow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunofluorescence, Immunofluorescence, Immunofluorescence	Marker	Hodgkin & Reed-Sternberg Cell Marker
Product Application Details Applications Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence Recommended Dilutions Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence	Specificity/Sensitivity	glycoprotein. In Hodgkins disease, CD30/Ki-1 antigen is expressed by mononuclear-Hodgkin and multinucleated Reed-Sternberg cells. It is also expressed by the tumor cells of a majority of anaplastic large cell lymphomas as well as by a varying proportion of activated T and B cells. This monoclonal antibody distinguishes large cell lymphomas derived from activated lymphoid cells from histiocytic malignancies and lymphomas derived from resting and precursor lymphoid cells or from anaplastic carcinomas. About one third of the
ApplicationsFlow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, ImmunofluorescenceRecommended DilutionsFlow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence	Immunogen	Recombinant human CD30/TNFRSF8 protein (Uniprot: P28908)
Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence Recommended Dilutions Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence	Product Application Details	
Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence	Applications	Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready,
	Recommended Dilutions	Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence,
Application NotesOptimal dilution of this antibody should be experimentally determined.	Application Notes	Optimal dilution of this antibody should be experimentally determined.
Application NotesOptimal dilution of this antibody should be experimentally determined.		Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence, CyTOF-ready





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-47906F

NBP1-43319F	Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1) [FITC]
NBP2-22660	Recombinant Human CD30/TNFRSF8 His Protein
210-TA-005	TNF-alpha [Unconjugated]
813-CD-100	CD30/TNFRSF8 [Unconjugated]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-47906F

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

