Product Datasheet

CD30/TNFRSF8 Antibody (Ki-1/779) [DyLight 680] NBP2-47906FR

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-47906FR

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-47906FR



NBP2-47906FR

CD30/TNFRSF8 Antibody (Ki-1/779) [DyLight 680]

Unit Size 0.1 ml Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark. Clonality Monoclonal Clonality Monoclonal Clona Kl-1779 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate DyLight 680 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description This conjugate is made on demand. Actual recovery may vary from the stated on the datasheet. Host Mouse Gene D 943 Gene Symbol TNFRSF8 Species Human Marker Hodgkin & Reed-Sternberg Cell Marker Specificity/Sensitivity Recognizes a single chain glycoprotein of 105/120kDa, identified as CD30/Ki-1. CD3 is synthesized as a 90kDa precursor, which is processed in the Golgi complexit not a membrane-bound phosphorytated mature 105/210kDa specificity/Sensitivity Recognizes a single chain glycoprotein of 105/120kDa, identified as CD30/Ki-1. Kreir Hodgkin & Reed-Sternberg Cell Marker <tr< th=""><th>Product Information</th><th></th></tr<>	Product Information	
services. Storage Store at 4C in the dark. Clonality Monoclonal Clone Ki-1/779 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate DyLight 680 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 943 Gene Symbol TNFRSF8 Species Human Marker Hodgkin & Reed-Sternberg Cell Marker Species Human Marker Hodgkin a membrane-bound phosphorylated mature 105/120kDa, identified as CD30/Ki-1. Specificity/Sensitivity Recognizes a single chain glycoprotein of 105/120kDa, identified as CD30/Ki-1. Specificity/Sensitivity Recognizes a single chain glycoprotein on 105/120kDa, identified as CD30/Ki-1. Specificity/Sensitivity Recognizes a single chain glycoprotein on 105/120kDa, identified as CD30/Ki-1. Specificity/Sensitivity Recombinant humor Cells of a majority of anaplastic large cell hymphomas as well as b	Unit Size	0.1 ml
ClonalityMonoclonalCloneKi-1/779Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateDyLight 680PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionThis conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.HostMouseGene ID943Gene SymbolTNFRSF8SpeciesHumanMarkerHodgkin & Reed-Sternberg Cell MarkerSpecificity/SensitivityRecognizes a single chain glycoprotein of 105/120kDa, identified as CD30/Ki-1. CD30 is synthesized as a 90kDa precursor, which is repressed by monouclear-Hodgkin and multinucleated Reed-Sternberg cell sing and antibody distinguishes large cell lymphomas derived from activated if monolater antibody distinguishes large cell lymphomas derived from activated if monolater antibody distinguishes large cell lymphomas derived from resting and antibody distinguishes large cell lymphomas derived from activated if monolater antibody distinguishes large cell lymphomas derived from resting and antibody distinguishes large cell lymphomas derived from activated if monolocal antibody distinguishes large cell lymphomas derived from activated if monolater antibody distinguishes large cell lymphomas derived from activated if monolocal antibody distinguishes large cell lymphomas derived from activated if monolocal antibody distinguishes large cell lymphomas derived from activated if monolocal antibody distinguishes large cell lymphomas derived form activated if monolocal antibody distinguishes large cell lymphomas derived from activated if mo	Concentration	•
Clone Ki-1/779 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate DyLight 680 Purity Protein A or G purified Buffer Sodium Borate Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 943 Gene Symbol TNFRSF8 Species Human Marker Hodgkin & Reed-Sternberg Cell Marker Specificity/Sensitivity Recognizes a single chain glycoprotein of 105/120kDa, identified as CD30/Ki-1. CD30 is synthesized as a 90kDa precursor, which is processed in the Golgi complex into a membrane-bound phosphonylated mature 105/120kDa glycoprotein. In Hodgkins disease, CD30/Ki-1 antigen is expressed by grononcuclear-Hodgkin and multinucleated Reed-Sternberg Cell Marker Specificity/Sensitivity Recognizes a single chain glycoprotein of 105/120kDa, identified as CD30/Ki-1. CD30 is synthesized as a 90kDa precursor, which is processed in the Golgi complex into a membrane-bound phosphonylated mature 105/120kDa glycoprotein. In Hodgkins disease, CD30/Ki-1 antigen is expressed by monocuclear-Hodgkin and multinucleated Reed-Sternberg Cells. It is also expressed by the tumor cells of a majority of anaplastic large cell lymphomas as expressed by form anaplastic large cells more sing and antibody distinguishes lar	Storage	Store at 4C in the dark.
Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate DyLight 680 Purity Protein A or G purified Buffer 60mM Sodium Borate Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 943 Gene Symbol TNFRSF8 Species Human Marker Hodgkin & Reed-Sternberg Cell Marker Specificity/Sensitivity Recognizes a single chain glycoprotein of 105/120kDa, identified as CD30/Ki-1. CD30 is synthesized as a 90kDa precursor, which is processed in the Golgi complex into a membrane-bound phosphorylated mature 105/120kDa, identified as CD30/Ki-1. Specificity/Sensitivity Recognizes a single chain glycoprotein of 105/120kDa, identified as CD30/Ki-1. Solg is synthesized as a 90kDa precursor, which is processed by mononuclear-Hodgkin and multinucleated Reed-Sternberg cells. It is also expressed by well as by a varying proportion of activated from resting and precursor lymphomas cells of rom nanplastic carrinomas. About one third of the ki-1 positive lymphomas lack the leukocyte common antigen (CD45). Immunogen Recombinant human CD30/TNFRSF8 protein (Uniprot: P28908) <	Clonality	Monoclonal
Isotype IgG1 Kappa Conjugate DyLight 680 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 943 Gene Symbol TNFRSF8 Species Human Marker Hodgkin & Reed-Sternberg Cell Marker Specificity/Sensitivity Recognizes a single chain glycoprotein of 105/120kDa, identified as CD30/Ki-1. Specificity/Sensitivity Recognizes a single chain glycoprotein of 105/120kDa, identified as CD30/Ki-1. Specificity/Sensitivity Recognizes a single chain glycoprotein of 105/120kDa, identified as CD30/Ki-1. Specificity/Sensitivity Recognizes a single chain glycoprotein of 105/120kDa, identified as CD30/Ki-1. Specificity/Sensitivity Recognizes a single chain glycoprotein of 105/120kDa, identified as CD30/Ki-1. Specificity/Sensitivity Recognizes a single chain glycoprotein of 105/120kDa, identified as CD30/Ki-1. Specificity/Sensitivity Recognizes ad single chain glycoprotein of 105/120kDa, identified as CD30/Ki-1. Specificity/Sensitivity Recognizes ad single	Clone	Ki-1/779
Conjugate DyLight 680 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 943 Gene Symbol TNFRSF8 Species Human Marker Hodgkin & Reed-Sternberg Cell Marker Specificity/Sensitivity Recognizes a single chain glycoprotein of 105/120kDa, identified as CD30/Ki-1. CD30 is synthesized as a 90kDa precursor, which is processed by mononuclear-Hodgkin and multinucleated Reed-Sternberg cells. It is also expressed by the tumor cells of a majority of anaplastic large cell lymphomas as well as by a varying proportion of activated T and B cells. This monoclonal antibody distinguishes large cell lymphomas derived from resting and precursor lymphoid cells or from anaplastic carcinomas. About one third of the Ki-1 positive lymphomas text the leukocyte common antigen (CD45). Immunogen Recombinant human CD30/TNFRSF8 protein (Uniprot: P28908) Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries. Product Application Details Flow Cytometry,	Preservative	0.05% Sodium Azide
Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 943 Gene Symbol TNFRSF8 Species Human Marker Hodgkin & Reed-Sternberg Cell Marker Specificity/Sensitivity Recognizes a single chain glycoprotein of 105/120kDa, identified as CD30/Ki-1. CD30 is synthesized as a 90kDa precursor, which is processed in the Golgi complex into a membrane-bound phosphorylated mature 106/120kDa, glycoprotein. In Hodgkins disease, CD30/Ki-1 antigen is expressed by mononuclear-Hodgkin and multinucleated Reed-Sternberg cells. It is also expressed by the tumor cells of a majority of anaplastic targe cell lymphomas as well as by a varying proportion of activated T and B cells. This monoclonal antibody distinguishes large cell lymphomas derived from resting and precursor lymphoid cells or from anaplastic carcionmas. About one third of the K-1 positive lymphomas tack the leukocyte common antigen (CD45). Immunogen Recombinant human CD30/TNFRSF8 protein (Uniprot: P28908) Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries. Product Application Details Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunonfluorescence, Immunohistochemistry-Paraffin, CyTOF-ready, Immuno	Isotype	IgG1 Kappa
Buffer 50mM Sodium Borate Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 943 Gene Symbol TNFRSF8 Species Human Marker Hodgkin & Reed-Sternberg Cell Marker Specificity/Sensitivity Recognizes a single chain glycoprotein of 105/120kDa, identified as CD30/Ki-1. CD30 is synthesized as a 90kDa precursor, which is processed in the Golgi complex into a membrane-bound phosphorylated mature 105/120kDa glycoprotein. In Hodgkins disease, CD30/Ki-1 antigen is expressed by mononuclear-Hodgkin and multinucleated Reed-Sternberg cells. It is also expressed by the tumor cells of a majority of anaplastic large cell lymphomas as well as by a varying proportion of a crivated T and B cells. This monoclonal antibody distinguishes large cell lymphomas derived from activated lymphoid cells from histiocytic malignancies and lymphomas About one third of the Ki-1 positive lymphoid acles of from anaplastic carcinomas. About one third of the Ki-1 positive lymphomas lack the leukocyte common antigen (CD45). Immunogen Flow Cytometry, Immunocytochemistry-Immunofluorescence, Immunofluorescence Product Application Details Flow Cytometry, Immunocytochemistry-Paraffin, CyTOF-ready, Immunofluorescence, CyTOF-ready	Conjugate	DyLight 680
Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 943 Gene Symbol TNFRSF8 Species Human Marker Hodgkin & Reed-Sternberg Cell Marker Specificity/Sensitivity Recognizes a single chain glycoprotein of 105/120kDa, identified as CD30/Ki-1. CD30 is synthesized as a 90kDa precursor, which is processed in the Golgi complex into a membrane-bound phosphorylated mature 105/120kDa glycoprotein. In Hodgkins disease, CD30/Ki-1 antigen is expressed by mononuclear-Hodgkin and multinucleated Reed-Sternberg cells. It is also expressed by the tumor cells of a majority of anaplastic large cell lymphomas as well as by a varying proportion of activated T and B cells. This monoclonal antibody distinguishes large cell lymphomas derived from activated lymphoid cells for minisquancies and lymphomas derived from resting and precursor lymphoid cells or from anaplastic carcinomas. About one third of the Ki-1 positive lymphomas lack the leukocyte common antigen (CD45). Immunogen Recombinant human CD30/TNFRSF8 protein (Uniprot: P28908) Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries. Product Application Details Flow Cytometry, Immunohistochemistry/Immunofluorescence, Immunohistochemistry. Paraffin, CyTOF-ready, Immunofluorescence, CyTOF-ready.	Purity	Protein A or G purified
DescriptionThis conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.HostMouseGene ID943Gene SymbolTNFRSF8SpeciesHumanMarkerHodgkin & Reed-Stemberg Cell MarkerSpecificity/SensitivityRecognizes a single chain glycoprotein of 105/120kDa, identified as CD30/Ki-1. CD30 is synthesized as a 90kDa precursor, which is processed in the Golgi complex into a membrane-bound phosphorylated mature 105/120kDa glycoprotein. In Hodgkins disease, CD30/Ki-1 antigen is expressed by mononuclear-Hodgkin and multinucleated Reed-Stemberg cells. It is also expressed by the tumor cells of a majority of anaplastic large cell lymphomas as well as by a varying proportion of activated T and B cells. This monoclonal antibody distinguishes large cell lymphomas derived from resting and precursor lymphoid cells or from anaplastic carcinomas. About one third of the Ki-1 positive lymphomas lack the leukocyte common antigen (CD45).ImmunogenRecombinant human CD30/TNFRSF8 protein (Uniprot: P28908)NotesDyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.Product Application DetailsFlow Cytometry, Immunohistochemistry/Immunofluorescence, ImmunofluorescenceRecommended DilutionsFlow Cytometry, Immunohistochemistry, Immunofluorescence, CyTOF-ready.	Buffer	50mM Sodium Borate
Volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.HostMouseGene ID943Gene SymbolTNFRSF8SpeciesHumanMarkerHodgkin & Reed-Sternberg Cell MarkerSpecificity/SensitivityRecognizes a single chain glycoprotein of 105/120kDa, identified as CD30/Ki-1. CD30 is synthesized as a 90kDa precursor, which is processed in the Golgi complex into a membrane-bound phosphorylated mature 105/120kDa glycoprotein. In Hodgkins disease, CD30/Ki-1 antigen is expressed by mononuclear-Hodgkin and multinucleated Reed-Sternberg cells. It is also expressed by the tumor cells of a majority of anaplastic large cell lymphomas as well as by a varying proportion of activated T and B cells. This monoclonal antibody distinguishes large cell lymphomas derived from esting and precursor lymphoid cells or from anaplastic carcinomas. About one third of the Ki-1 positive lymphomas lack the leukocyte common antigen (CD45).ImmunogenRecombinant human CD30/TNFRSF8 protein (Uniprot: P28908)NotesDyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.Product Application DetailsFlow Cytometry, Immunohistochemistry/Immunofluorescence, ImmunofluorescenceRecommended DilutionsFlow Cytometry, Immunohistochemistry-Paraffin, Immunofluorescence, CyTOF-ready.	Product Description	
Gene ID 943 Gene Symbol TNFRSF8 Species Human Marker Hodgkin & Reed-Sternberg Cell Marker Specificity/Sensitivity Recognizes a single chain glycoprotein of 105/120kDa, identified as CD30/Ki-1. CD30 is synthesized as a 90kDa precursor, which is processed in the Golgi complex into a membrane-bound phosphorylated mature 105/120kDa glycoprotein. In Hodgkins disease, CD30/Ki-1 antigen is expressed by mononuclear-Hodgkin and multinucleated Reed-Sternberg cells. It is also expressed by the tumor cells of a majority of anaplastic large cell lymphomas as well as by a varying proportion of activated 1 and B cells. This monoclonal antibody distinguishes large cell lymphomas derived from activated lymphoid cells from histiccytic malignancies and lymphomas derived from resting and precursor lymphoid cells or from anaplastic carcinomas. About one third of the Ki-1 positive lymphomas lack the leukocyte common antigen (CD45). Immunogen Recombinant human CD30/TNFRSF8 protein (Uniprot: P28908) Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries. Product Application Details Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunofluorescence Recommended Dilutions Flow Cytometry, Immunohistochemistry.Paraffin, CyTOF-ready, Immunofluorescence, Immunohistochemistry.Paraffin, Immunofluorescence, CyTOF-ready	Description	volume of this product. The volume will be greater than or equal to the unit size
Gene SymbolTNFRSF8SpeciesHumanMarkerHodgkin & Reed-Sternberg Cell MarkerSpecificity/SensitivityRecognizes a single chain glycoprotein of 105/120kDa, identified as CD30/Ki-1. CD30 is synthesized as a 90kDa precursor, which is processed in the Golgi complex into a membrane-bound phosphorylated mature 105/120kDa glycoprotein. In Hodgkins disease, CD30/Ki-1 antigen is expressed by mononuclear-Hodgkin and multinucleated Reed-Sternberg cells. It is also expressed by the tumor cells of a majority of anaplastic large cell lymphomas as well as by a varying proportion of activated T and B cells. This monoclonal antibody distinguishes large cell lymphomas derived from resting and precursor lymphoid cells or from anaplastic carcinomas. About one third of the Ki-1 positive lymphomas lack the leukocyte common antigen (CD45).ImmunogenRecombinant human CD30/TNFRSF8 protein (Uniprot: P28908)NotesDyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.Product Application DetailsFlow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunofluorescence, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence, CyTOF-ready	Host	Mouse
SpeciesHumanMarkerHodgkin & Reed-Sternberg Cell MarkerSpecificity/SensitivityRecognizes a single chain glycoprotein of 105/120kDa, identified as CD30/Ki-1. CD30 is synthesized as a 90kDa precursor, which is processed in the Golgi complex into a membrane-bound phosphorylated mature 105/120kDa glycoprotein. In Hodgkins disease, CD30/Ki-1 antigen is expressed by mononuclear-Hodgkin and multinucleated Reed-Sternberg cells. It is also expressed by the tumor cells of a majority of anaplastic large cell lymphomas as well as by a varying proportion of activated T and B cells. This monoclonal 	Gene ID	943
MarkerHodgkin & Reed-Sternberg Cell MarkerSpecificity/SensitivityRecognizes a single chain glycoprotein of 105/120kDa, identified as CD30/Ki-1. CD30 is synthesized as a 90kDa precursor, which is processed in the Golgi complex into a membrane-bound phosphorylated mature 105/120kDa glycoprotein. In Hodgkins disease, CD30/Ki-1 antigen is expressed by mononuclear-Hodgkin and multinucleated Reed-Sternberg cells. It is also expressed by the tumor cells of a majority of anaplastic large cell lymphomas as well as by a varying proportion of activated T and B cells. This monoclonal antibody distinguishes large cell lymphomas derived from activated lymphoid cells from histiocytic malignancies and lymphomas derived from resting and precursor lymphoid cells or from anaplastic carcinomas. About one third of the Ki-1 positive lymphomas lack the leukocyte common antigen (CD45).ImmunogenRecombinant human CD30/TNFRSF8 protein (Uniprot: P28908)NotesDyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.Product Application DetailsFlow Cytometry, Immunocytochemistry/ Immunofluorescence, ImmunofluorescenceRecommended DilutionsFlow Cytometry, Immunohistochemistry. Immunofluorescence, Immunofluorescence, Immunohistochemistry. Paraffin, Immunofluorescence, CyTOF-ready	Gene Symbol	TNFRSF8
Specificity/SensitivityRecognizes a single chain glycoprotein of 105/120kDa, identified as CD30/Ki-1. CD30 is synthesized as a 90kDa precursor, which is processed in the Golgi complex into a membrane-bound phosphorylated mature 105/120kDa glycoprotein. In Hodgkins disease, CD30/Ki-1 antigen is expressed by mononuclear-Hodgkin and multinucleated Reed-Sternberg cells. It is also expressed by the tumor cells of a majority of anaplastic large cell lymphomas as well as by a varying proportion of activated T and B cells. This monoclonal antibody distinguishes large cell lymphomas derived from activated lymphoid cells from histiocytic malignancies and lymphomas derived from resting and precursor lymphoid cells or from anaplastic carcinomas. About one third of the Ki-1 positive lymphomas lack the leukocyte common antigen (CD45).ImmunogenRecombinant human CD30/TNFRSF8 protein (Uniprot: P28908)NotesDyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.Product Application DetailsFlow Cytometry, Immunocytochemistry/ Immunofluorescence, ImmunofluorescenceRecommended DilutionsFlow Cytometry, Immunohistochemistry, Immunofluorescence, CyTOF-ready	Species	Human
CD30 is synthesized as a 90kDa precursor, which is processed in the Golgi complex into a membrane-bound phosphorylated mature 105/120kDa glycoprotein. In Hodgkins disease, CD30/Ki-1 antigen is expressed by mononuclear-Hodgkin and multinucleated Reed-Sternberg cells. It is also expressed by the tumor cells of a majority of anaplastic large cell lymphomas as well as by a varying proportion of activated T and B cells. This monoclonal antibody distinguishes large cell lymphomas derived from activated lymphoid cells from histiocytic malignancies and lymphomas derived from resting and precursor lymphoid cells or from anaplastic carcinomas. About one third of the Ki-1 positive lymphomas lack the leukocyte common antigen (CD45).ImmunogenRecombinant human CD30/TNFRSF8 protein (Uniprot: P28908)NotesDyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.Product Application DetailsFlow Cytometry, Immunocytochemistry/ Immunofluorescence, ImmunofluorescenceRecommended DilutionsFlow Cytometry, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence, RyTOF-ready	Marker	Hodgkin & Reed-Sternberg Cell Marker
Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries. Product Application Details Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence Recommended Dilutions Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, CyTOF-ready	Specificity/Sensitivity	CD30 is synthesized as a 90kDa precursor, which is processed in the Golgi complex into a membrane-bound phosphorylated mature 105/120kDa glycoprotein. In Hodgkins disease, CD30/Ki-1 antigen is expressed by mononuclear-Hodgkin and multinucleated Reed-Sternberg cells. It is also expressed by the tumor cells of a majority of anaplastic large cell lymphomas as well as by a varying proportion of activated T and B cells. This monoclonal antibody distinguishes large cell lymphomas derived from activated lymphoid cells from histiocytic malignancies and lymphomas derived from resting and precursor lymphoid cells or from anaplastic carcinomas. About one third of the
DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries. Product Application Details Applications Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence Recommended Dilutions Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, CyTOF-ready	Immunogen	Recombinant human CD30/TNFRSF8 protein (Uniprot: P28908)
ApplicationsFlow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, ImmunofluorescenceRecommended DilutionsFlow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence, CyTOF-ready	Notes	DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.
Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence Recommended Dilutions Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence, CyTOF-ready	Product Application Details	
Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence, CyTOF-ready	Applications	Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready,
Application NotesOptimal dilution of this antibody should be experimentally determined.	Recommended Dilutions	Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence,
	Application Notes	Optimal dilution of this antibody should be experimentally determined.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-47906FR

NBP1-43319FR-0.5ml	Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1) [DyLight 680]
NBP2-22660	Recombinant Human CD30/TNFRSF8 His Protein
210-TA-005	TNF-alpha [Unconjugated]
813-CD-100	CD30/TNFRSF8 [Unconjugated]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-47906FR

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

