Product Datasheet

Cytokeratin 7 Antibody (KRT7/1198) - Azide and BSA Free NBP2-47944-0.1mg

Unit Size: 0.1 mg

Store at -20 to -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-47944

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-47944



NBP2-47944-0.1mg


Cytokeratin 7 Antibody (KRT7/1198) - Azide and BSA Free

as cytokeratin 7. This monoclonal antibody is highly specific to cytokeratin 7 and shows no cross-reaction with other IFPs. Cytokeratin 7 is a basic cytokeratin, which is found in most glandular and transitional epithelia but not in the stratified squamous epithelia. Keratin 7 is expressed in the epithelial cells of ovary, lung, and breast but not of colon, prostate, or gastrointestinal tract. This monoclonal antibody is highly useful in distinguishing ovarian carcinomas (keratin 7+) from colon carcinomas (keratin 7-). Immunogen Recombinant full-length human Cytokeratin 7 protein (Uniprot: P08729) Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence Western Blot, Flow Cytometry 0.5 - 1 ug/million cells in 0.1 ml, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin 0.5 - 1.0 ug/ml, Immunofluorescence 0.5 - 1.0 ug/ml, CyTOF-ready Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining	Cytokeratin 7 Antibody (KRT7/1198) - Azide and BSA Free	
Storage Store at -20 to -80C. Avoid freeze-thaw cycles.	Product Information	
Storage Stora at -20 to -80C. Avoid freeze-thaw cycles.	Unit Size	0.1 mg
Clone KRT7/1198 Preservative No Preservative Isotype IgG1 Kappa Purity Protein A or G purified Buffer 10 mM PBS Target Molecular Weight 55 kDa Product Description 1.0 mg/ml of antibody purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WITH BSA & azide (NBP2-44820). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Mouse Gene ID 3855 Gene Symbol KRT7 Species Human, Rat Marker Glandular and Transitional Epithelial Marker Specificity/Sensitivity It recognizes an intermediate filament protein (IFP) of 55kDa, which is identified as cytokeratin 7. This monoclonal antibody is highly specific to cytokeratin 7 and shows no cross-reaction with other IFPs. Cytokeratin 7 is a basic cytokeratin which is found in most glandular and transitional epithelia but not in the stratified squamous epithelia. Keratin 7 is expressed in the epithelial cells of ovary, lung, and breast but not of colon, prostate, or gastrointestinal transitional epithelia but not in the stratified squamous epithelia. Keratin 7 is expressed in the epithelial cells of ovary, lung, and breast but not of colon, prostate, or gastrointestinal transitional epithelial cells of ovary, lung, and breast but not of colon, prostate, or gastrointestinal transitorional antibody is highly useful in distinguishing ovarian carcinomas (keratin 7+) from colon carcinomas (keratin 7-) is expressed in the epithelial cells of ovary, lung, and breast but not of colon, prostate, or gastrointestinal transitional epithelia but not in the stratifies squamous epithelia. Keratin 7 is expressed in the epithelial cells of ovary, lung, and breast but not of colon, prostate, or gastrointestinal transitional epithelia but not in the stratifier squamous epithelia. Keratin 7-) is expressed in the epithelial cells of ovary, lung, and breast but not of colon, prostate, or gastrointestinal transitional epithelia but not in the stratif	Concentration	1.0 mg/ml
Clone KRT7/1198 Preservative No Preservative IgG1 Kappa Purity Protein A or G purified Buffer 10 mM PBS Target Molecular Weight 55 kDa Product Description Description 1.0 mg/ml of antibody purified from Bioreactor Concentrate by Protein A/G, Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WITH BSA & azide (NBP2-44820). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Mouse Gene ID 3855 Gene Symbol KRT7 Species Human, Rat Marker Glandular and Transitional Epithelial Marker Specificity/Sensitivity It recognizes an intermediate filament protein (IFP) of 55kDa, which is identified as cytokeratin 7. This monoclonal antibody is highly specific to cytokeratin 7 and shows no cross-reaction with other IFPs. Cytokeratin 7 is a basic cytokeratin which is found in most glandular and transitional epithelial but not in the stratified squamous epithelia. Keratin 7 is expressed in the epithelial cells of ovary, lung, and breast but not of colon, prostate, or gastrointestinal tract. This monoclonal antibody is highly useful in distinguishing ovarian carcinomas (keratin 7+) from colon carcinomas (keratin 7 is expressed in the epithelial cells of ovary, lung, and breast but not of colon, prostate, or gastrointestinal tract. This monoclonal antibody is highly useful in distinguishing ovarian carcinomas (keratin 7+) from colon carcinomas (keratin 7). Immunogen Recombinant full-length human Cytokeratin 7 protein (Uniprot: P08729) Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Paraffin, CyTOF-ready, Immunofluorescence, Immunohistochemistry (Formalin-fixed): 1-2ug/mll for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 950 Cololing at RT for 20 minutes.	Storage	Store at -20 to -80C. Avoid freeze-thaw cycles.
Preservative No Preservative Isotype IgG1 Kappa Purity Protein A or G purified Buffer 10 mM PBS Target Molecular Weight 55 kDa Product Description Description 1.0 mg/ml of antibody purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WITH BSA & azide (NBP2-44820). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Mouse Gene ID 3855 Gene Symbol KRT7 Species Human, Rat Marker Glandular and Transitional Epithelial Marker Specificity/Sensitivity It recognizes an intermediate filament protein (IFP) of 55kDa, which is identified as cytokeratin 7. This monoclonal antibody is highly specific to cytokeratin 7 and shows no cross-reaction with other IFPs. Cytokeratin 7 is a basic cytokeratin, which is found in most glandular and transitional epithelia belated legisle of vary, lung, and breast but not of colon, prostate, or gastrointestinal tract. This monoclonal antibody is highly useful in distinguishing ovarian carcinomas (Keratin 7+) from colon carcinomas (Keratin 7-). Immunogen Recombinant full-length human Cytokeratin 7 protein (Uniprot: P08729) Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry Paraffin, CyTOF-ready, Immunofluorescence (Immunohistochemistry-Paraffin 0.5 - 1.0 ug/ml, Immunofluorescence, Immunohistochemistry, Immunocytochemistry Immunofluorescence 0.5 - 1.0 ug/ml, CyTOF-ready Application Notes Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minu	Clonality	Monoclonal
Isotype	Clone	KRT7/1198
Purity Protein A or G purified Buffer 10 mM PBS Target Molecular Weight 55 kDa Product Description Description 1.0 mg/ml of antibody purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WITH BSA & azide (NBP2-44820). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Mouse Gene ID 3855 Gene Symbol KRT7 Species Human, Rat Marker Glandular and Transitional Epithelial Marker Specificity/Sensitivity It recognizes an intermediate filament protein (IFP) of 55kDa, which is identified as cytokeratin 7. This monoclonal antibody is highly specific to cytokeratin 7 and shows no cross-reaction with other IFPs. Cytokeratin 7 is a basic cytokeratin, which is found in most glandular and transitional epithelial unto in the stratified squamous epithelial. Keratin 7 is expressed in the epithelial cells of ovary, lung, and breast but not of colon, prostate, or gastrointestinal tract. This monoclonal antibody is highly useful in distinguishing ovarian carcinomas (keratin 7+) from colon carcinomas (keratin 7-). Immunogen Recombinant full-length human Cytokeratin 7 protein (Uniprot: P08729) Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin 0.5 - 1.0 ug/mill in cells in 0.1 ml, Immunohistochemistry Paraffin 0.5 - 1.0 ug/mill in cells in 0.1 ml, Immunohistochemistry (Formalin-fixed): 1-2ug/mil for 30 minutes at RT. Staining of formalin-fixed itssues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Preservative	No Preservative
Buffer 10 mM PBS	Isotype	IgG1 Kappa
Target Molecular Weight Product Description Description 1.0 mg/ml of antibody purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WITH BSA & azide (NBP2-44820). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Mouse Gene ID 3855 Gene Symbol KRT7 Species Human, Rat Marker Glandular and Transitional Epithelial Marker Specificity/Sensitivity It recognizes an intermediate filament protein (IFP) of 55kDa, which is identified as cytokeratin 7. This monoclonal antibody is highly specific to cytokeratin 7 and shows no cross-reaction with other IFPs. Cytokeratin 7 is a basic cytokeratin, which is found in most glandular and transitional epithelial but not in the stratified squamous epithelia. Keratin 7 is expressed in the epithelial cells of ovary, lung, and breast but not of colon, prostate, or gastrointestinal trad. This monoclonal antibody is highly useful in distinguishing ovarian carcinomas (keratin 7+) from colon carcinomas (keratin 7-). Immunogen Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunofluorescence, Immunohistochemistry, Immunofluorescence, Immunohistochemistry, Immunofluorescence, Immunohistochemistry, Immunofluorescence, Immunohistochemistry, Immunofluorescence, Immunohistochemistry-Paraffin 0.5 - 1.0 ug/ml, Immunofluorescence 0.5 - 1.0 ug/ml, CyTOF-ready, Immunofluorescence 0.5 - 1.0 ug/ml, CyTOF-ready Immunofluorescence 0.5 - 1.0 ug/ml, CyTOF-up-dy Immunofluorescence 0.5 - 1.0 ug/ml, CyTOF-up-dy Immunof	Purity	Protein A or G purified
Product Description 1.0 mg/ml of antibody purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WITH BSA & azide (NBP2-44820). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Mouse Gene ID 3855 Gene Symbol KRT7 Species Human, Rat Marker Glandular and Transitional Epithelial Marker Specificity/Sensitivity It recognizes an intermediate filament protein (IFP) of 55kDa, which is identified as cytokeratin 7. This monoclonal antibody is highly specific to cytokeratin 7 ashows no cross-reaction with other IFPs. Cytokeratin 7 is a basic cytokeratin, which is found in most glandular and transitional epithelia Lote for ovary, lung, and breast but not of colon, prostate, or gastrointestinal tract. This monoclonal antibody is highly useful in distinguishing ovarian carcinomas (keratin 7+) from colon carcinomas (keratin 7-) from colon carcin	Buffer	10 mM PBS
Description	Target Molecular Weight	55 kDa
Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WITH BSA & azide (NBP2-44820). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Mouse Gene ID 3855 Gene Symbol KRT7 Species Human, Rat Marker Glandular and Transitional Epithelial Marker Specificity/Sensitivity It recognizes an intermediate filament protein (IFP) of 55kDa, which is identified as cytokeratin 7. This monoclonal antibody is highly specific to cytokeratin 7 and shows no cross-reaction with other IFPs. Cytokeratin 7 is a basic cytokeratin, which is found in most glandular and transitional epithelia but not in the stratified squamous epithelia. Keratin 7 is expressed in the epithelial cells of ovary, lung, and breast but not of colon, prostate, or gastrointestinal tract. This monoclonal antibody is highly useful in distinguishing ovarian carcinomas (keratin 7+) from colon carcinomas (keratin 7-). Immunogen Recombinant full-length human Cytokeratin 7 protein (Uniprot: P08729) Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin 0.5 - 1.0 ug/ml, Immunofluorescence 0.5 - 1.0 ug/ml, CyTOF-ready Application Notes Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissues sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Product Description	
Gene ID 3855 Gene Symbol KRT7 Species Human, Rat Marker Glandular and Transitional Epithelial Marker Specificity/Sensitivity It recognizes an intermediate filament protein (IFP) of 55kDa, which is identified as cytokeratin 7. This monoclonal antibody is highly specific to cytokeratin 7 and shows no cross-reaction with other IFPs. Cytokeratin 7 is a basic cytokeratin, which is found in most glandular and transitional epithelia but not in the stratified squamous epithelia. Keratin 7 is expressed in the epithelial cells of ovary, lung, and breast but not of colon, prostate, or gastrointestinal tract. This monoclonal antibody is highly useful in distinguishing ovarian carcinomas (keratin 7+) from colon carcinomas (keratin 7-). Immunogen Recombinant full-length human Cytokeratin 7 protein (Uniprot: P08729) Product Application Details Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Immunohistochemistry, Immunocytochemistry Immunofluorescence, Immunohistochemistry, Immunocytochemistry Immunofluorescence, Immunohistochemistry-Paraffin 0.5 - 1.0 ug/ml, Immunofluorescence 0.5 - 1.0 ug/ml, CyTOF-ready Application Notes Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Description	Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WITH BSA & azide (NBP2-44820). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to
Gene Symbol KRT7 Species Human, Rat Marker Glandular and Transitional Epithelial Marker Specificity/Sensitivity It recognizes an intermediate filament protein (IFP) of 55kDa, which is identified as cytokeratin 7. This monoclonal antibody is highly specific to cytokeratin 7 and shows no cross-reaction with other IFPs. Cytokeratin 7 is a basic cytokeratin, which is found in most glandular and transitional epithelia but not in the stratified squamous epithelia. Keratin 7 is expressed in the epithelial cells of ovary, lung, and breast but not of colon, prostate, or gastrointestinal tract. This monoclonal antibody is highly useful in distinguishing ovarian carcinomas (keratin 7+) from colon carcinomas (keratin 7-). Immunogen Recombinant full-length human Cytokeratin 7 protein (Uniprot: P08729) Product Application Details Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunofluorescence Recommended Dilutions Western Blot, Flow Cytometry 0.5 - 1 ug/million cells in 0.1 ml, Immunofluorescence, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin 0.5 - 1.0 ug/ml, Immunofluorescence 0.5 - 1.0 ug/ml, CyTOF-ready Application Notes Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Host	Mouse
Species	Gene ID	3855
Marker Glandular and Transitional Epithelial Marker Specificity/Sensitivity It recognizes an intermediate filament protein (IFP) of 55kDa, which is identified as cytokeratin 7. This monoclonal antibody is highly specific to cytokeratin 7 and shows no cross-reaction with other IFPs. Cytokeratin 7 is a basic cytokeratin, which is found in most glandular and transitional epithelia but not in the stratified squamous epithelia. Keratin 7 is expressed in the epithelial cells of ovary, lung, and breast but not of colon, prostate, or gastrointestinal tract. This monoclonal antibody is highly useful in distinguishing ovarian carcinomas (keratin 7+) from colon carcinomas (keratin 7-). Immunogen Recombinant full-length human Cytokeratin 7 protein (Uniprot: P08729) Product Application Details Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunofluorescence Recommended Dilutions Western Blot, Flow Cytometry 0.5 - 1 ug/million cells in 0.1 ml, Immunofistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin 0.5 - 1.0 ug/ml, Immunofluorescence 0.5 - 1.0 ug/ml, CyTOF-ready Application Notes Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Gene Symbol	KRT7
It recognizes an intermediate filament protein (IFP) of 55kDa, which is identified as cytokeratin 7. This monoclonal antibody is highly specific to cytokeratin 7 and shows no cross-reaction with other IFPs. Cytokeratin 7 is a basic cytokeratin, which is found in most glandular and transitional epithelia but not in the stratified squamous epithelia. Keratin 7 is expressed in the epithelial cells of ovary, lung, and breast but not of colon, prostate, or gastrointestinal tract. This monoclonal antibody is highly useful in distinguishing ovarian carcinomas (keratin 7+) from colon carcinomas (keratin 7-). Immunogen Recombinant full-length human Cytokeratin 7 protein (Uniprot: P08729) Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence Western Blot, Flow Cytometry 0.5 - 1 ug/million cells in 0.1 ml, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Paraffin 0.5 - 1.0 ug/ml, Immunofluorescence 0.5 - 1.0 ug/ml, CyTOF-ready Application Notes Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Species	Human, Rat
as cytokeratin 7. This monoclonal antibody is highly specific to cytokeratin 7 and shows no cross-reaction with other IFPs. Cytokeratin 7 is a basic cytokeratin, which is found in most glandular and transitional epithelia but not in the stratified squamous epithelia. Keratin 7 is expressed in the epithelial cells of ovary, lung, and breast but not of colon, prostate, or gastrointestinal tract. This monoclonal antibody is highly useful in distinguishing ovarian carcinomas (keratin 7+) from colon carcinomas (keratin 7-). Immunogen Recombinant full-length human Cytokeratin 7 protein (Uniprot: P08729) Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence Western Blot, Flow Cytometry 0.5 - 1 ug/million cells in 0.1 ml, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin 0.5 - 1.0 ug/ml, Immunofluorescence 0.5 - 1.0 ug/ml, CyTOF-ready Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Marker	Glandular and Transitional Epithelial Marker
Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence Recommended Dilutions Western Blot, Flow Cytometry 0.5 - 1 ug/million cells in 0.1 ml, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin 0.5 - 1.0 ug/ml, Immunofluorescence 0.5 - 1.0 ug/ml, CyTOF-ready Application Notes Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Specificity/Sensitivity	which is found in most glandular and transitional epithelia but not in the stratified squamous epithelia. Keratin 7 is expressed in the epithelial cells of ovary, lung, and breast but not of colon, prostate, or gastrointestinal tract. This monoclonal antibody is highly useful in distinguishing ovarian carcinomas (keratin 7+) from
Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence Western Blot, Flow Cytometry 0.5 - 1 ug/million cells in 0.1 ml, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin 0.5 - 1.0 ug/ml, Immunofluorescence 0.5 - 1.0 ug/ml, CyTOF-ready Application Notes Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Immunogen	Recombinant full-length human Cytokeratin 7 protein (Uniprot: P08729)
Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence Recommended Dilutions Western Blot, Flow Cytometry 0.5 - 1 ug/million cells in 0.1 ml, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin 0.5 - 1.0 ug/ml, Immunofluorescence 0.5 - 1.0 ug/ml, CyTOF-ready Application Notes Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Product Application Details	
Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin 0.5 - 1.0 ug/ml, Immunofluorescence 0.5 - 1.0 ug/ml, CyTOF-ready Application Notes Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Applications	Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready,
of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Recommended Dilutions	Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin 0.5 - 1.0 ug/ml, Immunofluorescence 0.5 - 1.0
	Application Notes	




Images


Western Blot: Cytokeratin 7 Antibody (KRT7/1198) - Azide and BSA Free [NBP2-47944] - HeLa cell lysate using Cytokeratin 7 Mouse Monoclonal Antibody (KRT7/1198).




Immunocytochemistry/Immunofluorescence: Cytokeratin 7 Antibody (KRT7/1198) - Azide and BSA Free [NBP2-47944] - MeOH-fixed HeLa cells labeling KRT7 with KRT7 Mouse Monoclonal Antibody (KRT7/1198) followed by Goat anti-Mouse IgG-CF488 (Green). The nuclear counterstain is Redot.




Immunohistochemistry-Paraffin: Cytokeratin 7 Antibody (KRT7/1198) - Azide and BSA Free [NBP2-47944] - Rat Lung stained with Cytokeratin 7 Mouse Monoclonal Antibody (KRT7/1198).



Immunohistochemistry-Paraffin: Cytokeratin 7 Antibody (KRT7/1198) - Azide and BSA Free [NBP2-47944] - Human Lung Carcinoma stained with Cytokeratin 7 Mouse Monoclonal Antibody (KRT7/1198).



Immunohistochemistry-Paraffin: Cytokeratin 7 Antibody (KRT7/1198) - Azide and BSA Free [NBP2-47944] - Human Ovarian Carcinoma stained with Cytokeratin 7 Mouse Monoclonal Antibody (KRT7/1198).





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP2-47944-0.1mg

HAF007 Goat anti-Mouse IgG Secondary Antibody [HRP]

NB720-B Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]

NBP1-43319-0.5mg Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1)

H00003855-P01-10ug Recombinant Human Cytokeratin 7 GST (N-Term) Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-47944

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

