Product Datasheet

Cytokeratin 19 Antibody (KRT19/800) [FITC] NBP2-47946F

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com

G

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-47946F

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-47946F



NBP2-47946F

Cytokeratin 19 Antibody (KRT19/800) [FITC]

Unit Size 0.1 ml Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark. Clonality Monoclonal Clonality Monoclonal Clonality Monoclonal Clone KRT19/800 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate FITC Purity Protein A or G purified Buffer PBS Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 3880 Gene Symbol KRT19 Species Human, Mouse, Rat Marker Pancreatic Stem Cell Marker Specificity/Sensitivity Recognizes a protein of 40kDa, identified as cytokeratin-19 (CK19), which is expressed in sweat gland, marmary gland ductal and secretory cells, bile ducts, gradiencies in lump in nodes, peripheral blood, bone marrow and breast cancer. Immunogen Recognizes a protein of 40kDa, identified as cytokeratin 9 (OK19), which is expre	Product Information	
services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneKRT19/800Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateFITCPurityProtein A or G purifiedBufferPBSProduct DescriptionThis conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.HostMouseGene ID3880Gene SymbolKRT19SpecifiesHuman, Mouse, RatMarkerPancreatic Stem Cell MarkerSpecifies.Recognizes a protein of 40kDa, identified as cytokeratin-19 (CK19), which is expressed in sweat gland, marmary gland ductal and secretory cells, bile ducts, gastrointesinal tract, bladder urothelium, oral epithelial, esophagus, and eccervical epithelium. Anti-CK19 reacts with a wide variety of epithelial malignancies including adenocarcinomas of the colon, stomach, pancreas, biliary carcinomas are also labeled. Anti-CK19 is a useful amplication is the identification of thyroid carcinoma of the papillary type, although 50%-60% of folicular malignancies including adenocarcinomas of the colon, stomach, pancreas, biliary carcinomas are also labeled. Anti-CK19 is a useful marker for detection of tumor carcinomas are also labeled. Anti-CK19 is a useful amplication is the identification of thyroid carcinoma of the papillary type, although 50%-60% of folicular malignancies including adenocarcinomas of the colon, stomach, pancreas, biliary carcinomas are also labeled. Anti-CK19 is a useful marker for detection of tumor carcinomas are also labeled. Anti-CK19 is a useful marker for d	Unit Size	0.1 ml
ClonalityMonoclonalCloneKRT19/800Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateFITCPurityProtein A or G purifiedBufferPBSProduct DescriptionThis conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.HostMouseGene ID3880Gene SymbolKRT19SpeciesHuman, Mouse, RatMarkerPancreatic Stem Cell MarkerSpecificity/SensitivityRecognizes a protein of 40kDa, identified as cytokeratin-19 (CK19), which is expressed in sweat gland, mammary gland ductal and secretory cells, bile ducts, gastrointestinal tract, biader urothelium, oral epithelia, esophagus, and ectocervical epithelium. Anti-CK19 reacts with a wide variety of epithelial malignancies including adencaricinomas are also labeled. Anti-CK19 is a useful marker for detection of turyoid carcinoma of the papillary type, although 50%-60% of folicular carcinomas are also labeled. Anti-CK19 is a useful marker for detection of turyoid carcinoma of the papillary type, although 50%-60% of folicular carcinomas are also labeled. Anti-CK19 is a useful marker for detection of turyoid carcinoma of the papillary type, although 50%-60% of folicular carcinomas are also labeled. Anti-CK19 is a useful marker for detection of turnor cells in Jymph nodes, peripheral blood, bone marrow and breast cancer.ImmunogenWestern Blot, Flow Cytometry, Flow (Intracellular), Immunohistochemistry/ Immunohistochemistry/ Immunohistochemistry/ Immunohistochemistry/ Immunohistochemistry/ Immunohistochemistry/ Immunohistochemistry/ Immunohi	Concentration	•
CloneKRT19/800Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateFITCPurityProtein A or G purifiedBufferPBSProduct DescriptionThis conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.HostMouseGene ID3880Gene SymbolKRT19SpeciesHuman, Mouse, RatMarkerPancreatic Stem Cell MarkerSpecificity/SensitivityRecognizes a protein of 40kDa, identified as cytokeratin-19 (CK19), which is expressed in sweat gland, marmary gland ductal and secretory cells, bile ducts, gastrointestinal tract, bladder urothelium, oral epithelial malignancies including adenocarcinomas of the colon, stomach, pancreas, biliary tract, liver, and breast. Perhaps the most useful application is the identification of thyroid carcinomas of the papillary type, although 50%-60% of follicular carcinomas are also labeled. Anti-CK19 is a useful marker for detection of thyroid carcinomas of the papillary type, although 50%-60% of follicular carcinomas are also labeled. Anti-CK19 is a useful marker for detection of thyroid carcinoma of the papillary type, although 50%-60% of follicular carcinomas are also labeled. Anti-CK19 is a useful marker for detection of thyroid carcinoma of the papillary type, although 50%-60% of follicular carcinomas are also labeled. Anti-CK19 is a useful marker for detection of thyroid carcinomas of the colon, stomach, pancreas, biliary tract, liver, and breast. Perhaps the most useful application is the identification of thyroid carcinomas are also labeled. Anti-CK19 is a useful marker for detection	Storage	Store at 4C in the dark.
Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateFITCPurityProtein A or G purifiedBufferPBSProduct DescriptionThis conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.HostMouseGene ID3880Gene SymbolKRT19SpeciesHuman, Mouse, RatMarkerPancreatic Stem Cell MarkerSpecificity/SensitivityRecognizes a protein of 40kDa, identified as cytokeratin-19 (CK19), which is expressed in sweat gland, mammary gland ductal and secretory cells, bile ducts, gastrointestinal tract, bladder urothelium, oral epithelia, esophagus, and ectocervical epithelium. Anti-CK19 reacts with a wide variety of epithelial maignancies including adenocarcinomas of the colon, stomach, pancreas, biliary tract, liver, and breast. Perhaps the most useful application is the identification of thyroid carcinoma of the papillary type, eithough 50%-60% of folicular carcinomas are also labeled. Anti-CK19 is a useful application of thyroid carcinoma are also labeled. Anti-CK19 is a useful marker for detection of tumor cells in lymph nodes, peripheral blood, bone marrow and breast cancer.ImmunogenWestern Blot, Flow Cytometry, Flow (Intracellular), Immunofucothemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry, ImmunofucorescenceRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry, Immunofucothemistry/ Immunofluorescence, Immunohistochemistry, Immunofucothemistry/ Immunofluorescence, Immunohistochemistry, Immunofucothemistry/ Immunofluorescence	Clonality	Monoclonal
IsotypeIgG1 KappaConjugateFITCPurityProtein A or G purifiedBufferPBSProduct DescriptionThis conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.HostMouseGene ID3880Gene SymbolKRT19SpeciesHuman, Mouse, RatMarkerPancreatic Stem Cell MarkerSpecificity/SensitivityRecognizes a protein of 40kDa, identified as cytokeratin-19 (CK19), which is expressed in sweat gland, marmary gland ductal and secretory cells, bile ducts, gastrointestinal tract, bladder urothelium, oral epithelia, esophagus, and ectocervical epithelium. Anti-CK19 reacts with a wide variety of epithelial malignancies including adenocarcinomas of the colon, stomach, pancreas, biliary tract, liver, and breast. Perhaps the most useful application is the identification of thyroid carcinoma af the papilary type, although 50%-60% of follicular carcinomas are also labeled. Anti-CK19 is a useful marker for detection of tumor cells in lymph nodes, peripheral blood, bone marrow and breast cancer.ImmunogenWestern Blot, Flow Cytometry, Flow (Intracellular), Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry/ ImmunofluorescenceRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence	Clone	KRT19/800
ConjugateFITCPurityProtein A or G purifiedBufferPBSProduct DescriptionThis conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.HostMouseGene ID3880Gene SymbolKRT19SpeciesHuman, Mouse, RatMarkerPancreatic Stem Cell MarkerSpecificity/SensitivityRecognizes a protein of 40kDa, identified as cytokeratin-19 (CK19), which is expressed in sweat gland, mammary gland ductal and secretory cells, bile ducts, gastrointestinal tract, bladder urothelium, oral epithelia, esophagus, and ectocervical epithelium. Anti-CK19 reacts with a wide variety of epithelial malignancies including adenocarcinomas of the colon, stomach, pancreas, bilary tract, liver, and breast. Perhaps the most useful application is the identification of thyroid carcinoma of the papilary type, although 50%-60% of follicular carcinomas are also labeled. Anti-CK19 is a useful marker for detection of tumor cells in lymph nodes, peripheral blood, bone marrow and breast cancer.ImmunogenRecombiant full-length human Cytokeratin 19 protein. (Uniprot: P08727)Product Application DetailsWestern Blot, Flow Cytometry, Flow (Intracellular), Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry. Paraffin, Flow (Intracellular), Elw (Intracellular), Flow (Intracellular), Flow (Intracellular), Flow (Intracellular), Flow (Intracellular), Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, Flow (Intracellular), Flow (Intracellular),	Preservative	0.05% Sodium Azide
Purity Protein A or G purified Buffer PBS Product Description Pescription Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 3880 Gene Symbol KRT19 Species Human, Mouse, Rat Marker Pancreatic Stem Cell Marker Specificity/Sensitivity Recognizes a protein of 40kDa, identified as cytokeratin-19 (CK19), which is expressed in sweat gland, mammary gland ductal and secretory cells, bile ducts, gastrointestinal tract, bladder urothelium, oral epithelia, esophagus, and ectocervical epithelium. Anti-CK19 reacts with a wide variety of epithelial malignancies including adenocarcinomas of the colon, stomach, pancreas, biliary tract, liver, and breast. Perhaps the most useful application is the identification of thyroid carcinoma of the papillary type, although 50%-60% of follicular carcinomas are also labeled. Anti-CK19 is a useful marker for detection of tumor cells in lymph nodes, peripheral blood, bone marrow and breast cancer. Immunogen Recombiant full-length human Cytokeratin 19 protein. (Uniprot: P08727) Product Application Details Vestern Blot, Flow Cytometry, Flow (Intracellular), Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunocytochemistry/Immunofluorescence, Imm	Isotype	IgG1 Kappa
Buffer PBS Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 3880 Gene Symbol KRT19 Species Human, Mouse, Rat Marker Pancreatic Stem Cell Marker Specificity/Sensitivity Recognizes a protein of 40kDa, identified as cytokeratin-19 (CK19), which is expressed in sweat gland, mammary gland ductal and secretory cells, bile ducts, gastrointestinal tract, bladder urothelium, oral epithelia, esophagus, and ectocervical epithelium. Anti-CK19 reacts with a wide variety of epithelial malignancies including adenocarcinomas of the colon, stomach, pancreas, biliary tract, liver, and breast. Perhaps the most useful application is the identification of thyroid carcinoma of the papillary type, atthough 50%-60% of follicular carcinomas are also labeled. Anti-CK19 is a useful marker for detection of tumor cells in lymph nodes, peripheral blood, bone marrow and breast cancer. Immunogen Recombinant full-length human Cytokeratin 19 protein. (Uniprot: P08727) Product Application Details Western Blot, Flow Cytometry, Flow (Intracellular), Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Paraffin, Flow (Intracellular), Immunofluorescence, Immunohistochemistry, Paraffin, Flow (Intracellular), Immunofluorescence, Immunohistoc	Conjugate	FITC
Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 3880 Gene Symbol KRT19 Species Human, Mouse, Rat Marker Pancreatic Stem Cell Marker Specificity/Sensitivity Recognizes a protein of 40kDa, identified as cytokeratin-19 (CK19), which is expressed in sweat gland, mammary gland ductal and secretory cells, bile ducts, gastrointestinal tract, bladder urothelium, oral epithelia, esophagus, and ectocervical epithelium. Anti-CK19 reacts with a wide variety of epithelial malignancies including adenocarcinomas of the colon, stomach, pancreas, biliary tract, liver, and breast. Perhaps the most useful application is the identification of thyroid carcinoma of the papillary type, although 50%-60% of follicular carcinomas are also labeled. Anti-CK19 is a useful marker for detection of tumor cells in lymph nodes, peripheral blood, bone marrow and breast cancer. Immunogen Recombinant full-length human Cytokeratin 19 protein. (Uniprot: P08727) Product Application Details Western Blot, Flow Cytometry, Flow (Intracellular), Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Paraffin, Cy/TOF-ready, Immunofluorescence, Immunohistochemistry. Paraffin, Flow (Intracellular), Section 2005	Purity	Protein A or G purified
DescriptionThis conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.HostMouseGene ID3880Gene SymbolKRT19SpeciesHuman, Mouse, RatMarkerPancreatic Stem Cell MarkerSpecificity/SensitivityRecognizes a protein of 40kDa, identified as cytokeratin-19 (CK19), which is expressed in sweat gland, mammary gland ductal and secretory cells, bile ducts, gastrointestinal tract, bladder urothelium, oral epithelia, esophagus, and ectocervical epithelium. Anti-CK19 reacts with a wide variety of epithelial malignancies including adenocarcinomas of the colon, stomach, pancreas, biliary tract, liver, and breast. Perhaps the most useful application is the identification of thyroid carcinoma of the papillary type, although 50%-60% of follicular carcinomas are also labeled. Anti-CK19 is a useful marker for detection of tumor cells in lymph nodes, peripheral blood, bone marrow and breast cancer.ImmunogenWestern Blot, Flow Cytometry, Flow (Intracellular), Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry/ ImmunofluorescenceRecommended DilutionsWestern Blot, Flow Cytometry, Inmunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, Flow (Intracellular), Immunofluorescence, Immunohistochemist	Buffer	PBS
volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.HostMouseGene ID3880Gene SymbolKRT19SpeciesHuman, Mouse, RatMarkerPancreatic Stem Cell MarkerSpecificity/SensitivityRecognizes a protein of 40kDa, identified as cytokeratin-19 (CK19), which is expressed in sweat gland, mammary gland ductal and secretory cells, bile ducts, gastrointestinal tract, bladder urothelium, oral epithelia, esophagus, and ectocervical epithelium. Anti-CK19 reacts with a wide variety of epithelial malignancies including adenocarcinomas of the colon, stomach, pancreas, biliary tract, liver, and breast. Perhaps the most useful application is the identification of thyroid carcinoma of the papillary type, although 50%-60% of follicular carcinomas are also labeled. Anti-CK19 is a useful marker for detection of tumor cells in lymph nodes, peripheral blood, bone marrow and breast cancer.ImmunogenWestern Blot, Flow Cytometry, Flow (Intracellular), Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry/ ImmunofluorescenceRecommended DilutionsWestern Blot, Flow Cytometry, Inov Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, Flow (Intracellular), Immunofluorescence, Immunohistochemistry-Paraffin,	Product Description	
Gene ID3880Gene SymbolKRT19SpeciesHuman, Mouse, RatMarkerPancreatic Stem Cell MarkerSpecificity/SensitivityRecognizes a protein of 40kDa, identified as cytokeratin-19 (CK19), which is expressed in sweat gland, mammary gland ductal and secretory cells, bile ducts, gastrointestinal tract, bladder urothelium, oral epithelia, esophagus, and ectocervical epithelium. Anti-CK19 reacts with a wide variety of epithelial malignancies including adenocarcinomas of the colon, stomach, pancreas, biliary tract, liver, and breast. Perhaps the most useful application is the identification of thyroid carcinoma of the papillary type, although 50%-60% of follicular carcinomas are also labeled. Anti-CK19 is a useful marker for detection of tumor cells in lymph nodes, peripheral blood, bone marrow and breast cancer.ImmunogenRecombinant full-length human Cytokeratin 19 protein. (Uniprot: P08727)Product Application DetailsWestern Blot, Flow Cytometry, Flow (Intracellular), Immunocytochemistry/ ImmunofluorescenceApplicationsWestern Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry. Flow (Intracellular), Flow (Intracellular), Flow (Intracellular), Sectoremistry/ Immunofluorescence	Description	volume of this product. The volume will be greater than or equal to the unit size
Gene SymbolKRT19SpeciesHuman, Mouse, RatMarkerPancreatic Stem Cell MarkerSpecificity/SensitivityRecognizes a protein of 40kDa, identified as cytokeratin-19 (CK19), which is expressed in sweat gland, mammary gland ductal and secretory cells, bile ducts, gastrointestinal tract, bladder urothelium, oral epithelia, esophagus, and ectocervical epithelium. Anti-CK19 reacts with a wide variety of epithelial malignancies including adenocarcinomas of the colon, stomach, pancreas, biliary tract, liver, and breast. Perhaps the most useful application is the identification of thyroid carcinoma of the papillary type, although 50%-60% of follicular carcinomas are also labeled. Anti-CK19 is a useful marker for detection of tumor cells in lymph nodes, peripheral blood, bone marrow and breast cancer.ImmunogenRecombinant full-length human Cytokeratin 19 protein. (Uniprot: P08727)Product Application DetailsWestern Blot, Flow Cytometry, Flow (Intracellular), Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Flow (Intracellular), Flow (Intracellu	Host	Mouse
SpeciesHuman, Mouse, RatMarkerPancreatic Stem Cell MarkerSpecificity/SensitivityRecognizes a protein of 40kDa, identified as cytokeratin-19 (CK19), which is expressed in sweat gland, mammary gland ductal and secretory cells, bile ducts, gastrointestinal tract, bladder urothelium, oral epithelia, esophagus, and ectocervical epithelium. Anti-CK19 reacts with a wide variety of epithelial malignancies including adenocarcinomas of the colon, stomach, pancreas, biliary tract, liver, and breast. Perhaps the most useful application is the identification of thyroid carcinoma of the papillary type, although 50%-60% of follicular carcinomas are also labeled. Anti-CK19 is a useful marker for detection of tumor cells in lymph nodes, peripheral blood, bone marrow and breast cancer.ImmunogenRecombinant full-length human Cytokeratin 19 protein. (Uniprot: P08727)Product Application DetailsWestern Blot, Flow Cytometry, Flow (Intracellular), Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry-Paraffin, CyTOF-ready, ImmunofluorescenceRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Flow (Intracellular), Flow (Gene ID	3880
MarkerPancreatic Stem Cell MarkerSpecificity/SensitivityRecognizes a protein of 40kDa, identified as cytokeratin-19 (CK19), which is expressed in sweat gland, mammary gland ductal and secretory cells, bile ducts, gastrointestinal tract, bladder urothelium, oral epithelia, esophagus, and ectocervical epithelium. Anti-CK19 reacts with a wide variety of epithelial malignancies including adenocarcinomas of the colon, stomach, pancreas, biliary tract, liver, and breast. Perhaps the most useful application is the identification of thyroid carcinoma of the papillary type, although 50%-60% of follicular carcinomas are also labeled. Anti-CK19 is a useful marker for detection of tumor cells in lymph nodes, peripheral blood, bone marrow and breast cancer.ImmunogenRecombinant full-length human Cytokeratin 19 protein. (Uniprot: P08727)Product Application DetailsWestern Blot, Flow Cytometry, Flow (Intracellular), Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, ImmunofluorescenceRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry. Paraffin, Flow (Intracellular), Sector Sector)	Gene Symbol	KRT19
Specificity/SensitivityRecognizes a protein of 40kDa, identified as cytokeratin-19 (CK19), which is expressed in sweat gland, mammary gland ductal and secretory cells, bile ducts, gastrointestinal tract, bladder urothelium, oral epithelia, esophagus, and ectocervical epithelium. Anti-CK19 reacts with a wide variety of epithelial malignancies including adenocarcinomas of the colon, stomach, pancreas, biliary tract, liver, and breast. Perhaps the most useful application is the identification of thyroid carcinoma of the papillary type, although 50%-60% of follicular carcinomas are also labeled. Anti-CK19 is a useful marker for detection of tumor cells in lymph nodes, peripheral blood, bone marrow and breast cancer.ImmunogenRecombinant full-length human Cytokeratin 19 protein. (Uniprot: P08727)Product Application Details ApplicationsWestern Blot, Flow Cytometry, Flow (Intracellular), Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Paraffin, Flow (Intracellular), Flow (Intr	Species	Human, Mouse, Rat
expressed in sweat gland, mammary gland ductal and secretory cells, bile ducts, gastrointestinal tract, bladder urothelium, oral epithelia, esophagus, and ectocervical epithelium. Anti-CK19 reacts with a wide variety of epithelial malignancies including adenocarcinomas of the colon, stomach, pancreas, biliary tract, liver, and breast. Perhaps the most useful application is the identification of thyroid carcinoma of the papillary type, although 50%-60% of follicular carcinomas are also labeled. Anti-CK19 is a useful marker for detection of tumor cells in lymph nodes, peripheral blood, bone marrow and breast cancer.ImmunogenRecombinant full-length human Cytokeratin 19 protein. (Uniprot: P08727)Product Application DetailsWestern Blot, Flow Cytometry, Flow (Intracellular), Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Immunocytochemistry/ ImmunofluorescenceRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Paraffin, Flow (Intracellular), Flow (Intracellul	Marker	Pancreatic Stem Cell Marker
Product Application Details Applications Western Blot, Flow Cytometry, Flow (Intracellular), Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence Recommended Dilutions Western Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, Flow (Intracellular),	Specificity/Sensitivity	expressed in sweat gland, mammary gland ductal and secretory cells, bile ducts, gastrointestinal tract, bladder urothelium, oral epithelia, esophagus, and ectocervical epithelium. Anti-CK19 reacts with a wide variety of epithelial malignancies including adenocarcinomas of the colon, stomach, pancreas, biliary tract, liver, and breast. Perhaps the most useful application is the identification of thyroid carcinoma of the papillary type, although 50%-60% of follicular carcinomas are also labeled. Anti-CK19 is a useful marker for detection of tumor
ApplicationsWestern Blot, Flow Cytometry, Flow (Intracellular), Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, ImmunofluorescenceRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, Flow (Intracellular),	Immunogen	Recombinant full-length human Cytokeratin 19 protein. (Uniprot: P08727)
Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, ImmunofluorescenceRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, Flow (Intracellular),	Product Application Details	
Immunofluorescence, Immunohistochemistry-Paraffin, Flow (Intracellular),	Applications	Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin,
	Recommended Dilutions	Immunofluorescence, Immunohistochemistry-Paraffin, Flow (Intracellular),
Application NotesOptimal dilution of this antibody should be experimentally determined.	Application Notes	Optimal dilution of this antibody should be experimentally determined.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-47946F

NBP1-43319F	Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1) [FITC]
NBP1-78278PEP	Cytokeratin 19 Antibody Blocking Peptide
1129-ER-050	ErbB2/Her2 [Unconjugated]
NB100-687PEP	Cytokeratin 19 Antibody Blocking Peptide

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-47946F

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

