Product Datasheet

NF-L Antibody (NR-4) [Alexa Fluor® 750] NBP2-47969AF750

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-47969AF750

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-47969AF750



NBP2-47969AF750

NF-L Antibody (NR-4) [Alexa Fluor® 750]

Unit Size0.1 mlConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneNR-4Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 750PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description4747HostMoseGene BymbolNEFLSpeciesHuman, Rat, Porcine, Bovine, ChickenMarkerNeuronal MarkerSpecificity/SensitivityThis monoclonal antibody reacts with a 68kDa protein, identified as light sub-unit of neurofilaments consist of three major subunits with melecular weights of 68kDa (NF-L), 160kDa (NF-H), Anti- neurofilaments consist of three major subunits with melecular weights of 68kDa (NF-L), 160kDa (NF-H), Anti- neurofilaments sof as an unber of neurofilaments consist of three major subunits with melecular weights of 68kDa (NF-L), 160kDa (NF-H), Anti- neurofilaments sof as an unber of neurofilaments. Neuromas, ganglioneuromas, ganglioneuroblastomas, and neurofilaments servis as and endocrine carcinomas of the skin, and oard cell carcinomas of the lung also express neurofilament. Neurofilaments. Neuronal servis of the lung also express neurofilament.ImmunogenCrude neurofilament preparation from porcine spinal cord	Product Information	
StorageStore at 4C in the dark.ClonalityMonoclonalCloneNR-4Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 750PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID4747Gene SymbolNEFLSpeciesHuman, Rat, Porcine, Bovine, ChickenMarkerNeuronal MarkerSpecificity/SensitivityThis monoclonal antibody reacts with a 68kDa protein, identified as light sub-unit of neurofilaments (NF-L). Neurofilaments make up the main structural elements of axons and dendrites and are found in neurons, peripheral nerves, and sympathetic ganglion cells. Neurofilaments consist of three major subunits with molecular weights of 68kDa (NF-L), 160kDa (NF-M) and 200kDa (NF-H). Anti-neurofilament stains a number of neurofilament scient etamors. Neuromas, and neuroblastomas, and neuroblastomas stain positively for anti-neurofilament. Neurofilaments are also present in paragangiomas as well as adrenal and extra-adrenal pheochromocytomas. Carcinoids, neuroendocrine, carcinomas of the skin, and oat cell carcinomas of the lung also express neurofilament.	Unit Size	0.1 ml
ClonalityMonoclonalCloneNR-4Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 750PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID4747Gene SymbolNEFLSpeciesHuman, Rat, Porcine, Bovine, ChickenMarkerNeuronal MarkerSpecificity/SensitivityThis monoclonal antibody reacts with a 68kDa protein, identified as light sub-unit of neurofilaments (NF-L). Neurofilaments make up the main structural elements of axons and dendrites and are found in neurons, peripheral nerves, and sympathetic ganglion cells. Neurofilaments consist of three major subunits with molecular weights of 68kDa (NF-L), 160kDa (NF-H). Anti-neurofilaments (NF-L). Neurofilaments consist of three major subunits with molecular weights of 68kDa (NF-L), alokDa and extra-adrenal pheochromocytomas stain positively for anti-neurofilaments are also present in paraganglionas as well as adrenal and extra-adrenal pheochromocytomas Carcinoids, neuroendocrine carcinomas of the skin, and oat cell carcinomas of the lung also express neurofilament.	Concentration	
CloneNR-4Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 750PurityProtein A or G purifiedBuffer50MM Sodium BorateProduct DescriptionMouseHostMouseGene ID4747Gene SymbolNEFLSpeciesHuman, Rat, Porcine, Bovine, ChickenMarkerNeuronal MarkerSpecificity/SensitivityThis monoclonal antibody reacts with a 68kDa protein, identified as light sub-unit of neurofilaments (NF-L). Neurofilaments make up the main structural elements of axons and dendrites and are found in neurons, peripheral nerves, and sympathetic ganglion cells. Neurofilaments consist of three major subunits with molecular weights of 68kDa (NF-L). 160kDa (NF-M) and 200kDa (NF-H). Arti- neurofilaments stain positively for anti-neurofilament sare also present in paragangliomas as well as adrenal and extra-adrenal pheochromocytomas. Carcinoids, neuroendocrine, carcinomas of the skin, and oat cell carcinomas of the lung also express neurofilament.	Storage	Store at 4C in the dark.
Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 750PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description4747HostMouseGene ID4747Gene SymbolNEFLSpeciesHuman, Rat, Porcine, Bovine, ChickenMarkerNeuronal MarkerSpecificity/SensitivityThis monoclonal antibody reacts with a 68kDa protein, identified as light sub-unit of neurofilaments (NF-L). Neurofilaments make up the main structural elements of axons and dendrites and are found in neurons, peripheral nerves, and sympathetic ganglion cells. Neurofilaments consist of three major subunits with molecular weights of 68kDa (NF-L), 160kDa (NF-M) and 200kDa (NF-H). Anti-neurofilament stains a number of neurofilament. Neurofilaments are also present in paragangliomas as well as adrenal and extra-adrenal pheochromocytomas. Carcinoids, neuroendocrine carcinomas of the skin, and oat cell carcinomas of the lung also express neurofilament.	Clonality	Monoclonal
IsotypeIgG1 KappaConjugateAlexa Fluor 750PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID4747Gene SymbolNEFLSpeciesHuman, Rat, Porcine, Bovine, ChickenMarkerNeuronal MarkerSpecificity/SensitivityThis monoclonal antibody reacts with a 68kDa protein, identified as light sub-unit of neurofilaments (NF-L). Neurofilaments make up the main structural elements of axons and dendrites and are found in neurons, peripheral nerves, and sympathetic ganglion cells. Neurofilaments consist of three major subunits with molecular weights of 68kDa (NF-L), 160kDa (NF-M) and 200kDa (NF-H). Anti- neurofilament stains a number of neurafinaments are also present in paragangliomas as well as adrenal and extra-adrenal pheochromocytomas. Carcinoids, neuroendocrine carcinomas of the skin, and oat cell carcinomas of the lung also express neurofilament.	Clone	NR-4
ConjugateAlexa Fluor 750PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID4747Gene SymbolNEFLSpeciesHuman, Rat, Porcine, Bovine, ChickenMarkerNeuronal MarkerSpecificity/SensitivityThis monoclonal antibody reacts with a 68kDa protein, identified as light sub-unit of neurofilaments (NF-L). Neurofilaments make up the main structural elements of axons and dendrites and are found in neurons, peripheral nerves, and sympathetic ganglion cells. Neurofilaments consist of three major subunits with molecular weights of 68kDa (NF-L). 160kDa (NF-M) and 200kDa (NF-H). Anti-neurofilament stains a number of neural, neuroendocrine, and endocrine tumors. Neuromas, ganglioneuromas, ganglioneuroblastomas, and neuroblastomas stain positively for anti-neurofilament. Neurofilaments are also present in paragangliomas as well as drenal and extra-adrenal pheochromocytomas. Carcinoids, neuroendocrine carcinomas of the skin, and oat cell carcinomas of the lung also express neurofilament.	Preservative	0.05% Sodium Azide
PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID4747Gene SymbolNEFLSpeciesHuman, Rat, Porcine, Bovine, ChickenMarkerNeuronal MarkerSpecificity/SensitivityThis monoclonal antibody reacts with a 68kDa protein, identified as light sub-unit of neurofilaments (NF-L). Neurofilaments make up the main structural elements of axons and dendrites and are found in neurons, peripheral nerves, and sympathetic ganglion cells. Neurofilaments consist of three major subunits with molecular weights of 68kDa (NF-L), 160kDa (NF-M) and 200kDa (NF-H). Antineurofilament stains a number of neural, neuroendocrine, and endocrine tumors. Neuromas, ganglioneuromas, ganglioneuroblastomas, and neuroblastomas stain positively for anti-neurofilament. Neurofilaments are also present in paragangliomas as well as adrenal and extra-adrenal pheochromocytomas. Carcinoids, neuroendocrine carcinomas of the skin, and oat cell carcinomas of the lung also express neurofilament.	Isotype	IgG1 Kappa
Buffer50mM Sodium BorateProduct DescriptionHostMouseGene ID4747Gene SymbolNEFLSpeciesHuman, Rat, Porcine, Bovine, ChickenMarkerNeuronal MarkerSpecificity/SensitivityThis monoclonal antibody reacts with a 68kDa protein, identified as light sub-unit of neurofilaments (NF-L). Neurofilaments make up the main structural elements of axons and dendrites and are found in neurons, peripheral nerves, and sympathetic ganglion cells. Neurofilaments consist of three major subunits with molecular weights of 68kDa (NF-L), 160kDa (NF-M) and 200kDa (NF-H). Anti-neurofilament stains a number of neural, neuroendocrine, and endocrine tumors. Neuromas, ganglioneuroblastomas, and neuroblastomas stain positively for anti-neurofilaments are also present in paragangliomas as well as adrenal and extra-adrenal pheochromocytomas. Carcinoids, neuroendocrine carcinomas of the skin, and oat cell carcinomas of the lung also express neurofilament.	Conjugate	Alexa Fluor 750
Product DescriptionHostMouseGene ID4747Gene SymbolNEFLSpeciesHuman, Rat, Porcine, Bovine, ChickenMarkerNeuronal MarkerSpecificity/SensitivityThis monoclonal antibody reacts with a 68kDa protein, identified as light sub-unit of neurofilaments (NF-L). Neurofilaments make up the main structural elements of axons and dendrites and are found in neurons, peripheral nerves, and sympathetic ganglion cells. Neurofilaments consist of three major subunits with molecular weights of 68kDa (NF-L), 160kDa (NF-M) and 200kDa (NF-H). Anti- neurofilament stains a number of neural, neuroendocrine, and endocrine tumors. Neuromas, ganglioneuromas, gangliojemas, ganglioneuroblastomas, and neuroblastomas stain positively for anti-neurofilament. Neurofilaments are also present in paragangliomas as well as adrenal and extra-adrenal pheochromocytomas. Carcinoids, neuroendocrine carcinomas of the skin, and oat cell carcinomas of the lung also express neurofilament.	Purity	Protein A or G purified
HostMouseGene ID4747Gene SymbolNEFLSpeciesHuman, Rat, Porcine, Bovine, ChickenMarkerNeuronal MarkerSpecificity/SensitivityThis monoclonal antibody reacts with a 68kDa protein, identified as light sub-unit of neurofilaments (NF-L). Neurofilaments make up the main structural elements of axons and dendrites and are found in neurons, peripheral nerves, and sympathetic ganglion cells. Neurofilaments consist of three major subunits with molecular weights of 68kDa (NF-L), 160kDa (NF-M) and 200kDa (NF-H). Anti- neurofilament stains a number of neural, neuroendocrine, and endorrine tumors. Neuromas, ganglioneuromas, gangliogliomas, ganglioneuroblastomas, and neuroblastomas stain positively for anti-neurofilament. Neurofilaments are also present in paragangliomas as well as adrenal and extra-adrenal pheochromocytomas. Carcinoids, neuroendocrine carcinomas of the skin, and oat cell carcinomas of the lung also express neurofilament.	Buffer	50mM Sodium Borate
Gene ID4747Gene SymbolNEFLSpeciesHuman, Rat, Porcine, Bovine, ChickenMarkerNeuronal MarkerSpecificity/SensitivityThis monoclonal antibody reacts with a 68kDa protein, identified as light sub-unit of neurofilaments (NF-L). Neurofilaments make up the main structural elements of axons and dendrites and are found in neurons, peripheral nerves, and sympathetic ganglion cells. Neurofilaments consist of three major subunits with molecular weights of 68kDa (NF-L), 160kDa (NF-M) and 200kDa (NF-H). Anti- neurofilament stains a number of neural, neuroendocrine, and endocrine tumors. Neuromas, ganglioneuromas, gangliogliomas, ganglioneuroblastomas, and neuroblastomas stain positively for anti-neurofilament. Neurofilaments are also present in paragangliomas as well as adrenal and extra-adrenal pheochromocytomas. Carcinoids, neuroendocrine carcinomas of the skin, and oat cell carcinomas of the lung also express neurofilament.	Product Description	
Gene SymbolNEFLSpeciesHuman, Rat, Porcine, Bovine, ChickenMarkerNeuronal MarkerSpecificity/SensitivityThis monoclonal antibody reacts with a 68kDa protein, identified as light sub-unit of neurofilaments (NF-L). Neurofilaments make up the main structural elements of axons and dendrites and are found in neurons, peripheral nerves, and sympathetic ganglion cells. Neurofilaments consist of three major subunits with molecular weights of 68kDa (NF-L), 160kDa (NF-M) and 200kDa (NF-H). Anti- neurofilament stains a number of neural, neuroendocrine, and endocrine tumors. Neuromas, ganglioneuromas, gangliogliomas, ganglioneuroblastomas, and neuroblastomas stain positively for anti-neurofilament. Neurofilaments are also present in paragangliomas as well as adrenal and extra-adrenal pheochromocytomas. Carcinoids, neuroendocrine carcinomas of the skin, and oat cell carcinomas of the lung also express neurofilament.	Host	Mouse
SpeciesHuman, Rat, Porcine, Bovine, ChickenMarkerNeuronal MarkerSpecificity/SensitivityThis monoclonal antibody reacts with a 68kDa protein, identified as light sub-unit of neurofilaments (NF-L). Neurofilaments make up the main structural elements of axons and dendrites and are found in neurons, peripheral nerves, and sympathetic ganglion cells. Neurofilaments consist of three major subunits with molecular weights of 68kDa (NF-L), 160kDa (NF-M) and 200kDa (NF-H). Anti- neurofilament stains a number of neural, neuroendocrine, and endocrine tumors. Neuromas, ganglioneuromas, gangliogliomas, ganglioneuroblastomas, and neuroblastomas stain positively for anti-neurofilament. Neurofilaments are also present in paragangliomas as well as adrenal and extra-adrenal pheochromocytomas. Carcinoids, neuroendocrine carcinomas of the skin, and oat cell carcinomas of the lung also express neurofilament.	Gene ID	4747
MarkerNeuronal MarkerSpecificity/SensitivityThis monoclonal antibody reacts with a 68kDa protein, identified as light sub-unit of neurofilaments (NF-L). Neurofilaments make up the main structural elements of axons and dendrites and are found in neurons, peripheral nerves, and sympathetic ganglion cells. Neurofilaments consist of three major subunits with molecular weights of 68kDa (NF-L), 160kDa (NF-M) and 200kDa (NF-H). Anti- neurofilament stains a number of neural, neuroendocrine, and endocrine tumors. Neuromas, ganglioneuromas, gangliogliomas, ganglioneuroblastomas, and neuroblastomas stain positively for anti-neurofilament. Neurofilaments are also present in paragangliomas as well as adrenal and extra-adrenal pheochromocytomas. Carcinoids, neuroendocrine carcinomas of the skin, and oat cell carcinomas of the lung also express neurofilament.	Gene Symbol	NEFL
Specificity/Sensitivity This monoclonal antibody reacts with a 68kDa protein, identified as light sub-unit of neurofilaments (NF-L). Neurofilaments make up the main structural elements of axons and dendrites and are found in neurons, peripheral nerves, and sympathetic ganglion cells. Neurofilaments consist of three major subunits with molecular weights of 68kDa (NF-L), 160kDa (NF-M) and 200kDa (NF-H). Antineurofilament stains a number of neural, neuroendocrine, and endocrine tumors. Neuromas, ganglioneuromas, gangliogliomas, ganglioneuroblastomas, and neuroblastomas stain positively for anti-neurofilament. Neurofilaments are also present in paragangliomas as well as adrenal and extra-adrenal pheochromocytomas. Carcinoids, neuroendocrine carcinomas of the skin, and oat cell carcinomas of the lung also express neurofilament.	Species	Human, Rat, Porcine, Bovine, Chicken
of neurofilaments (NF-L). Neurofilaments make up the main structural elements of axons and dendrites and are found in neurons, peripheral nerves, and sympathetic ganglion cells. Neurofilaments consist of three major subunits with molecular weights of 68kDa (NF-L), 160kDa (NF-M) and 200kDa (NF-H). Anti- neurofilament stains a number of neural, neuroendocrine, and endocrine tumors. Neuromas, ganglioneuromas, gangliogliomas, ganglioneuroblastomas, and neuroblastomas stain positively for anti-neurofilament. Neurofilaments are also present in paragangliomas as well as adrenal and extra-adrenal pheochromocytomas. Carcinoids, neuroendocrine carcinomas of the skin, and oat cell carcinomas of the lung also express neurofilament.	Marker	Neuronal Marker
Immunogen Crude neurofilament preparation from porcine spinal cord	Specificity/Sensitivity	of neurofilaments (NF-L). Neurofilaments make up the main structural elements of axons and dendrites and are found in neurons, peripheral nerves, and sympathetic ganglion cells. Neurofilaments consist of three major subunits with molecular weights of 68kDa (NF-L), 160kDa (NF-M) and 200kDa (NF-H). Anti- neurofilament stains a number of neural, neuroendocrine, and endocrine tumors. Neuromas, ganglioneuromas, gangliogliomas, ganglioneuroblastomas, and neuroblastomas stain positively for anti-neurofilament. Neurofilaments are also present in paragangliomas as well as adrenal and extra-adrenal pheochromocytomas. Carcinoids, neuroendocrine carcinomas of the skin, and
	Immunogen	Crude neurofilament preparation from porcine spinal cord

www.novusbio.com



	Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.
Product Application Details	
Applications	Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence
Recommended Dilutions	Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence, CyTOF-ready
Application Notes	Optimal dilution of this antibody should be experimentally determined.
	•

Notes





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-47969AF750

IC002SMouse IgG1 Isotype Control (11711) [Alexa Fluor® 750]NBP2-59061PEPNF-L Recombinant Protein Antigen7954-GM-010/CFGM-CSF [Unconjugated]NBP2-80299Mouse NF-L ELISA Kit (Colorimetric)

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-47969AF750

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

