# **Product Datasheet**

# S100A9 Antibody (47-8D3 (DF3973)) [DyLight 550] NBP2-47980R

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-47980R

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP2-47980R



### NBP2-47980R

S100A9 Antibody (47-8D3 (DF3973)) [DyLight 550]

Stated on the datasheet.	3 100A9 Allibody (47-6D3 (DF	Saraj) [DyLight Sou]
Please see the vial label for concentration. If unlisted please contact technical services.	Product Information	
Storage Store at 4C in the dark.  Clonality Monoclonal  Clone 47-8D3 (DF3973)  Preservative 0.05% Sodium Azide  Isotype IgG1 Kappa  Conjugate DyLight 550  Purity Protein A or G purified  Buffer 50mM Sodium Borate  Product Description  Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.  Host Mouse  Gene ID 6280  Gene Symbol S100A9  Species Human, Mouse, Rat  Reactivity Notes Use in Mouse reported in scientific literature (PMID:32191647).  Marker Macrophage Marker  Specificity/Sensitivity This monoclonal antibody stains the cytoplasm of macrophages and histiccytes in hematopoietic organs, Kupffers cells of the skin. It also stains the mantle zon B-lymphocytes of the lymph node and spleen, spermatogonia, and chief cells on the stomach. S100A9 is also expressed by macrophages in acutely inflammed tissues and in chronic inflammation. It is detected in peripheral blood leukocytes, in neutrophils and granulocytes. It is present at sites of vascular inflammation. S100A9 is also expressed in epithelial cells constitutively or induced during dermatoses. S100A9 is also expressed in the cell constitutively or induced during dermatoses. S100A9 is a Calcium-binding protein. It has antimicrobial activity towards bacteria and fungi. It is important for resistance to invasion by pathogenic bacteria. It up-regulates the answerping and infiltration of granulocytes at sites of wounding. It plays a role as a pro-inflammatory mediator in acute and chronic inflammation and up-regulates the control inflammator and up-regulates the formation and up-fregulates the formation and up-f	Unit Size	0.1 ml
Clonality Monoclonal Clone 47-8D3 (DF3973) Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate DyLight 550 Purity Protein A or G purified Buffer 50mM Sodium Borate  Product Description Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.  Host Mouse Gene ID 6280 Gene Symbol S100A9 Species Human, Mouse, Rat Reactivity Notes Use in Mouse reported in scientific literature (PMID:32191647).  Marker Macrophage Marker Specificity/Sensitivity This monoclonal antibody stains the cytoplasm of macrophages and histiocytes in hematopoietic organs, Kupffers cells of the skin. It also stains the mantle zon B-lymphocytes of the lymph node and spleen, spermatogonia, and chief cells of the stomach. S100A9 is expressed by macrophages in actuely inflamed tissues and in chronic inflammation. It is detected in periphelial cells constitutively or induced during dermatoses. S100A9 is also expressed in epithelial cells constitutively or induced during dermatoses. S100A9 is a Calcium-binding protein. It has antimicrobial activity towards bacteria and fungi. It is important for resistance to invasion by pathogenic bacteria. It up-regulates the invasion by pathogenic bacteria. It up-regulates transcription of genes that are under the control of NF-kappa-B. S100A9 plays a role in the development of endotoxic shock in response to bacterial lipopolysaccharide (LPS). It promotes tubulin polymerization when unphosphorylated. It also promotes phagocyte migration and infiltration of granulocytes at sites of wounding. It plays a role as a pro-inflammatory mediator in acute and chronic inflammation and up-regulates the	Concentration	·
Clone 47-8D3 (DF3973)  Preservative 0.05% Sodium Azide  Isotype IgG1 Kappa  Conjugate DyLight 550  Purity Protein A or G purified  Buffer 50mM Sodium Borate  Product Description  Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.  Host Mouse  Gene ID 6280  Gene Symbol S100A9  Species Human, Mouse, Rat  Reactivity Notes Use in Mouse reported in scientific literature (PMID:32191647).  Marker Macrophage Marker  Specificity/Sensitivity  This monoclonal antibody stains the cytoplasm of macrophages and histiocytes in hematopoletic organs, Kupffers cells of the skin. It also stains the mantle zon B-lymphocytes of the lymph node and spleen, spermatogonia, and chief cells of the stomach. S100A9 is expressed by macrophages in acutely inflamed tissue and in chronic inflammation. It is detected in peripheral blood leukocytes, in neutrophils and granulocytes. It is present at sites of vascular inflammation. S100A9 is also expressed in epithelial cells constitutively or induced during dermatoses. S100A9 is a Calcium-binding protein. It has antimicrobial activity towards bacteria and fungi. It is important for restance to invasion by pathogenic bacteria. It up-regulates transcription of genes that are under the control of NF-kappa-B. S100A9 plays a role in the development of endotoxic shock in response to bacterial inpoplysacharide (IPS). It promotes tubulin polymerization when unphosphorylated. It also promotes phagocyte migration and infiltration of granulocytes at sites of wounding. It plays a role as a pro-inflammatory mediator in acute and chronic inflammation and up-regulates the	Storage	Store at 4C in the dark.
Preservative   0.05% Sodium Azide   Isotype   IgG1 Kappa   Conjugate   DyLight 550   Purity   Protein A or G purified   Buffer   50mM Sodium Borate   Product Description   Description   This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.  Host   Mouse   Gene ID   6280   Gene Symbol   S100A9   Species   Human, Mouse, Rat   Reactivity Notes   Use in Mouse reported in scientific literature (PMID:32191647).  Marker   Macrophage Marker   Specificity/Sensitivity   This monoclonal antibody stains the cytoplasm of macrophages and histiocytes in hematopoietic organs, Kupffers cells of the skin. It also stains the mantle zon B-lymphocytes of the lymph node and spleen, spermatogonia, and chief cells or the stomach. \$100A9 is expressed by macrophages in acutely inflamed tissues and in chronic inflammation. It is detected in peripheral blood leukocytes, in neutrophils and granulocytes. It is present at sites of vascular inflammation. \$100A9 is also expressed in epithelial cells constitutively or induced during dermatoses. \$100A9 is a Calcium-binding protein. It has antimicrobial activity towards bacteria and fungi. It is important for resistance to invasion by pathogenic bacteria. It up-regulates transcription of genes that are under the control of NF-kappa-B. \$100A9 plays a role in the development of endotoxic shock in response to bacterial lipopolysaccharide (LPS). It promotes tubulin polymerization when unphosphorylated. It also promotes phagocyte migration and infiltration of granulocytes at sites of wounding. It plays a role as a pro-infilammatory mediator in acute and chronic inflammation and up-regulates the	Clonality	Monoclonal
IgG1 Kappa	Clone	47-8D3 (DF3973)
Conjugate DyLight 550 Purity Protein A or G purified Buffer 50mM Sodium Borate  Product Description  This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.  Host Mouse Gene ID 6280 Gene Symbol S100A9 Species Human, Mouse, Rat Human, Mouse, Rat Reactivity Notes Use in Mouse reported in scientific literature (PMID:32191647).  Marker Specificity/Sensitivity This monoclonal antibody stains the cytoplasm of macrophages and histiocytes in hematopoietic organs, Kupffers cells of the skin. It also stains the mantle zon B-lymphocytes of the lymph node and spleen, spermatogonia, and chief cells of the stomach. S100A9 is expressed by macrophages in acutely inflamed tissues and in chronic inflammation. It is detected in peripheral blood leukocytes, in neutrophils and granulocytes. It is present at sites of vascular inflammation. S100A9 is a Calcium-binding protein. It has antimicrobial activity towards bacteria and fungi. It is important for resistance to invasion by pathogenic bacteria. It up-regulates transcription of genes that are under the control of NF-kappa-B. S100A9 plays a role in the development of endotoxic shock in response to bacterial lipopolysaccharide (LPS). It promotes tubulin polymerization when unphosphorylated. It also promotes phagocyte migration and infiltration of granulocytes at sites of wounding. It plays a role as a pro-inflammatory mediator in acute and chronic inflammation and up-regulates the	Preservative	0.05% Sodium Azide
Purity Protein A or G purified Buffer 50mM Sodium Borate  Product Description  This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.  Host Mouse  Gene ID 6280  Gene Symbol S100A9  Species Human, Mouse, Rat  Reactivity Notes Use in Mouse reported in scientific literature (PMID:32191647).  Marker Macrophage Marker  Specificity/Sensitivity  This monoclonal antibody stains the cytoplasm of macrophages and histiocytes in hematopoietic organs, Kupffers cells of the skin. It also stains the mantle zon B-lymphocytes of the lymph node and spleen, spermatogonia, and chief cells of the stomach. \$100A9 is expressed by macrophages in acutely inflamed tissues and in chronic inflammation. It is detected in peripheral blood leukocytes, in neutrophils and granulocytes. It is present at sites of vascular inflammation. S100A9 is also expressed in epithelial cells constitutively or induced during dermatoses. \$100A9 is a Calcium-binding protein. It has antimicrobial activity towards bacteria and fungi. It is important for resistance to invasion by pathogenic bacteria. It up-regulates transcription of genes that are under the control of NF-kappa-B. \$100A9 plays a role in the development of endotoxic shock in response to bacterial lipopolysaccharide (LPS). It promotes tubulin polymerization when unphosphorylated. It also promotes phagocyte migration and infiltration of granulocytes at sites of wounding. It plays a role as a pro-inflammatory mediator in acute and chronic inflammation and up-regulates the	Isotype	IgG1 Kappa
Product Description  This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.  Host  Mouse  Gene ID  6280  Gene Symbol  S100A9  Species  Human, Mouse, Rat  Reactivity Notes  Use in Mouse reported in scientific literature (PMID:32191647).  Marker  Macrophage Marker  Specificity/Sensitivity  This monoclonal antibody stains the cytoplasm of macrophages and histiocytes in hematopoietic organs, Kupffers cells of the skin. It also stains the mantle zon B-lymphocytes of the lymph node and spleen, spermatogonia, and chief cells on the stomach. S100A9 is expressed by macrophages in acutely inflamed tissues and in chronic inflammation. It is detected in peripheral blood leukocytes, in neutrophilis and granulocytes. It is present at sites of vascular inflammation. S100A9 is also expressed in epithelial cells constitutively or induced during dermatoses. S100A9 is a Calcium-binding protein. It has antimicrobial activity towards bacteria and fungi. It is important for resistance to invasion by pathogenic bacteria. It up-regulates transcription of genes that are under the control of NF-kappa-B. S100A9 plays a role in the development of endotoxic shock in response to bacterial lipopolysaccharide (LPS). It promotes tubulin polymerization when unphosphorylated. It also promotes phagocyte migration and infiltration of granulocytes at sites of wounding. It plays a role as a pro-inflammatory mediator in acute and chronic inflammation and up-regulates the	Conjugate	DyLight 550
Product Description  This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.  Mouse  Gene ID  6280  Gene Symbol  S100A9  Species  Human, Mouse, Rat  Reactivity Notes  Use in Mouse reported in scientific literature (PMID:32191647).  Marker  Macrophage Marker  Specificity/Sensitivity  This monoclonal antibody stains the cytoplasm of macrophages and histiocytes in hematopoietic organs, Kupffers cells of the skin. It also stains the mantle zon B-lymphocytes of the lymph node and spleen, spermatogonia, and chief cells o the stomach. S100A9 is expressed by macrophages in acutely inflamed tissues and in chronic inflammation. It is detected in peripheral blood leukocytes, in neutrophils and granulocytes. It is present at sites of vascular inflammation. S100A9 is also expressed in epithelial cells constitutively or induced during dermatoses. S100A9 is a Calcium-binding protein. It has antimicrobial activity towards bacteria and fungi. It is important for resistance to invasion by pathogenic bacteria. It up-regulates transcription of genes that are under the control of NF-kappa-B. S100A9 plays a role in the development of endotoxic shock in response to bacterial lipopolysaccharide (LPS). It promotes tubulin polymerization when unphosphorylated. It also promotes phagocyte migration and infiltration of granulocytes at sites of wounding. It plays a role as a proinflammation and up-regulates the	Purity	Protein A or G purified
This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.  Mouse  Gene ID  6280  Gene Symbol  S100A9  Species  Human, Mouse, Rat  Reactivity Notes  Use in Mouse reported in scientific literature (PMID:32191647).  Marker  Macrophage Marker  Specificity/Sensitivity  This monoclonal antibody stains the cytoplasm of macrophages and histiocytes in hematopoietic organs, Kupffers cells of the skin. It also stains the mantle zon B-lymphocytes of the lymph node and spleen, spermatogonia, and chief cells of the stomach. S100A9 is expressed by macrophages in acutely inflamed tissues and in chronic inflammation. It is detected in peripheral blood leukocytes, in neutrophils and granulocytes. It is present at sites of vascular inflammation. S100A9 is also expressed in epithelial cells constitutively or induced during dermatoses. S100A9 is a Calcium-binding protein. It has antimicrobial activity towards bacteria and fungi. It is important for resistance to invasion by pathogenic bacteria. It up-regulates transcription of genes that are under the control of NF-kappa-B. S100A9 plays a role in the development of endotoxic shock in response to bacterial lipopolysaccharide (LPS). It promotes tubulin polymerization when unphosphorylated. It also promotes phagocyte migration and infiltration of granulocytes at sites of wounding. It plays a role as a pro-inflammatory mediator in acute and chronic inflammation and up-regulates the	Buffer	50mM Sodium Borate
volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.  Mouse  Gene ID  6280  Gene Symbol  S100A9  Species  Human, Mouse, Rat  Reactivity Notes  Use in Mouse reported in scientific literature (PMID:32191647).  Marker  Macrophage Marker  Specificity/Sensitivity  This monoclonal antibody stains the cytoplasm of macrophages and histiocytes in hematopoietic organs, Kupffers cells of the skin. It also stains the mantle zon B-lymphocytes of the lymph node and spleen, spermatogonia, and chief cells of the stomach. S100A9 is expressed by macrophages in acutely inflamed tissues and in chronic inflammation. It is detected in peripheral blood leukocytes, in neutrophils and granulocytes. It is present at sites of vascular inflammation. S100A9 is also expressed in epithelial cells constitutively or induced during dermatoses. S100A9 is a Calcium-binding protein. It has antimicrobial activity towards bacteria and fungi. It is important for resistance to invasion by pathogenic bacteria. It up-regulates transcription of genes that are under the control of NF-kappa-B. S100A9 plays a role in the development of endotoxic shock in response to bacterial lipopolysaccharide (LPS). It promotes tubulin polymerization when unphosphorylated. It also promotes phagocyte migration and infiltration of granulocytes at sites of wounding. It plays a role as a proinflammatory mediator in acute and chronic inflammation and up-regulates the	Product Description	
Gene Symbol  Species  Human, Mouse, Rat  Reactivity Notes  Use in Mouse reported in scientific literature (PMID:32191647).  Marker  Specificity/Sensitivity  This monoclonal antibody stains the cytoplasm of macrophages and histiocytes in hematopoietic organs, Kupffers cells of the skin. It also stains the mantle zon B-lymphocytes of the lymph node and spleen, spermatogonia, and chief cells o the stomach. S100A9 is expressed by macrophages in acutely inflamed tissues and in chronic inflammation. It is detected in peripheral blood leukocytes, in neutrophils and granulocytes. It is present at sites of vascular inflammation. S100A9 is also expressed in epithelial cells constitutively or induced during dermatoses. S100A9 is a Calcium-binding protein. It has antimicrobial activity towards bacteria and fungi. It is important for resistance to invasion by pathogenic bacteria. It up-regulates transcription of genes that are under the control of NF-kappa-B. S100A9 plays a role in the development of endotoxic shock in response to bacterial lipopolysaccharide (LPS). It promotes tubullin polymerization when unphosphorylated. It also promotes phagocyte migration and infiltration of granulocytes at sites of wounding. It plays a role as a pro-inflammatory mediator in acute and chronic inflammation and up-regulates the	Description	volume of this product. The volume will be greater than or equal to the unit size
Gene Symbol  Species  Human, Mouse, Rat  Use in Mouse reported in scientific literature (PMID:32191647).  Marker  Macrophage Marker  This monoclonal antibody stains the cytoplasm of macrophages and histiocytes in hematopoietic organs, Kupffers cells of the skin. It also stains the mantle zon B-lymphocytes of the lymph node and spleen, spermatogonia, and chief cells of the stomach. S100A9 is expressed by macrophages in acutely inflamed tissues and in chronic inflammation. It is detected in peripheral blood leukocytes, in neutrophils and granulocytes. It is present at sites of vascular inflammation. S100A9 is also expressed in epithelial cells constitutively or induced during dermatoses. S100A9 is a Calcium-binding protein. It has antimicrobial activity towards bacteria and fungi. It is important for resistance to invasion by pathogenic bacteria. It up-regulates transcription of genes that are under the control of NF-kappa-B. S100A9 plays a role in the development of endotoxic shock in response to bacterial lipopolysaccharide (LPS). It promotes tubulin polymerization when unphosphorylated. It also promotes phagocyte migration and infiltration of granulocytes at sites of wounding. It plays a role as a pro-inflammatory mediator in acute and chronic inflammation and up-regulates the	Host	Mouse
Species  Human, Mouse, Rat  Use in Mouse reported in scientific literature (PMID:32191647).  Marker  Macrophage Marker  This monoclonal antibody stains the cytoplasm of macrophages and histiocytes in hematopoietic organs, Kupffers cells of the skin. It also stains the mantle zon B-lymphocytes of the lymph node and spleen, spermatogonia, and chief cells of the stomach. \$100A9 is expressed by macrophages in acutely inflamed tissues and in chronic inflammation. It is detected in peripheral blood leukocytes, in neutrophils and granulocytes. It is present at sites of vascular inflammation. \$100A9 is also expressed in epithelial cells constitutively or induced during dermatoses. \$100A9 is a Calcium-binding protein. It has antimicrobial activity towards bacteria and fungi. It is important for resistance to invasion by pathogenic bacteria. It up-regulates transcription of genes that are under the control of NF-kappa-B. \$100A9 plays a role in the development of endotoxic shock in response to bacterial lipopolysaccharide (LPS). It promotes tubulin polymerization when unphosphorylated. It also promotes phagocyte migration and infiltration of granulocytes at sites of wounding. It plays a role as a pro-inflammatory mediator in acute and chronic inflammation and up-regulates the	Gene ID	6280
Reactivity Notes  Use in Mouse reported in scientific literature (PMID:32191647).  Marker  Macrophage Marker  This monoclonal antibody stains the cytoplasm of macrophages and histiocytes in hematopoietic organs, Kupffers cells of the skin. It also stains the mantle zon B-lymphocytes of the lymph node and spleen, spermatogonia, and chief cells of the stomach. S100A9 is expressed by macrophages in acutely inflamed tissues and in chronic inflammation. It is detected in peripheral blood leukocytes, in neutrophils and granulocytes. It is present at sites of vascular inflammation. S100A9 is also expressed in epithelial cells constitutively or induced during dermatoses. S100A9 is a Calcium-binding protein. It has antimicrobial activity towards bacteria and fungi. It is important for resistance to invasion by pathogenic bacteria. It up-regulates transcription of genes that are under the control of NF-kappa-B. S100A9 plays a role in the development of endotoxic shock in response to bacterial lipopolysaccharide (LPS). It promotes tubulin polymerization when unphosphorylated. It also promotes phagocyte migration and infiltration of granulocytes at sites of wounding. It plays a role as a proinflammatory mediator in acute and chronic inflammation and up-regulates the	Gene Symbol	S100A9
Marker  Specificity/Sensitivity  This monoclonal antibody stains the cytoplasm of macrophages and histiocytes in hematopoietic organs, Kupffers cells of the skin. It also stains the mantle zon B-lymphocytes of the lymph node and spleen, spermatogonia, and chief cells of the stomach. S100A9 is expressed by macrophages in acutely inflamed tissues and in chronic inflammation. It is detected in peripheral blood leukocytes, in neutrophils and granulocytes. It is present at sites of vascular inflammation. S100A9 is also expressed in epithelial cells constitutively or induced during dermatoses. S100A9 is a Calcium-binding protein. It has antimicrobial activity towards bacteria and fungi. It is important for resistance to invasion by pathogenic bacteria. It up-regulates transcription of genes that are under the control of NF-kappa-B. S100A9 plays a role in the development of endotoxic shock in response to bacterial lipopolysaccharide (LPS). It promotes tubulin polymerization when unphosphorylated. It also promotes phagocyte migration and infiltration of granulocytes at sites of wounding. It plays a role as a pro-inflammatory mediator in acute and chronic inflammation and up-regulates the	Species	Human, Mouse, Rat
This monoclonal antibody stains the cytoplasm of macrophages and histiocytes in hematopoietic organs, Kupffers cells of the skin. It also stains the mantle zon B-lymphocytes of the lymph node and spleen, spermatogonia, and chief cells of the stomach. S100A9 is expressed by macrophages in acutely inflamed tissues and in chronic inflammation. It is detected in peripheral blood leukocytes, in neutrophils and granulocytes. It is present at sites of vascular inflammation. S100A9 is also expressed in epithelial cells constitutively or induced during dermatoses. S100A9 is a Calcium-binding protein. It has antimicrobial activity towards bacteria and fungi. It is important for resistance to invasion by pathogenic bacteria. It up-regulates transcription of genes that are under the control of NF-kappa-B. S100A9 plays a role in the development of endotoxic shock in response to bacterial lipopolysaccharide (LPS). It promotes tubulin polymerization when unphosphorylated. It also promotes phagocyte migration and infiltration of granulocytes at sites of wounding. It plays a role as a pro-inflammatory mediator in acute and chronic inflammation and up-regulates the	Reactivity Notes	Use in Mouse reported in scientific literature (PMID:32191647).
in hematopoietic organs, Kupffers cells of the skin. It also stains the mantle zon B-lymphocytes of the lymph node and spleen, spermatogonia, and chief cells of the stomach. S100A9 is expressed by macrophages in acutely inflamed tissues and in chronic inflammation. It is detected in peripheral blood leukocytes, in neutrophils and granulocytes. It is present at sites of vascular inflammation. S100A9 is also expressed in epithelial cells constitutively or induced during dermatoses. S100A9 is a Calcium-binding protein. It has antimicrobial activity towards bacteria and fungi. It is important for resistance to invasion by pathogenic bacteria. It up-regulates transcription of genes that are under the control of NF-kappa-B. S100A9 plays a role in the development of endotoxic shock in response to bacterial lipopolysaccharide (LPS). It promotes tubulin polymerization when unphosphorylated. It also promotes phagocyte migration and infiltration of granulocytes at sites of wounding. It plays a role as a pro-inflammatory mediator in acute and chronic inflammation and up-regulates the	Marker	Macrophage Marker
release of IL8 and cell-surface expression of ICAM1.	Specificity/Sensitivity	neutrophils and granulocytes. It is present at sites of vascular inflammation. S100A9 is also expressed in epithelial cells constitutively or induced during dermatoses. S100A9 is a Calcium-binding protein. It has antimicrobial activity towards bacteria and fungi. It is important for resistance to invasion by pathogenic bacteria. It up-regulates transcription of genes that are under the control of NF-kappa-B. S100A9 plays a role in the development of endotoxic shock in response to bacterial lipopolysaccharide (LPS). It promotes tubulin polymerization when unphosphorylated. It also promotes phagocyte migration and infiltration of granulocytes at sites of wounding. It plays a role as a pro-
Immunogen Human peripheral blood monocyte components (Uniprot: P06702)	Immunogen	Human peripheral blood monocyte components (Uniprot: P06702)
Notes  DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Notes	DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.

## **Product Application Details**



Applications	Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence
Recommended Dilutions	Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence, CyTOF-ready
Application Notes	Optimal dilution of this antibody should be experimentally determined.





### **Novus Biologicals USA**

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom

Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

### **Products Related to NBP2-47980R**

NBP1-43319R Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1) [DyLight 550]

NBP1-44500 Recombinant Human S100A9 His Protein

210-TA-005 TNF-alpha [Unconjugated]

DY5578 S100A9 [Biotin]

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-47980R

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

