# **Product Datasheet**

# Von Willebrand Factor Antibody (IIIE2.34) [Alexa Fluor® 488] NBP2-48008AF488

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-48008AF488

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.



Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-48008AF488

#### NBP2-48008AF488

Von Willebrand Factor Antibody (IIIE2.34) [Alexa Fluor® 488]

Product Information           Unit Size         0.1 ml           Concentration         Please see the vial label for concentration. If unlisted please contact technical services.           Storage         Store at 4C in the dark.           Clonality         Monoclonal           Clone         IIIE2.34           Preservative         0.05% Sodium Azide           Isotype         IgG1 Kappa           Conjugate         Alexa Fluor 488           Purity         Protein A or G purified           Buffer         500M Sodium Borate           Product Description         Gene ID           Host         Mouse           Gene ID         7450           Gene Symbol         VWF           Species         Human           Marker         Endothelial Marker           Specificity/Sensitivity         von Willebrand Factor (WF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. WP undergoes a variety of postranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propablie and formation of N-terminal disulfide bords. This artibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposi s sarcoma and cardiac myxoma. It is widely used for differentitating vacuular lesions from those of other tissue differentiatio	von willebrand Factor Antibody (	IIIE2.34) [Alexa Fluor® 488]	
ConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneIIIE2.34Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 488PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseGene ID7450Gene SymbolVWFSpeciesHumanMarkerEndothelial MarkerSpecificity/Sensitivityvon Willebrand Factor (VWF) is a multimeric glycoprotein that is found in endothelial cells, plasm and platelets. It acts as a carrier protein for Factor VIII and promotes platelet achesion and agregation. WF undergoes a variety of opostranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposi s sarrom and cardiac myxoma. It is widely used for differentiating vascular lesions from those of other tissue differentiation within a panel of other vascular markers although not all tumors of endothelial differentiation within a panel of other vascular markers although not all tumors of endothelial differentiation contain this antigen.	Product Information		
services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneIIIE2.34Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 488PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseGene ID7450Gene SymbolVWFSpeciesHumanMarkerEndothelial MarkerSpecificity/Sensitivityvon Willebrand Factor (VWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet achesions and acardiac myxoma. It is widely used for differentiation within a panel of other vascular markers although not all turnors of endothelial differentiation within a panel of other vascular markers although not all turnors of endothelial differentiation within a panel of other vascular markers although not all turnors of endothelial differentiation within a panel of other vascular markers although not all turnors of endothelial differentiation within a panel of other vascular markers although not all turnors of endothelial differentiation within a panel of other vascular markers although not all turnors of endothelial differentiation within a panel of other vascular markers although not all turnors of endothelial differentiation contain this antigen.	Unit Size	0.1 ml	
ClonalityMonoclonalCloneIIIE2.34Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 488PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID7450Gene SymbolVWFSpeciesHumanMarkerEndothelial MarkerSpeciesHumanMarkerEndothelial MarkerSpecificity/Sensitivityvon Wilebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet athesion and aggregation. WF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial mature of some lesions of disputed histogenesis, e.g. Kaposi s sarcoma and cardiac myxoma. It is widely used for differentiating vascular markers although not all tumors of endothelial differentiation contain this antigen.ImmunogenRecombinant human Von Willebrand Factor fragment spanning around aa 845-	Concentration		
CloneIIIE2.34Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 488PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostHostMouseGene ID7450Gene SymbolVWFSpeciesHumanMarkerEndothelial MarkerSpecificity/Sensitivityvon Willebrand Factor (WVF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propertie and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposi s sarcoma and cardiac myxoma. It is widely used for differentiating vascular markers although not all tumors of endothelial differentiation contain this antigen.ImmunogenRecombinant human Von Willebrand Factor fragment spanning around aa 845-	Storage	Store at 4C in the dark.	
Preservative       0.05% Sodium Azide         Isotype       IgG1 Kappa         Conjugate       Alexa Fluor 488         Purity       Protein A or G purified         Buffer       50mM Sodium Borate         Product Description       Host         Host       Mouse         Gene ID       7450         Species       Human         Marker       Endothelial Marker         Specificity/Sensitivity       von Willebrand Factor (VWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. VWF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposi's sarcoma and cardiac myxoma. It is widely used for differentiating vascular markers although not all tumors of endothelial differentiation contain this antigen.         Immunogen       Recombinant human Von Willebrand Factor fragment spanning around aa 845-	Clonality	Monoclonal	
IsotypeIgG1 KappaConjugateAlexa Fluor 488PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID7450Gene SymbolVWFSpeciesHumanMarkerEndothelial MarkerSpecificity/Sensitivityvon Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kapos is sarcoma and cardiac myxoma. It is widely used for differentiation contain this antigen.ImmunogenRecombinant human Von Willebrand Factor fragment spanning around aa 845-	Clone	IIIE2.34	
ConjugateAlexa Fluor 488PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID7450Gene SymbolVWFSpeciesHumanMarkerEndothelial MarkerSpecificity/Sensitivityvon Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposi s sarcoma and cardiac myxoma. It is widely used for differentiating vascular markers although not all tumors of endothelial differentiation contain this antigen.ImmunogenRecombinant human Von Willebrand Factor fragment spanning around aa 845-	Preservative	0.05% Sodium Azide	
Purity       Protein A or G purified         Buffer       50mM Sodium Borate         Product Description       Mouse         Gene ID       7450         Gene Symbol       VWF         Species       Human         Marker       Endothelial Marker         Specificity/Sensitivity       von Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposi s sarcoma and cardiac myxoma. It is widely used for differentiating vascular markers although not all tumors of endothelial differentiation contain this antigen.         Immunogen       Recombinant human Von Willebrand Factor fragment spanning around aa 845-	Isotype	IgG1 Kappa	
Buffer       50mM Sodium Borate         Product Description       Mouse         Host       Mouse         Gene ID       7450         Gene Symbol       VWF         Species       Human         Marker       Endothelial Marker         Specificity/Sensitivity       von Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposi s sarcoma and cardiac myxoma. It is widely used for differentiating vascular lesions from those of other tissue differentiation within a panel of other vascular markers although not all tumors of endothelial differentiation contain this antigen.         Immunogen       Recombinant human Von Willebrand Factor fragment spanning around aa 845-	Conjugate	Alexa Fluor 488	
Product DescriptionHostMouseGene ID7450Gene SymbolVWFSpeciesHumanMarkerEndothelial MarkerSpecificity/Sensitivityvon Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposi s sarcoma and cardiac myxoma. It is widely used for differentiating vascular lesions from those of other tissue differentiation within a panel of other vascular markers although not all tumors of endothelial differentiation contain this antigen.ImmunogenRecombinant human Von Willebrand Factor fragment spanning around aa 845-	Purity	Protein A or G purified	
HostMouseGene ID7450Gene SymbolVWFSpeciesHumanMarkerEndothelial MarkerSpecificity/Sensitivityvon Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposi s sarcoma and cardiac myxoma. It is widely used for differentiating vascular lesions from those of other tissue differentiation within a panel of other vascular markers although not all tumors of endothelial differentiation contain this antigen.ImmunogenRecombinant human Von Willebrand Factor fragment spanning around aa 845-	Buffer	50mM Sodium Borate	
Gene ID7450Gene SymbolVWFSpeciesHumanMarkerEndothelial MarkerSpecificity/Sensitivityvon Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposi s sarcoma and cardiac myxoma. It is widely used for differentiating vascular lesions from those of other tissue differentiation within a panel of other vascular markers although not all tumors of endothelial differentiation contain this antigen.ImmunogenRecombinant human Von Willebrand Factor fragment spanning around aa 845-	Product Description		
Gene SymbolVWFSpeciesHumanMarkerEndothelial MarkerSpecificity/Sensitivityvon Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposi s sarcoma and cardiac myxoma. It is widely used for differentiating vascular lesions from those of other tissue differentiation within a panel of other vascular markers although not all tumors of endothelial differentiation contain this antigen.ImmunogenRecombinant human Von Willebrand Factor fragment spanning around aa 845-	Host	Mouse	
SpeciesHumanMarkerEndothelial MarkerSpecificity/Sensitivityvon Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposi s sarcoma and cardiac myxoma. It is widely used for differentiating vascular lesions from those of other tissue differentiation within a panel of other vascular markers although not all tumors of endothelial differentiation contain this antigen.ImmunogenRecombinant human Von Willebrand Factor fragment spanning around aa 845-	Gene ID	7450	
MarkerEndothelial MarkerSpecificity/Sensitivityvon Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposi s sarcoma and cardiac myxoma. It is widely used for differentiating vascular lesions from those of other tissue differentiation within a panel of other vascular markers although not all tumors of endothelial differentiation contain this antigen.ImmunogenRecombinant human Von Willebrand Factor fragment spanning around aa 845-	Gene Symbol	VWF	
Specificity/Sensitivityvon Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposi s sarcoma and cardiac myxoma. It is widely used for differentiating vascular lesions from those of other tissue differentiation within a panel of other vascular markers although not all tumors of endothelial differentiation contain this antigen.ImmunogenRecombinant human Von Willebrand Factor fragment spanning around aa 845-	Species	Human	
<ul> <li>endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposi s sarcoma and cardiac myxoma. It is widely used for differentiating vascular lesions from those of other tissue differentiation within a panel of other vascular markers although not all tumors of endothelial differentiation contain this antigen.</li> <li>Immunogen</li> </ul>	Marker	Endothelial Marker	
ImmunogenRecombinant human Von Willebrand Factor fragment spanning around aa 845- 949 (exact sequence is proprietary) (Uniprot: P04275 )		endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposi s sarcoma and cardiac myxoma. It is widely used for differentiating vascular lesions from those of other tissue differentiation within a panel of other vascular markers although not all tumors of endothelial differentiation contain this antigen.	
	Immunogen	Recombinant human Von Willebrand Factor fragment spanning around aa 845- 949 (exact sequence is proprietary) (Uniprot: P04275)	



Notes	Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.
Product Application Details	
Applications	Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation, CyTOF-ready, Immunofluorescence
Recommended Dilutions	Western Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunoprecipitation, Immunohistochemistry-Paraffin, Immunofluorescence, CyTOF-ready
Application Notes	Optimal dilution of this antibody should be experimentally determined.





#### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

## **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

## **General Contact Information**

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

#### Products Related to NBP2-48008AF488

IC002G	Mouse IgG1 Isotype Control (11711) [Alexa Fluor® 488]
NBP1-84762PEP	Von Willebrand Factor Recombinant Protein Antigen
210-TA-005	TNF-alpha [Unconjugated]
KA0512	Human Von Willebrand Factor ELISA Kit (Colorimetric)

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-48008AF488

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

