Product Datasheet

TFF1/pS2 Antibody (SPM313) [Alexa Fluor® 647] NBP2-48009AF647

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-48009AF647

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-48009AF647



NBP2-48009AF647

TFF1/pS2 Antibody (SPM313) [Alexa Fluor® 647]

Product InformationUnit Size0.1 mlConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneSPM313Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 647PurityProtein A or G purifiedBufferSomM Sodium BorateProduct DescriptionTFF1HostMouseGene ID7031Gene SymbolTFF1SpeciesHuman, Cynomolgus MonkeySpeciesIt recognized perites dong through the gut hough to play a role in its protein. Its epitope is localized between as57-84 of human p52 progulated throughout the gut that play a role in mucosal healing. These peptides contain three intra-chain disulifie bords, form dimerization is throught to play a role in its protective and healing properties. About 60% of breast carcinomas are positive for p52. Staning is cytoplasmic, often with localization to the Golgi apparatus. pS2 is shown to be localized in normal storach mucosal peating. These peptides contain three intra-chain disulifie bords, forming the trefol mucosal peoties contain three intra-chain disulifie bords, forming the trefol mucosal peoties contain three intra-chain disulifie bords, forming the trefol mucosal peoties contain three intra-chain disulifie bords, forming the trefol mucosal peoties contain three intra-chain disulifie bords, forming the trefol mucosal peoties contain three intra-chain disulifie bords, forming the trefol motion. p52 is known to form dimers and this dimerization is throught to play a role in its protective and healing properties. About 60% of breast carcinomas are positive fo		-
ConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneSPM313Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseGene ID7031Gene ID7031Gene SymbolTFF1SpeciesHuman, Cynomolgus MonkeySpeciesHuman, Cynomolgus MonkeySpeciesIt recognizes a polypeptide of 6.5kDa, identified as pS2 estrogen-regulated protein. Its epitope is localized between aa57-84 of human pS2 is a trefoil petide. Trefoil petides are protease resistant molecules screted throughout the gut that play a role in mucosal healing. These peptides contain three intra-chain disulfide bonds, forming the trefoil motif, or P-domain. pS2 is a trefoil petides are protease resistant molecules screted throughout the gut that play a role in mucosal healing. These peptides contain three intra-chain disulfide bonds, forming the trefoil motif, or P-domain. pS2 is shown to be localized in normal stomach mucosa, gastric fluid, goblet cells in the colon and small intestine, and in ulcerations of the gastrointestinal tract. Several studies have shown that pS2 is primarily expressed in estrogen- receptor-positive breast tumors and it may define a subset of estrogen- dependent tumors that displays an increased likelihood of response to endocrine therapy.ImmunogenA synthetic peptide (around aa 57-84) of human TFF1/pS2 polypeptide (exact	Product Information	
services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneSPM313Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 647PurityProtein A or G purifiedBufferSomM Sodium BorateProduct DescriptionTFF1HostMouseGene ID7031Gene SymbolTFF1Specificity/SensitivityIt recognizes a polypeptide of 6.5kDa, identified as pS2 estrogen-regulated protein. Its epitope is localized between aa57-84 of human pS2 protein. pS2 is a normal somath mousal notability of proses and this dimerization is thought to play a role in its protective and healing properties. About 60% of breast carcinomas are positive for pS2. Staining is cytoplasmic, often with localization to the Golgi apparatus. Sp2 is shown to be localized in ulceration is thought to play a prote in its protective and healing properties. About 60% of breast carcinomas are positive for pS2. Staining is cytoplasmic, often with localization to the Golgi apparatus. Sp2 is shown to be localized in normal stomach mucosa, gastric fluid, goblet cells in the colon and small intestine, and in ulcerations of the gastrointestinal tract. Several studies have shown that pS2 is primarily expressed in estrogen-dependent tumors that displays an increased likelihood of response to endocrine therapy.ImmunogenA synthetic peptide (around aa 57-84) of human TFF1/pS2 polypeptide (exact	Unit Size	0.1 ml
ClonalityMonoclonalCloneSPM313Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseGene ID7031Gene SymbolTFF1SpeciesHuman, Cynomolgus MonkeySpeciesHuman, Cynomolgus MonkeySpeciestrecognizes a polypeptide of 6.5kDa, identified as pS2 estrogen-regulated protein. Its epitope is localized between aa57-84 of human pS2 protein. pS2 is a trefoil peptide. Trefoil peptides. Forming the trefoil motif, or P-domain. pS2 is known to form dimers and this dimerization is thought to play a role in its protective and healing properties. About 60% of breast carcinomas are positive for pS2. Staining is cytoplasmic, often with localization to the Golgi apparatus. pS2 is shown to be localized in normal stomach mucosa, gastric fluid, goblet cells in the colon and small intestine, and in ulcerations of the gastrointestinal tract. Several studies have shown that pS2 is primarily expressed in estrogen receptor-positive breast tumors and it may define a subset of estrogen-dependent tumors that displays an increased likelihood of response to endocrine therapy.ImmunogenA synthetic peptide (around as 57-84) of human TFF1/pS2 polypeptide (exact	Concentration	
CloneSPM313Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description6ene IDHostMouseGene ID7031Gene SymbolTFF1SpeciesHuman, Cynomolgus MonkeySpeciestrecognizes a polypeptide of 6.5kDa, identified as pS2 estrogen-regulated protein. Its epitope is localized between aa57-84 of human pS2 rotein. pS2 is a trefoil peptide. Trefoil peptides are protease resistant molecules secreted through ut the gut that play a role in mucosal healing. These peptides contain three intra-chain disulfide bonds, forming the trefoil motif, or P-domain. pS2 is known to form dimers and this dimerization is thought to play a role in its protective and healing properties. About 60% of breast carcinomas are positive for pS2. Staining is cytoplasmic, often with localization to the Golgi apparatus. pS2 is shown to be localized in normal stomach mucosa, gastric fluid, goblet cells in the colon and small intestine, and in ulcerations of the gastrointestinal tract. Everal studies have shown that pS2 is primarily expressed in estrogen receptor-positive breast tumors and it may define a subset of estrogen-dependent tumors that displays an increased likelihood of response to endocrine therapy.ImmunogenA synthetic peptide (around aa 57-84) of human TFF1/pS2 polypeptide (exact	Storage	Store at 4C in the dark.
Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID7031Gene SymbolTFF1SpeciesHuman, Cynomolgus MonkeySpecificity/SensitivityIt recognizes a polypeptide of 6.5kDa, identified as pS2 estrogen-regulated protein. Its epitope is localized between aa57-84 of human pS2 protein. pS2 is a trefoil peptide. Trefoil peptides are protease resistant molecules secreted through out the gut that play a role in mucosal healing. These peptides contain three intra-chain disulfide bonds, forming the trefoil motif, or P-domain. pS2 is known to form dimers and this dimerization is thought to play a role in its protective and healing properties. About 60% of breast carcinomas are positive for pS2. Staining is cytoplasmic, often with localization to the Golgi apparatus. pS2 is shown to be localized in normal stomach mucosa, gastric fluid, goblet cells in the colon and small intestine, and in ulcerations of the gastrointestinal tract. Several studies have shown that pS2 is primarily expressed in estrogen receptor-positive breast tumors and it may define a subset of estrogen-dependent tumors that displays an increased likelihood of response to endocrine therapy.ImmunogenA synthetic peptide (around aa 57-84) of human TFF1/pS2 polypeptide (exact	Clonality	Monoclonal
IsotypeIgG1 KappaConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID7031Gene SymbolTFF1SpeciesHuman, Cynomolgus MonkeySpeciesHuman, Cynomolgus MonkeySpecieshuman, Cynomolgus Monkeyspecificity/SensitivityIt recognizes a polypeptide of 6.5kDa, identified as pS2 estrogen-regulated protein. Its epitope is localized between aa57-84 of human pS2 protein. pS2 is a trefoil peptide. Trefoil peptides are protease resistant molecules secreted throughout the gut that play a role in mucosal healing. These peptides contain three intra-chain disulfide bonds, forming the trefoil motif, or P-domain. pS2 is known to form dimers and this dimerization is thought a pole in its protective and healing properties. About 60% of breast carcinomas are positive for pS2. Staining is cytoplasmic, often with localization to the Golgi apparatus. pS2 is shown to be localized in normal stomach mucosa, gastric fluid, goblet cells in the colon and small intestine, and in ulcerations of the gastrointestinal tract. Several studies have shown that pS2 is primarily expressed in estrogen receptor-positive breast tumors and it may define a subset of estrogen-dependent tumors that displays an increased likelihood of response to endocrine therapy.ImmunogenA synthetic peptide (around aa 57-84) of human TFF1/pS2 polypeptide (exact	Clone	SPM313
ConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseGene ID7031Gene SymbolTFF1SpeciesHuman, Cynomolgus MonkeySpecificity/SensitivityIt recognizes a polypeptide of 6.5kDa, identified as pS2 estrogen-regulated protein. Its epitope is localized between aa57-84 of human pS2 protein. pS2 is a trefol peptide. Trefol peptides are protease resistant molecules secreted throughout the gut that play a role in mucosal healing. These peptides contain three intra-chain disulfide bonds, forming the trefoil motif, or P-domain. pS2 is known to form dimers and this dimerization is though to play a role in its protective and healing properties. About 60% of breast carcinomas are positive for pS2. Staining is cytoplasmic, often with localization to the Golgi apparatus. pS2 is shown to be localized in normal stomach mucosa, gastric fluid, goblet cells in the colo and small intestine, and in ulcerations of the gastrointestinal tract. Several studies have shown that pS2 is primarily expressed in estrogen- dependent tumors that displays an increased likelihood of response to endocrine therapy.ImmunogenA synthetic peptide (around aa 57-84) of human TFF1/pS2 polypeptide (exact	Preservative	0.05% Sodium Azide
PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID7031Gene SymbolTFF1SpeciesHuman, Cynomolgus MonkeySpecificity/SensitivityIt recognizes a polypeptide of 6.5kDa, identified as pS2 estrogen-regulated protein. Its epitope is localized between aa57-84 of human pS2 protein. pS2 is a trefoil peptide. Trefoil peptides are protease resistant molecules secreted throughout the gut that play a role in mucosal healing. These peptides contain three intra-chain disulfide bonds, forming the trefoil motif, or P-domain. pS2 is known to form dimers and this dimerization is though to play a role in its protective and healing properties. About 60% of breast carcinomas are positive for pS2. Staining is cytoplasmic, often with localization to the Golgi apparatus. pS2 is shown to be localized in normal stomach mucosa, gastric fluid, goblet cells in the colon and small intestine, and in ulcerations of the gastrointestinal tract. Several studies have shown that pS2 is primarily expressed in estrogen- dependent tumors that displays an increased likelihood of response to endocrine thereapy.ImmunogenA synthetic peptide (around aa 57-84) of human TFF1/pS2 polypeptide (exact	Isotype	IgG1 Kappa
Buffer50mM Sodium BorateProduct DescriptionHostMouseGene ID7031Gene SymbolTFF1SpeciesHuman, Cynomolgus MonkeySpecificity/SensitivityIt recognizes a polypeptide of 6.5kDa, identified as pS2 estrogen-regulated protein. Its epitope is localized between aa57-84 of human pS2 protein. pS2 is a trefoil peptide. Trefoil peptides are protease resistant molecules secreted throughout the gut that play a role in mucosal healing. These peptides contain three intra-chain disulfide bonds, forming the trefoil motif, or P-domain. pS2 is known to form dimers and this dimerization is thought to play a role in its protective and healing properties. About 60% of breast carcinomas are positive for pS2. Staining is cytoplasmic, often with localization to the Golgi apparatus. pS2 is shown to be localized in normal stomach mucosa, gastric fluid, goblet cells in the colon and small intestine, and in ulcerations of the gastrointestinal tract. Several studies have shown that pS2 is primarily expressed in estrogen- dependent tumors that displays an increased likelihood of response to endocrine therapy.ImmunogenA synthetic peptide (around aa 57-84) of human TFF1/pS2 polypeptide (exact	Conjugate	Alexa Fluor 647
Product DescriptionHostMouseGene ID7031Gene SymbolTFF1SpeciesHuman, Cynomolgus MonkeySpecificity/SensitivityIt recognizes a polypeptide of 6.5kDa, identified as pS2 estrogen-regulated protein. Its epitope is localized between aa57-84 of human pS2 protein. pS2 is a trefoil peptide. Trefoil peptides are protease resistant molecules secreted throughout the gut that play a role in mucosal healing. These peptides contain three intra-chain disulfide bonds, forming the trefoil motif, or P-domain. pS2 is known to form dimers and this dimerization is thought to play a role in its protective and healing properties. About 60% of breast carcinomas are positive for pS2. Staining is cytoplasmic, often with localization to the Golgi apparatus. pS2 is shown to be localized in normal stomach mucosa, gastric fluid, goblet cells in the colon and small intestine, and in ulcerations of the gastrointestinal tract. Several studies have shown that pS2 is primarily expressed in estrogen receptor-positive breast tumors and it may define a subset of estrogen- dependent tumors that displays an increased likelihood of response to endocrine therapy.ImmunogenA synthetic peptide (around as 57-84) of human TFF1/pS2 polypeptide (exact	Purity	Protein A or G purified
HostMouseGene ID7031Gene SymbolTFF1SpeciesHuman, Cynomolgus MonkeySpecificity/SensitivityIt recognizes a polypeptide of 6.5kDa, identified as pS2 estrogen-regulated protein. Its epitope is localized between aa57-84 of human pS2 protein. pS2 is a trefoil peptide. Trefoil peptides are protease resistant molecules secreted throughout the gut that play a role in mucosal healing. These peptides contain three intra-chain disulfide bonds, forming the trefoil motif, or P-domain. pS2 is known to form dimers and this dimerization is thought to play a role in its protective and healing properties. About 60% of breast carcinomas are positive for pS2. Staining is cytoplasmic, often with localization to the Golgi apparatus. pS2 is shown to be localized in normal stomach mucosa, gastric fluid, goblet cells in the colon and small intestine, and in ulcerations of the gastrointestinal tract. Several studies have shown that pS2 is primarily expressed in estrogen- dependent tumors that displays an increased likelihood of response to endocrine therapy.ImmunogenA synthetic peptide (around aa 57-84) of human TFF1/pS2 polypeptide (exact	Buffer	50mM Sodium Borate
Gene ID7031Gene SymbolTFF1SpeciesHuman, Cynomolgus MonkeySpecificity/SensitivityIt recognizes a polypeptide of 6.5kDa, identified as pS2 estrogen-regulated protein. Its epitope is localized between aa57-84 of human pS2 protein. pS2 is a trefoil peptide. Trefoil peptides are protease resistant molecules secreted throughout the gut that play a role in mucosal healing. These peptides contain three intra-chain disulfide bonds, forming the trefoil motif, or P-domain. pS2 is known to form dimers and this dimerization is thought to play a role in its protective and healing properties. About 60% of breast carcinomas are positive for pS2. Staining is cytoplasmic, often with localization to the Golgi apparatus. pS2 is shown to be localized in normal stomach mucosa, gastric fluid, goblet cells in the colon and small intestine, and in ulcerations of the gastrointestinal tract. Several studies have shown that pS2 is primarily expressed in estrogen- dependent tumors that displays an increased likelihood of response to endocrine therapy.ImmunogenA synthetic peptide (around aa 57-84) of human TFF1/pS2 polypeptide (exact	Product Description	
Gene SymbolTFF1SpeciesHuman, Cynomolgus MonkeySpecificity/SensitivityIt recognizes a polypeptide of 6.5kDa, identified as pS2 estrogen-regulated protein. Its epitope is localized between aa57-84 of human pS2 protein. pS2 is a trefoil peptide. Trefoil peptides are protease resistant molecules secreted throughout the gut that play a role in mucosal healing. These peptides contain three intra-chain disulfide bonds, forming the trefoil motif, or P-domain. pS2 is known to form dimers and this dimerization is thought to play a role in its protective and healing properties. About 60% of breast carcinomas are positive for pS2. Staining is cytoplasmic, often with localization to the Golgi apparatus. pS2 is shown to be localized in normal stomach mucosa, gastric fluid, goblet cells in the colon and small intestine, and in ulcerations of the gastrointestinal tract. Several studies have shown that pS2 is primarily expressed in estrogen receptor-positive breast tumors and it may define a subset of estrogen- dependent tumors that displays an increased likelihood of response to endocrine therapy.ImmunogenA synthetic peptide (around aa 57-84) of human TFF1/pS2 polypeptide (exact	Host	Mouse
SpeciesHuman, Cynomolgus MonkeySpecificity/SensitivityIt recognizes a polypeptide of 6.5kDa, identified as pS2 estrogen-regulated protein. Its epitope is localized between aa57-84 of human pS2 protein. pS2 is a trefoil peptide. Trefoil peptides are protease resistant molecules secreted throughout the gut that play a role in mucosal healing. These peptides contain three intra-chain disulfide bonds, forming the trefoil motif, or P-domain. pS2 is known to form dimers and this dimerization is thought to play a role in its protective and healing properties. About 60% of breast carcinomas are positive for pS2. Staining is cytoplasmic, often with localization to the Golgi apparatus. pS2 is shown to be localized in normal stomach mucosa, gastric fluid, goblet cells in the colon and small intestine, and in ulcerations of the gastrointestinal tract. Several studies have shown that pS2 is primarily expressed in estrogen receptor-positive breast tumors and it may define a subset of estrogen- dependent tumors that displays an increased likelihood of response to endocrine therapy.ImmunogenA synthetic peptide (around aa 57-84) of human TFF1/pS2 polypeptide (exact	Gene ID	7031
Specificity/SensitivityIt recognizes a polypeptide of 6.5kDa, identified as pS2 estrogen-regulated protein. Its epitope is localized between aa57-84 of human pS2 protein. pS2 is a trefoil peptide. Trefoil peptides are protease resistant molecules secreted throughout the gut that play a role in mucosal healing. These peptides contain three intra-chain disulfide bonds, forming the trefoil motif, or P-domain. pS2 is known to form dimers and this dimerization is thought to play a role in its protective and healing properties. About 60% of breast carcinomas are positive for pS2. Staining is cytoplasmic, often with localization to the Golgi apparatus. pS2 is shown to be localized in normal stomach mucosa, gastric fluid, goblet cells in the colon and small intestine, and in ulcerations of the gastrointestinal tract. Several studies have shown that pS2 is primarily expressed in estrogen- dependent tumors that displays an increased likelihood of response to endocrine therapy.ImmunogenA synthetic peptide (around aa 57-84) of human TFF1/pS2 polypeptide (exact	Gene Symbol	TFF1
 protein. Its epitope is localized between aa57-84 of human pS2 protein. pS2 is a trefoil peptide. Trefoil peptides are protease resistant molecules secreted throughout the gut that play a role in mucosal healing. These peptides contain three intra-chain disulfide bonds, forming the trefoil motif, or P-domain. pS2 is known to form dimers and this dimerization is thought to play a role in its protective and healing properties. About 60% of breast carcinomas are positive for pS2. Staining is cytoplasmic, often with localization to the Golgi apparatus. pS2 is shown to be localized in normal stomach mucosa, gastric fluid, goblet cells in the colon and small intestine, and in ulcerations of the gastrointestinal tract. Several studies have shown that pS2 is primarily expressed in estrogen receptor-positive breast tumors and it may define a subset of estrogen-dependent tumors that displays an increased likelihood of response to endocrine therapy. Immunogen 	Species	Human, Cynomolgus Monkey
	Specificity/Sensitivity	protein. Its epitope is localized between aa57-84 of human pS2 protein. pS2 is a trefoil peptide. Trefoil peptides are protease resistant molecules secreted throughout the gut that play a role in mucosal healing. These peptides contain three intra-chain disulfide bonds, forming the trefoil motif, or P-domain. pS2 is known to form dimers and this dimerization is thought to play a role in its protective and healing properties. About 60% of breast carcinomas are positive for pS2. Staining is cytoplasmic, often with localization to the Golgi apparatus. pS2 is shown to be localized in normal stomach mucosa, gastric fluid, goblet cells in the colon and small intestine, and in ulcerations of the gastrointestinal tract. Several studies have shown that pS2 is primarily expressed in estrogen receptor-positive breast tumors and it may define a subset of estrogen-dependent tumors that displays an increased likelihood of response to endocrine
	Immunogen	



Notes	Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.
Product Application Details	
Applications	Immunohistochemistry, Immunohistochemistry-Paraffin
Recommended Dilutions	Immunohistochemistry, Immunohistochemistry-Paraffin
Application Notes	Optimal dilution of this antibody should be experimentally determined.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-48009AF647

IC002R	Mouse IgG1 Isotype Control (11711) [Alexa Fluor® 647]
NBP2-35042-5ug	Recombinant Human TFF1/pS2 Protein
236-EG-200	EGF [Unconjugated]
DY5237	TFF1/pS2 [Biotin]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-48009AF647

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

