Product Datasheet

alpha-Fetoprotein/AFP Antibody (C2) - IHC-Prediluted NBP2-48121

Unit Size: 7 ml

Store at 4C.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-48121

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-48121



NBP2-48121


alpha-Fetoprotein/AFP Antibody (C2) - IHC-Prediluted

,
7 ml
Please see the vial label for concentration. If unlisted please contact technical services.
Store at 4C.
Monoclonal
C2
0.05% Sodium Azide
IgG1 Kappa
Protein A or G purified
10 mM PBS with 0.05% BSA
70 kDa
The prediluted antibody does not require any mixing, dilution, reconstitution, or titration; the antibody is ready-to-use and optimized for staining.
Mouse
174
AFP
Human, Mouse, Bovine (Negative), Canine (Negative), Rat (Negative)
Does not react with Bovine, Canine or Rat.
Hepatocellular/Germ Cell Tumor Marker
It recognizes an oncofetal glycoprotein with a single chain of 70kDa, which is identified as alpha fetoprotein (AFP) (ISOBM TD-2 workshop). This monoclonal antibody is highly specific to AFP and shows no cross-reaction with other oncofetal antigens or serum albumin. AFP is normally synthesized in the liver, intestinal tract, and yolk sac of the fetus. Antibody to AFP has been shown to be useful in detecting hepatocellular carcinomas (HCC) and germ cell neoplasms, especially yolk sac tumors.
Alpha feto protein (AFP) Purified from serum of a hepatoma patient (Uniprot: P02771)
Immunohistochemistry, Immunohistochemistry-Paraffin
Immunohistochemistry, Immunohistochemistry-Paraffin
Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes. Optimal dilution for a specific application should be determined.



Images

Immunohistochemistry-Paraffin: alpha-Fetoprotein/AFP Antibody (C2) -IHC-Prediluted [NBP2-48121] - Human Fetal Liver stained with AFP Mouse Monoclonal Antibody (C2).







Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-48121

NBP2-52171-0.05mg	Recombinant Human alpha-Fetoprotein/AFP His Protein
NBP1-43319-0.5mg	Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1)
NB720-B	Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]
HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-48121

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

