

Product Datasheet

CD20 Antibody (IGEL/1497R) [Biotin] NBP2-49873B

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-49873B

Updated 10/26/2023 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-49873B



NBP2-49873B

CD20 Antibody (IGEL/1497R) [Biotin]

| Product Information | |
|------------------------------------|---|
| Unit Size | 0.1 ml |
| Concentration | Please see the vial label for concentration. If unlisted please contact technical services. |
| Storage | Store at 4C in the dark. |
| Clonality | Monoclonal |
| Clone | IGEL/1497R |
| Preservative | 0.05% Sodium Azide |
| Isotype | IgG |
| Conjugate | Biotin |
| Purity | Protein A purified |
| Buffer | PBS |
| Product Description | |
| Host | Rabbit |
| Gene ID | 931 |
| Gene Symbol | MS4A1 |
| Species | Human |
| Reactivity Notes | Does not react with Canine. |
| Marker | B-Cell Marker |
| Specificity/Sensitivity | Recognizes a protein of 30-33kDa, which is identified as CD20. It is a non-Ig differentiation antigen of B-cells and its expression is restricted to normal and neoplastic B-cells, being absent from all other leukocytes and tissues. CD20 is expressed by pre B-cells and persists during all stages of B-cell maturation but is lost upon terminal differentiation into plasma cells. This monoclonal antibody can be used for immunophenotyping of leukemia and malignant cells, B lymphocyte detection in peripheral blood and B cell localization in tissues. It reacts with the majority of B-cells present in peripheral blood and lymphoid tissues and their derived lymphomas. In lymphoid tissue, germinal center blasts and B-immunoblasts are particularly reactive. It is a reliable antibody for ascribing a B-cell phenotype in known lymphoid tissues. Rarely, CD20-positive T-cell lymphomas have been reported. Reactivity has also been noted with Reed-Sternberg cells in cases of Hodgkins disease, particularly of lymphocyte predominant type. |
| Immunogen | Recombinant human full-length CD20 protein (Uniprot: P11836) |
| Product Application Details | |
| Applications | Flow Cytometry, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready |
| Recommended Dilutions | Flow Cytometry, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready |
| Application Notes | Optimal dilution of this antibody should be experimentally determined. |





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-49873B

| | |
|-----------------|-------------------------------------|
| NBP2-24891B | Rabbit IgG Isotype Control [Biotin] |
| NBP2-61455-10ug | Recombinant Human CD20 His Protein |
| 210-TA-005 | TNF-alpha [Unconjugated] |
| NBL1-13312 | CD20 Overexpression Lysate |

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-49873B

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

