

Product Datasheet

eRF1 Antibody (4F9H12) - BSA Free NBP2-52552

Unit Size: 0.1 mg

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 1


Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-52552

Updated 4/13/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-52552



NBP2-52552

eRF1 Antibody (4F9H12) - BSA Free

Product Information

Unit Size	0.1 mg
Concentration	1 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	4F9H12
Preservative	0.05% Sodium Azide
Isotype	IgG1
Purity	Protein G purified
Buffer	PBS
Target Molecular Weight	49 kDa

Product Description


Host	Mouse
Gene ID	2107
Gene Symbol	ETF1
Species	Human
Reactivity Notes	Use in Human reported in scientific literature (PMID:33764477).
Immunogen	Purified recombinant fragment of human eRF1 (AA: 288-437) expressed in E. Coli.

Product Application Details


Applications	Western Blot, Simple Western, ELISA, Flow Cytometry, CyTOF-ready
Recommended Dilutions	Western Blot 1:500-1:2000, Simple Western 1:100, Flow Cytometry 1:200-1:400, ELISA 1:10000, CyTOF-ready
Application Notes	This antibody is Cytof ready.

Images


Western Blot: eRF1 Antibody (4F9H12) - BSA Free [NBP2-52552] - Effect of CC-885 and CC-90009 on COL17A1 PTC readthrough. JEB01 keratinocytes derived from a JEB patient with a COL17A1 nonsense mutation (R688X/R688X) were exposed to the indicated concentrations of CC-90009 for 72 h and eRF3a, eRF3b and eRF1 (NBP2-52552) levels were determined using automated capillary electrophoresis western analysis. GAPDH was used as loading control. Image collected and cropped by CiteAb from the following publication ([//pubmed.ncbi.nlm.nih.gov/33764477/](https://pubmed.ncbi.nlm.nih.gov/33764477/)) licensed under a CC-BY license.




Flow Cytometry: eRF1 Antibody (4F9H12) - BSA Free [NBP2-52552] - Analysis of HepG2 cells using RF1 mouse mAb (green) and negative control (red).




Western Blot: eRF1 Antibody (4F9H12) - BSA Free [NBP2-52552] - Analysis using RF1 mAb against HEK293 (1) and RF1 (AA: 288-437)-hlgGfc transfected HEK293 (2) cell lysate.




Western Blot: eRF1 Antibody (4F9H12) - BSA Free [NBP2-52552] - Analysis using RF1 mAb against human RF1 (AA: 288-437) recombinant protein. (Expected MW is 43 kDa)




Western Blot: eRF1 Antibody (4F9H12) - BSA Free [NBP2-52552] - Analysis using eRF1 mouse mAb against MCF-7 (1), T47D (2), MOLT4 (3), and Raji (4) cell lysate.




Flow Cytometry: eRF1 Antibody (4F9H12) - BSA Free [NBP2-52552] - Analysis of Hela cells using RF1 mouse mAb (green) and negative control (red).



ELISA: eRF1 Antibody (4F9H12) - BSA Free [NBP2-52552] - Black line: Control Antigen (100 ng); Purple line: Antigen (10ng); Blue line: Antigen (50 ng); Red line: Antigen (100 ng).



Simple Western: eRF1 Antibody (4F9H12) - BSA Free [NBP2-52552] - Effect of CC-885 & CC-90009 on COL17A1 PTC readthrough. (A, B) HaCaT keratinocytes derived from an unaffected individual were exposed to various concentrations of CC-885 (A) or CC-90009 (B) with or without 100 μ g/ml gentamicin sulfate for 48 h & cell viability was measured using the MTT assay in triplicate samples ($\pm$ SD). Concentrations of CC-885 & CC-90009 are in nanomolar. (C, D) JEB01 keratinocytes derived from a JEB patient with a COL17A1 nonsense mutation (R688X/R688X) were exposed to the indicated concentrations of CC-885 (C) or CC-90009 (D) for 72 h & eRF3a, eRF3b & eRF1 levels were determined using automated capillary electrophoresis western analysis. GAPDH was used as loading control. (E, F) JEB01 keratinocytes were exposed to the indicated concentrations of CC-885 (E) or CC-90009 (F) in the presence or absence of 100 or 300 μ g/ml gentamicin sulfate for 72 h & Collagen XVII levels were measured by SDS-PAGE followed by traditional western blotting. β -actin was used as loading control. Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/33764477>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Publications

Baradaran-Heravi A, Balgi AD, Hosseini-Farahabadi S et al. Effect of small molecule eRF3 degraders on premature termination codon readthrough Nucleic acids research 2021-03-25 [PMID: 33764477] (WB, Human)

**Novus Biologicals USA**

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-52552

HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB720-B	Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]
NBP1-97005-0.5mg	Mouse IgG1 Isotype Control (MG1)
H00002107-P01-10ug	Recombinant Human eRF1 GST (N-Term) Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-52552

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

