Product Datasheet

CD4 Antibody (CE9.1 (Clenoliximab)) [PE/Cy7] - Chimeric NBP2-52681PECY7

Unit Size: 0.1 ml

Store at 4C in the dark. Do not freeze.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-52681PECY7

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.





NBP2-52681PECY7

CD4 Antibody (CE9.1 (Clenoliximab)) [PE/Cy7] - Chimeric

services.StorageStore at 4C in the dark. Do not freeze.ClonalityMonoclonalCloneCE9.1 (Clenoliximab)Preservative0.05% Sodium AzideIsotypeIgG LambdaConjugatePE/Cy7PurityProtein A purifiedBufferPBSProduct DescriptionRabbitHostRabbitGene ID920Gene SymbolCD4SpeciesHuman, ChimpanzeeReactivity NotesHuman, ChimpanzeeSpecificity/SensitivityThe antibody binds to human CD4 with a Kd of 1.0 nM and an ED50 of 0.2 ug/ml.ImmunogenDomain 1 of human CD4.Product Application DetailsFlow CytometryApplicationsFlow CytometryApplication NotesOptimal results using our Tandem dyes, please avoid prolonged exposure to 1 or extreme temperature fluctuations. These can lead to irreversible degradatio or decoupling. When staining intracellular targets, specific attention to the fixa and permeabilization steps in your flow protocol may be required. Please con our technical support team at technical@novusbie.com if you have any		
Concentration Please see the vial label for concentration. If unlisted please contact technica services. Storage Store at 4C in the dark. Do not freeze. Clonality Monoclonal Clone CE9.1 (Clenoliximab) Preservative 0.05% Sodium Azide Isotype IgG Lambda Conjugate PE/Cy7 Purity Protein A purified Buffer PBS Product Description Host Host Rabbit Gene ID 920 Gene Symbol CD4 Species Human, Chimpanzee Reactivity Notes Human, Chimpanzee Specificity/Sensitivity The antibody binds to human CD4 with a Kd of 1.0 nM and an ED50 of 0.2 ug/ml. Immunogen Domain 1 of human CD4. Product Application Details Flow Cytometry Applications Flow Cytometry Recommended Dilutions Flow Cytometry Application Notes Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to resulte degradatio or exterme temperature fluctuations. These can lead to irreverishele degradatio or ex	Product Information	
services. Storage Store at 4C in the dark. Do not freeze. Clonality Monoclonal Clone CE9.1 (Clenoliximab) Preservative 0.05% Sodium Azide Isotype IgG Lambda Conjugate PE/Cy7 Purity Protein A purified Buffer PBS Product Description Rabbit Gene ID 920 Gene Symbol CD4 Species Human, Chimpanzee Reactivity Notes Human, Chimpanzee Specificity/Sensitivity The antibody binds to human CD4 with a Kd of 1.0 nM and an ED50 of 0.2 ug/ml. Immunogen Domain 1 of human CD4. Product Application Details Flow Cytometry Applications Flow Cytometry Application Notes Optimal results using our Tandem dyes, please avoid prolonged exposure to 1 or extreme temperature fluctuations. These can lead to irreversible degradatio or decoupling. When staining intracellular targets, specific attention to the fixa and permeabilization steps in your flow protocol may be required. Please com our technical support team at technical@novusbio.com if you have any	Unit Size	0.1 ml
Clonality Monoclonal Clone CE9.1 (Clenoliximab) Preservative 0.05% Sodium Azide Isotype IgG Lambda Conjugate PE/Cy7 Purity Protein A purified Buffer PBS Product Description Host Host Rabbit Gene ID 920 Gene Symbol CD4 Species Human, Chimpanzee Reactivity Notes Human, Chimpanzee Species Human, Chimpanzee Specificity/Sensitivity The antibody binds to human CD4 with a Kd of 1.0 nM and an ED50 of 0.2 ug/ml. Immunogen Domain 1 of human CD4. Product Application Details Flow Cytometry Applications Flow Cytometry Application Notes Optimal results using our Tandem dyes, please avoid prolonged exposure to 1 or extreme temperature fluctuations. These can lead to irreversible degradatio or decoupling. When staining intracellular targets, specific attention to the fix and permeabilization steps in your flow protocol may be required. Please con or our technical support team at technical@envousbic.com if you have any	Concentration	Please see the vial label for concentration. If unlisted please contact technical services.
CloneCE9.1 (Clenoliximab)Preservative0.05% Sodium AzideIsotypeIgG LambdaConjugatePE/Cy7PurityProtein A purifiedBufferPBSProduct DescriptionHostHostRabbitGene ID920Gene SymbolCD4SpeciesHuman, ChimpanzeeReactivity NotesHuman, ChimpanzeeSpecificity/SensitivityThe antibody binds to human CD4 with a Kd of 1.0 nM and an ED50 of 0.2 ug/ml.ImmunogenDomain 1 of human CD4.Product Application DetailsFlow CytometryApplicationsFlow CytometryApplication NotesOptimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to 1 or extreme temperature fluctuations. These can lead to irreversible degradation of decoupling. When staining intracellular targets, specific attention to the fixa and permeabilization steps in your flow protocol may be required. Please con our technical support team at technical@novusbio.com if you have any	Storage	Store at 4C in the dark. Do not freeze.
Preservative 0.05% Sodium Azide Isotype IgG Lambda Conjugate PE/Cy7 Purity Protein A purified Buffer PBS Product Description Host Host Rabbit Gene ID 920 Gene Symbol CD4 Species Human, Chimpanzee Reactivity Notes Human, Chimpanzee Specificity/Sensitivity The antibody binds to human CD4 with a Kd of 1.0 nM and an ED50 of 0.2 ug/ml. Immunogen Domain 1 of human CD4. Product Application Details Flow Cytometry Applications Flow Cytometry Application Notes Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to 1 or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixa and permeabilization steps in your flow protocol may be required. Please con our technical support team at technical@novusbio.com if you have any	Clonality	Monoclonal
Isotype IgG Lambda Conjugate PE/Cy7 Purity Protein A purified Buffer PBS Product Description Host Rabbit Gene ID 920 Gene Symbol CD4 Species Human, Chimpanzee Reactivity Notes Human, Chimpanzee Specificity/Sensitivity The antibody binds to human CD4 with a Kd of 1.0 nM and an ED50 of 0.2 ug/ml. Immunogen Domain 1 of human CD4. Product Application Details Flow Cytometry Applications Flow Cytometry Recommended Dilutions Flow Cytometry Application Notes Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to 1 or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixa and permeabilization steps in your flow protocol may be required. Please con our technical support team at technical@novusbio.com if you have any	Clone	CE9.1 (Clenoliximab)
Conjugate PE/Cy7 Purity Protein A purified Buffer PBS Product Description Rabbit Host Rabbit Gene ID 920 Gene Symbol CD4 Species Human, Chimpanzee Reactivity Notes Human, Chimpanzee Specificity/Sensitivity The antibody binds to human CD4 with a Kd of 1.0 nM and an ED50 of 0.2 ug/ml. Immunogen Domain 1 of human CD4. Product Application Details Flow Cytometry Applications Flow Cytometry Application Notes Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to 1 or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixa and permeabilization steps in your flow protocol may be required. Please com our technical support team at technical@novusbio.com if you have any	Preservative	0.05% Sodium Azide
Purity Protein A purified Buffer PBS Product Description Rabbit Host Rabbit Gene ID 920 Gene Symbol CD4 Species Human, Chimpanzee Reactivity Notes Human, Chimpanzee Specificity/Sensitivity The antibody binds to human CD4 with a Kd of 1.0 nM and an ED50 of 0.2 ug/ml. Immunogen Domain 1 of human CD4. Product Application Details Flow Cytometry Applications Flow Cytometry Recommended Dilutions Flow Cytometry Application Notes Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to 1 or extreme temperature fluctuations. These can lead to irreversible degradatio or decoupling. When staining intracellular targets, specific attention to the fixa and permeabilization steps in your flow protocol may be required. Please con our technical support team at technical@novusbio.com if you have any	Isotype	IgG Lambda
Buffer PBS Product Description Rabbit Host Rabbit Gene ID 920 Gene Symbol CD4 Species Human, Chimpanzee Reactivity Notes Human, Chimpanzee Specificity/Sensitivity The antibody binds to human CD4 with a Kd of 1.0 nM and an ED50 of 0.2 ug/ml. Immunogen Domain 1 of human CD4. Product Application Details Flow Cytometry Applications Flow Cytometry Application Notes Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to 1 or extreme temperature fluctuations. These can lead to irreversible degradatito or decoupling. When staining intracellular targets, specific attention to the fixa and permeabilization steps in your flow protocol may be required. Please con our technical support team at technical@novusbio.com if you have any	Conjugate	PE/Cy7
Product Description Host Rabbit Gene ID 920 Gene Symbol CD4 Species Human, Chimpanzee Reactivity Notes Human, Chimpanzee Specificity/Sensitivity The antibody binds to human CD4 with a Kd of 1.0 nM and an ED50 of 0.2 ug/ml. Immunogen Domain 1 of human CD4. Product Application Details Applications Applications Flow Cytometry Recommended Dilutions Flow Cytometry Application Notes Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to 1 or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixe and permeabilization steps in your flow protocol may be required. Please con our technical support team at technical@novusbio.com if you have any	Purity	Protein A purified
HostRabbitGene ID920Gene SymbolCD4SpeciesHuman, ChimpanzeeReactivity NotesHuman, ChimpanzeeSpecificity/SensitivityThe antibody binds to human CD4 with a Kd of 1.0 nM and an ED50 of 0.2 ug/ml.ImmunogenDomain 1 of human CD4.Product Application DetailsFlow CytometryApplicationsFlow CytometryRecommended DilutionsFlow CytometryApplication NotesOptimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to I or extreme temperature fluctuations. These can lead to irreversible degradatio or decoupling. When staining intracellular targets, specific attention to the fixa and permeabilization steps in your flow protocol may be required. Please con our technical support team at technical@novusbio.com if you have any	Buffer	PBS
Gene ID920Gene SymbolCD4SpeciesHuman, ChimpanzeeReactivity NotesHuman, ChimpanzeeSpecificity/SensitivityThe antibody binds to human CD4 with a Kd of 1.0 nM and an ED50 of 0.2 ug/ml.ImmunogenDomain 1 of human CD4.Product Application DetailsFlow CytometryApplicationsFlow CytometryRecommended DilutionsFlow CytometryApplication NotesOptimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to I or extreme temperature fluctuations. These can lead to irreversible degradatio or decoupling. When staining intracellular targets, specific attention to the fixa and permeabilization steps in your flow protocol may be required. Please con our technical support team at technical@novusbio.com if you have any	Product Description	
Gene Symbol CD4 Species Human, Chimpanzee Reactivity Notes Human, Chimpanzee Specificity/Sensitivity The antibody binds to human CD4 with a Kd of 1.0 nM and an ED50 of 0.2 ug/ml. Immunogen Domain 1 of human CD4. Product Application Details Applications Applications Flow Cytometry Recommended Dilutions Flow Cytometry Application Notes Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to I or extreme temperature fluctuations. These can lead to irreversible degradation or ecoupling. When staining intracellular targets, specific attention to the fixa and permeabilization steps in your flow protocol may be required. Please con our technical support team at technical@novusbio.com if you have any	Host	Rabbit
SpeciesHuman, ChimpanzeeReactivity NotesHuman, ChimpanzeeSpecificity/SensitivityThe antibody binds to human CD4 with a Kd of 1.0 nM and an ED50 of 0.2 ug/ml.ImmunogenDomain 1 of human CD4.Product Application DetailsFlow CytometryApplicationsFlow CytometryRecommended DilutionsFlow CytometryApplication NotesOptimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to I or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixa and permeabilization steps in your flow protocol may be required. Please com our technical support team at technical@novusbio.com if you have any	Gene ID	920
Reactivity NotesHuman, ChimpanzeeSpecificity/SensitivityThe antibody binds to human CD4 with a Kd of 1.0 nM and an ED50 of 0.2 ug/ml.ImmunogenDomain 1 of human CD4.Product Application DetailsFlow CytometryApplicationsFlow CytometryRecommended DilutionsFlow CytometryApplication NotesOptimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to I or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixa and permeabilization steps in your flow protocol may be required. Please com our technical support team at technical@novusbio.com if you have any	Gene Symbol	CD4
Specificity/SensitivityThe antibody binds to human CD4 with a Kd of 1.0 nM and an ED50 of 0.2 ug/ml.ImmunogenDomain 1 of human CD4.Product Application DetailsFlow CytometryApplicationsFlow CytometryRecommended DilutionsFlow CytometryApplication NotesOptimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to I or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixation decoupling. When staining intracellular targets, specific attention to the fixation our technical support team at technical@novusbio.com if you have any	Species	Human, Chimpanzee
ug/ml.ImmunogenDomain 1 of human CD4.Product Application DetailsApplicationsFlow CytometryRecommended DilutionsFlow CytometryApplication NotesOptimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to I or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixa and permeabilization steps in your flow protocol may be required. Please com our technical support team at technical@novusbio.com if you have any	Reactivity Notes	Human, Chimpanzee
Product Application Details Applications Flow Cytometry Recommended Dilutions Flow Cytometry Application Notes Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to I or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixe and permeabilization steps in your flow protocol may be required. Please comour technical support team at technical@novusbio.com if you have any	Specificity/Sensitivity	
ApplicationsFlow CytometryRecommended DilutionsFlow CytometryApplication NotesOptimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to I or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixe and permeabilization steps in your flow protocol may be required. Please com- our technical support team at technical@novusbio.com if you have any	Immunogen	Domain 1 of human CD4.
Recommended Dilutions Flow Cytometry Application Notes Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to I or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixe and permeabilization steps in your flow protocol may be required. Please comour technical support team at technical@novusbio.com if you have any	Product Application Details	
Application NotesOptimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to I or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixat and permeabilization steps in your flow protocol may be required. Please com- our technical support team at technical@novusbio.com if you have any	Applications	Flow Cytometry
optimal results using our Tandem dyes, please avoid prolonged exposure to l or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixat and permeabilization steps in your flow protocol may be required. Please con- our technical support team at technical@novusbio.com if you have any	Recommended Dilutions	Flow Cytometry
	Application Notes	optimal results using our Tandem dyes, please avoid prolonged exposure to light or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixation and permeabilization steps in your flow protocol may be required. Please contact





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-52681PECY7

NBP1-19371PEP	CD4 Antibody Blocking Peptide
210-TA-005	TNF-alpha [Unconjugated]
514-CD-050/CF	CD4 [Unconjugated]
M6000B-1	IL-6 [HRP]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-52681PECY7

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

