# **Product Datasheet**

# Complement Factor H Antibody (028B-244.2.10X (10-10)) [Janelia Fluor® 669] NBP2-52743JF669

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-52743JF669

Updated 8/20/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP2-52743JF669



# NBP2-52743JF669

Complement Factor H Antibody (028B-244.2.10X (10-10)) [Janelia Fluor® 669]

0.1 ml
Please see the vial label for concentration. If unlisted please contact technical services.
Store at 4C in the dark.
Monoclonal
028B-244.2.10X (10-10)
0.05% Sodium Azide
IgG1
Janelia Fluor 669
Protein A purified
50mM Sodium Borate
Mouse
3075
CFH
Human
Reacts with Human.
NB100-64342 recognizes human complement factor H which exists in 2 forms. The most common form, of 150 kDa, and the less common form of 43 kDa. Factor H is secreted by the liver into the blood serum. It is important in regulating the complement pathway, preventing unnecessary inflammation which can
damage the host tissue. Complement factor H functions as a cofactor in the inactivation of C3b by factor I. It makes C3b susceptible to cleavage by factor I, resulting in iC3b. Factor H also inhibits the formation of the C3bBb complex (C3 convertase) and increases the rate of dissociation of both C3 convertase and the (C3b)NBB complex (C5 convertase). This prevents these components of the classical and of alternative complement pathways from forming a positive feedback loop. Mutations in factor H are associated with hemolytic-uremic syndrome, age-related macular degeneration, membranoproliferative glomerulonephritis (MPGN) type II and chronic hypocomplementemic nephropathy.
inactivation of C3b by factor I. It makes C3b susceptible to cleavage by factor I, resulting in iC3b. Factor H also inhibits the formation of the C3bBb complex (C3 convertase) and increases the rate of dissociation of both C3 convertase and the (C3b)NBB complex (C5 convertase). This prevents these components of the classical and of alternative complement pathways from forming a positive feedback loop. Mutations in factor H are associated with hemolytic-uremic syndrome, age-related macular degeneration, membranoproliferative glomerulonephritis (MPGN) type II and chronic hypocomplementemic
inactivation of C3b by factor I. It makes C3b susceptible to cleavage by factor I, resulting in iC3b. Factor H also inhibits the formation of the C3bBb complex (C3 convertase) and increases the rate of dissociation of both C3 convertase and the (C3b)NBB complex (C5 convertase). This prevents these components of the classical and of alternative complement pathways from forming a positive feedback loop. Mutations in factor H are associated with hemolytic-uremic syndrome, age-related macular degeneration, membranoproliferative glomerulonephritis (MPGN) type II and chronic hypocomplementemic nephropathy.
inactivation of C3b by factor I. It makes C3b susceptible to cleavage by factor I, resulting in iC3b. Factor H also inhibits the formation of the C3bBb complex (C3 convertase) and increases the rate of dissociation of both C3 convertase and the (C3b)NBB complex (C5 convertase). This prevents these components of the classical and of alternative complement pathways from forming a positive feedback loop. Mutations in factor H are associated with hemolytic-uremic syndrome, age-related macular degeneration, membranoproliferative glomerulonephritis (MPGN) type II and chronic hypocomplementemic nephropathy.  Purified human complement factor H from serum  Sold under license from the Howard Hughes Medical Institute, Janelia Research
inactivation of C3b by factor I. It makes C3b susceptible to cleavage by factor I, resulting in iC3b. Factor H also inhibits the formation of the C3bBb complex (C3 convertase) and increases the rate of dissociation of both C3 convertase and the (C3b)NBB complex (C5 convertase). This prevents these components of the classical and of alternative complement pathways from forming a positive feedback loop. Mutations in factor H are associated with hemolytic-uremic syndrome, age-related macular degeneration, membranoproliferative glomerulonephritis (MPGN) type II and chronic hypocomplementemic nephropathy.  Purified human complement factor H from serum  Sold under license from the Howard Hughes Medical Institute, Janelia Research
inactivation of C3b by factor I. It makes C3b susceptible to cleavage by factor I, resulting in iC3b. Factor H also inhibits the formation of the C3bBb complex (C3 convertase) and increases the rate of dissociation of both C3 convertase and the (C3b)NBB complex (C5 convertase). This prevents these components of the classical and of alternative complement pathways from forming a positive feedback loop. Mutations in factor H are associated with hemolytic-uremic syndrome, age-related macular degeneration, membranoproliferative glomerulonephritis (MPGN) type II and chronic hypocomplementemic nephropathy.  Purified human complement factor H from serum  Sold under license from the Howard Hughes Medical Institute, Janelia Research Campus.  Western Blot, ELISA, Flow Cytometry, Immunohistochemistry,





## Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

### Products Related to NBP2-52743JF669

NBP2-38695PEP Complement Factor H Recombinant Protein Antigen

210-TA-005 TNF-alpha [Unconjugated]
DY4779 Complement Factor H [Biotin]

M6000B-1 IL-6 [HRP]

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-52743JF669

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

