

# Product Datasheet

## FUS Antibody (CL0190) - BSA Free NBP2-52874

Unit Size: 0.1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

[www.novusbio.com](http://www.novusbio.com)



[technical@novusbio.com](mailto:technical@novusbio.com)

### Publications: 3

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:  
[www.novusbio.com/NBP2-52874](http://www.novusbio.com/NBP2-52874)

Updated 3/5/2025 v.20.1

Earn rewards for product  
reviews and publications.

Submit a publication at [www.novusbio.com/publications](http://www.novusbio.com/publications)

Submit a review at [www.novusbio.com/reviews/destination/NBP2-52874](http://www.novusbio.com/reviews/destination/NBP2-52874)



**NBP2-52874**


FUS Antibody (CL0190) - BSA Free

<b>Product Information</b>	
<b>Unit Size</b>	0.1 ml
<b>Concentration</b>	Concentrations vary lot to lot. See vial label for concentration. If unlisted please contact technical services.
<b>Storage</b>	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
<b>Clonality</b>	Monoclonal
<b>Clone</b>	CL0190
<b>Preservative</b>	0.02% Sodium Azide
<b>Isotype</b>	IgG1
<b>Purity</b>	Protein A purified
<b>Buffer</b>	PBS (pH 7.2) and 40% Glycerol
<b>Product Description</b>	
<b>Host</b>	Mouse
<b>Gene ID</b>	2521
<b>Gene Symbol</b>	FUS
<b>Species</b>	Human
<b>Reactivity Notes</b>	Please note that this antibody is reactive to Mouse and derived from the same host, Mouse. Mouse-On-Mouse blocking reagent may be needed for IHC and ICC experiments to reduce high background signal. You can find these reagents under catalog numbers PK-2200-NB and MP-2400-NB. Please contact Technical Support if you have any questions
<b>Immunogen</b>	This antibody was developed using a recombinant protein derived from P35637, with the exact immunogen sequence remaining proprietary.
<b>Product Application Details</b>	
<b>Applications</b>	Western Blot, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Knockdown Validated, Knockout Validated
<b>Recommended Dilutions</b>	Western Blot 1 ug/ml, Immunohistochemistry 1:2500 - 1:5000, Immunocytochemistry/ Immunofluorescence 2-10 ug/ml, Immunohistochemistry-Paraffin 1:2500 - 1:5000, Knockout Validated, Knockdown Validated
<b>Application Notes</b>	For IHC-Paraffin, HIER pH 6 retrieval is recommended. ICC/IF Fixation Permeabilization: Use PFA/Triton X-100.




## Images


**Immunocytochemistry/ Immunofluorescence: FUS Antibody (CL0190) [NBP2-52874]** - Immunofluorescence of HeLa wildtype (WT) and FUS knockout (KO) cells labelled with a green or a far-red fluorescent dye, respectively. WT and KO cells were mixed and plated to a 1:1 ratio on coverslips. Cells were stained with Mouse Anti-FUS Monoclonal Antibody (Catalog #NBP2-52874) at 1:1000, followed by the corresponding Alexa Fluor 555-coupled Secondary Antibody. Acquisition of the green (identification of WT cells), red (antibody staining) and far-red (identification of KO cells) channels was performed. Representative image of the red (grayscale) channel is shown. WT and KO cells are outlined with yellow and magenta dashed line, respectively. Image, protocol and testing courtesy of YCharOS Inc. (ycharos.com).




**Western Blot: FUS Antibody (CL0190) [NBP2-52874]** - Western blot using lysates of HeLa parental cell line (WT) and FUS knockout HeLa cell line (KO), collected in RIPA Buffer. Lysates were prepared using 30 ug of protein and the nitrocellulose membrane was probed with Mouse Anti-FUS Monoclonal Antibody (Catalog #NBP2-52874) at 1:5000 O/N at 4C, followed by peroxidase-conjugated Anti-Mouse Secondary Antibody and ECL detection. A specific band at ~70 kDa was detected in the FUS WT cell line but is not observed in the FUS KO cell line. The Ponceau stained transfer of the blot is shown to confirm equal protein loading. Image, protocol, and testing courtesy of YCharOS Inc. (ycharos.com).




**Immunohistochemistry-Paraffin: FUS Antibody (CL0190) [NBP2-52874]** - Staining in human cerebral cortex and liver tissues. Corresponding FUS RNA-seq data are presented for the same tissues.




**Knockdown Validated: FUS Antibody (CL0190) [NBP2-52874]** - Analysis of extracts from U-251 cells, transfected with: control siRNA, target specific siRNA probe #1, target specific siRNA probe #2, using Anti-FUS monoclonal antibody. Downregulation of antibody signal confirms target specificity. Remaining intensity is presented. Loading control: Anti-GAPDH.




Western Blot: FUS Antibody (CL0190) [NBP2-52874] - Lane 1: Marker [kDa], Lane 2: Human cell line HeLa cytoplasmic fraction , Lane 3: Human cell line HeLa membrane fraction , Lane 4: Human cell line HeLa nuclear fraction , Lane 5: Human cell line HeLa chromatin fraction , Lane 6: Human cell line HeLa cytos




Immunocytochemistry/Immunofluorescence: FUS Antibody (CL0190) [NBP2-52874] - Staining in U2OS cell line with Anti-FUS monoclonal antibody, showing clear nuclear (without nucleoli) staining in green. Microtubule- and nuclear probes are visualized in red and blue respectively (where available). Antibody staining shown in green.




Western Blot: FUS Antibody (CL0190) [NBP2-52874] - Lane 1: Marker [kDa] Lane 2: Human cell line HeLa Lane 3: Human cell line A-431 Lane 4: Human cell line MCF-7 Lane 5: Human cell line U2-OS Lane 6: Human cell line Hep-G2




Immunocytochemistry/Immunofluorescence: FUS Antibody (CL0190) [NBP2-52874] - Staining in A431 cell line with Anti-FUS monoclonal antibody, showing clear nuclear (without nucleoli) staining in green. Microtubule- and nuclear probes are visualized in red and blue respectively (where available). Antibody staining is shown in green.




Immunocytochemistry/Immunofluorescence: FUS Antibody (CL0190) [NBP2-52874] - Staining in HeLa cell line with Anti-FUS monoclonal antibody, showing clear nuclear (without nucleoli) staining in green. Microtubule- and nuclear probes are visualized in red and blue respectively (where available). Antibody staining is shown in green.




Immunocytochemistry/Immunofluorescence: FUS Antibody (CL0190) [NBP2-52874] - Staining in MCF7 cell line with Anti-FUS monoclonal antibody, showing clear nuclear (without nucleoli) staining in green. Microtubule- and nuclear probes are visualized in red and blue respectively (where available). Antibody staining is shown in green.




Immunocytochemistry/Immunofluorescence: FUS Antibody (CL0190) [NBP2-52874] - Staining in U251 cell line with Anti-FUS monoclonal antibody, showing clear nuclear (without nucleoli) staining in green. Microtubule- and nuclear probes are visualized in red and blue respectively (where available). Antibody staining is shown in green.




Immunohistochemistry-Paraffin: FUS Antibody (CL0190) [NBP2-52874] - Staining of human endometrium shows strong nuclear positivity in glandular cells.




Immunohistochemistry-Paraffin: FUS Antibody (CL0190) [NBP2-52874] - Staining of human cerebral cortex shows strong nuclear positivity in neurons.



Immunohistochemistry-Paraffin: FUS Antibody (CL0190) [NBP2-52874] - Staining of human testis shows strong nuclear positivity in cells in seminiferous ducts.



Immunohistochemistry-Paraffin: FUS Antibody (CL0190) [NBP2-52874] - Staining of human liver shows only very weak nuclear positivity in hepatocytes.



## Publications

Alshalfie, W;Fotouhi, M;Ayoubi, R;You, Z;Southern, K;McPherson, PS;Laflamme, C;NeuroSGC/YCharOS/EDDU collaborative group, ; The identification of high-performing antibodies for RNA-binding protein FUS for use in Western Blot, immunoprecipitation, and immunofluorescence F1000Research 2023-06-26 [PMID: 37384305]

Mamontova, EM;Clément, MJ;Sukhanova, MV;Joshi, V;Bouhss, A;Rengifo-Gonzalez, JC;Desforges, B;Hamon, L;Lavrik, OI;Pastré, D; FUS RRM regulates poly(ADP-ribose) levels after transcriptional arrest and PARP-1 activation on DNA damage Cell reports 2023-10-05 [PMID: 37804508]

Vakil PR PARP-1 Activation Directs FUS to DNA Damage Sites to Form PARG-Reversible Compartments Enriched in Damaged DNA Cell Rep 2019-05-07 [PMID: 31067465]



### **Novus Biologicals USA**

10730 E. Briarwood Avenue  
Centennial, CO 80112  
USA  
Phone: 303.730.1950  
Toll Free: 1.888.506.6887  
Fax: 303.730.1966  
nb-customerservice@bio-techne.com

### **Bio-Techne Canada**

21 Canmotor Ave  
Toronto, ON M8Z 4E6  
Canada  
Phone: 905.827.6400  
Toll Free: 855.668.8722  
Fax: 905.827.6402  
canada.inquires@bio-techne.com

### **Bio-Techne Ltd**

19 Barton Lane  
Abingdon Science Park  
Abingdon, OX14 3NB, United Kingdom  
Phone: (44) (0) 1235 529449  
Free Phone: 0800 37 34 15  
Fax: (44) (0) 1235 533420  
info.EMEA@bio-techne.com

### **General Contact Information**

www.novusbio.com  
Technical Support: nb-technical@bio-techne.com  
Orders: nb-customerservice@bio-techne.com  
General: novus@novusbio.com

### **Products Related to NBP2-52874**

---

NBP2-52874PEP	FUS Recombinant Protein Antigen
HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB720-B	Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]
NBP1-97005-0.5mg	Mouse IgG1 Isotype Control (MG1)

---

### **Limitations**

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit [www.novusbio.com/guarantee](http://www.novusbio.com/guarantee)

Earn gift cards/discounts by submitting a review: [www.novusbio.com/reviews/submit/NBP2-52874](http://www.novusbio.com/reviews/submit/NBP2-52874)

Earn gift cards/discounts by submitting a publication using this product:  
[www.novusbio.com/publications](http://www.novusbio.com/publications)

