

# Product Datasheet

**CD27/TNFRSF7 Antibody (LPFS2/1611)**

**NBP2-53196-100ug**

Unit Size: 100 ug

Store at 4C.

[www.novusbio.com](http://www.novusbio.com)



[technical@novusbio.com](mailto:technical@novusbio.com)

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:  
[www.novusbio.com/NBP2-53196](http://www.novusbio.com/NBP2-53196)

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at [www.novusbio.com/publications](http://www.novusbio.com/publications)

Submit a review at [www.novusbio.com/reviews/destination/NBP2-53196](http://www.novusbio.com/reviews/destination/NBP2-53196)



**NBP2-53196-100ug**

CD27/TNFRSF7 Antibody (LPFS2/1611)

Product Information	
Unit Size	100 ug
Concentration	0.2 mg/ml
Storage	Store at 4C.
Clonality	Monoclonal
Clone	LPFS2/1611
Preservative	0.05% Sodium Azide
Isotype	IgG1 Kappa
Purity	Protein A or G purified
Buffer	10 mM PBS with 0.05% BSA
Target Molecular Weight	120 kDa


Product Description	
Description	200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP2-54582)  Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C.
Host	Mouse
Gene ID	939
Gene Symbol	CD27
Species	Human
Immunogen	Recombinant full-length human CD27/TNFRSF7 protein (Uniprot: P26842)

Product Application Details	
Applications	ELISA, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Protein Array
Recommended Dilutions	Flow Cytometry 1-2 ug/million cells, ELISA, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry-Paraffin 1-2 ug/ml, Protein Array
Application Notes	ELISA: For coating, order antibody without BSA). Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris buffer with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes. Optimal dilution for a specific application should be determined.




## Images


Immunocytochemistry/Immunofluorescence: CD27/TNFRSF7 Antibody (LPFS2/1611) [NBP2-53196] - Immunofluorescence staining of Ramos cells using CD27/TNFRSF7 Antibody (LPFS2/1611) followed by goat anti-Mouse IgG conjugated to CF488 (green). Nuclei are stained with Red Dot.




Immunohistochemistry-Paraffin: CD27/TNFRSF7 Antibody (LPFS2/1611) [NBP2-53196] - Formalin-fixed, paraffin-embedded human Colon stained with CD27/TNFRSF7 Antibody (LPFS2/1611).




Flow Cytometry: CD27/TNFRSF7 Antibody (LPFS2/1611) [NBP2-53196] - Flow Cytometric Analysis of Ramos cells using CD27/TNFRSF7 Mouse Monoclonal Antibody (LPFS2/1611) followed by goat anti-Mouse IgG-CF488 (Blue); Isotype Control (Red).




Immunohistochemistry-Paraffin: CD27/TNFRSF7 Antibody (LPFS2/1611) [NBP2-53196] - Formalin-fixed, paraffin-embedded human Tonsil stained with CD27 Monoclonal Antibody (LPFS2/1611).




Immunohistochemistry-Paraffin: CD27/TNFRSF7 Antibody (LPFS2/1611) [NBP2-53196] - Formalin-fixed, paraffin-embedded human Tonsil stained with CD27/TNFRSF7 Antibody (LPFS2/1611).



Immunohistochemistry-Paraffin: CD27/TNFRSF7 Antibody (LPFS2/1611) [NBP2-53196] - Formalin-fixed, paraffin-embedded human Spleen stained with CD27/TNFRSF7 Antibody (LPFS2/1611).



Protein Array: CD27/TNFRSF7 Antibody (LPFS2/1611) [NBP2-53196] - Analysis of Protein Array containing more than 19,000 full-length human proteins using CD27/TNFRSF7 Antibody (LPFS2/1611) Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD?s) above the mean value of all signals generated on that array. If targets on HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD?s) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5.





### **Novus Biologicals USA**

10730 E. Briarwood Avenue  
Centennial, CO 80112  
USA  
Phone: 303.730.1950  
Toll Free: 1.888.506.6887  
Fax: 303.730.1966  
nb-customerservice@bio-techne.com

### **Bio-Techne Canada**

21 Canmotor Ave  
Toronto, ON M8Z 4E6  
Canada  
Phone: 905.827.6400  
Toll Free: 855.668.8722  
Fax: 905.827.6402  
canada.inquires@bio-techne.com

### **Bio-Techne Ltd**

19 Barton Lane  
Abingdon Science Park  
Abingdon, OX14 3NB, United Kingdom  
Phone: (44) (0) 1235 529449  
Free Phone: 0800 37 34 15  
Fax: (44) (0) 1235 533420  
info.EMEA@bio-techne.com

### **General Contact Information**

www.novusbio.com  
Technical Support: nb-technical@bio-techne.com  
Orders: nb-customerservice@bio-techne.com  
General: novus@novusbio.com

### **Products Related to NBP2-53196-100ug**

---

HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB720-B	Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]
NBP1-43319-0.5mg	Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1)
NBP2-38434PEP	CD27/TNFRSF7 Recombinant Protein Antigen

---

### **Limitations**

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit [www.novusbio.com/guarantee](http://www.novusbio.com/guarantee)

Earn gift cards/discounts by submitting a review: [www.novusbio.com/reviews/submit/NBP2-53196](http://www.novusbio.com/reviews/submit/NBP2-53196)

Earn gift cards/discounts by submitting a publication using this product:  
[www.novusbio.com/publications](http://www.novusbio.com/publications)

