

Product Datasheet

Desmin Antibody (DES/1711) NBP2-53383-100ug

Unit Size: 100 ug

Store at 4C.

www.novusbio.com



technical@novusbio.com

Publications: 6

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-53383

Updated 10/23/2024 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-53383



NBP2-53383-100ug

Desmin Antibody (DES/1711)

Product Information	
Unit Size	100 ug
Concentration	0.2 mg/ml
Storage	Store at 4C.
Clonality	Monoclonal
Clone	DES/1711
Preservative	0.05% Sodium Azide
Isotype	IgG1 Kappa
Purity	Protein A or G purified
Buffer	10 mM PBS with 0.05% BSA
Target Molecular Weight	52 kDa

Product Description	
Description	200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP2-54503) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C.
Host	Mouse
Gene ID	1674
Gene Symbol	DES
Species	Human, Mouse, Rat, Chicken, Hamster
Marker	Muscle Cell Marker
Specificity/Sensitivity	Cytoskeletal intermediate filaments (IFs) constitute a diverse group of proteins that are expressed in a highly tissue-specific manner. IFs are constructed from two-chain -helical coiled-coil molecules arranged on an imperfect helical lattice, and have been widely used as markers for distinguishing individual cell types within a tissue and identifying the origins of metastatic tumors. Vimentin is an IF general marker of cells originating in the mesenchyme. Vimentin and Desmin, a related class III IF, are both expressed during skeletal muscle development. Desmin, a 469 amino acid protein found near the Z line in sarcomeres, is expressed more frequently in adult differentiated state tissues. Anti-desmin detects cells of normal smooth, skeletal, and cardiac muscles. Antibody reacts with leiomyomas, leiomyosarcoma, rhabdomyomas, rhabdomyosarcoma, and perivascular cells of glomus tumors of the skin.
Immunogen	Recombinant human desmin protein (Uniprot: P17661)

Product Application Details	
Applications	Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin
Recommended Dilutions	Western Blot 1-2 ug/ml, Flow Cytometry 1-2 ug/million cells, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry-Paraffin 1-2 ug/ml




Application Notes

Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes. Optimal dilution for a specific application should be determined.

Images

Immunohistochemistry-Paraffin: Desmin Antibody (DES/1711) [NBP2-53383] - Formalin-fixed, paraffin-embedded human Uterus stained with Desmin Monoclonal Antibody (DES/1711)

**Publications**

Dubner AM, Lu S, Jolly AJ et al. Smooth muscle-derived adventitial progenitor cells promote key cell type transitions controlling plaque stability in atherosclerosis in a Klf4-dependent manner bioRxiv 2023-07-23 [PMID: 37503181] (Flow Cytometry, Mouse)

Hu K, Palmieri E, Samnuan K Et al. Generalized Modules for Membrane Antigens (GMMA), an outer membrane vesicle-based vaccine platform, for efficient viral antigen delivery J Extracell Vesicles 2022-11-15 [PMID: 36377074] (FLOW, Mouse)

Details:

Citation using the Alexa Fluor 647 version of this antibody.

Piera L, Szymanski J, Juszcak M et al. Histamine is involved in the regulation of collagen content in cultured heart myofibroblasts via H(2), H(3) and H(4) histamine receptors Biomed Rep 2021-08-01 [PMID: 34276989] (Flow)

Details:

Citation using the Azide and BSA Free format of this antibody.

Rajani SF, Faghihi M, Imani Post-infarct morphine treatment reduces apoptosis and myofibroblast density in a rat model of cardiac ischemia-reperfusion Eur J Pharmacol 2020-09-25 [PMID: 32976827] (FLOW, Rat)

Details:

Citation using the Azide and BSA Free version of this antibody.

Wolak M, Bojanowska E, Staszewska T et al. Histamine augments collagen content via H1 receptor stimulation in cultures of myofibroblasts taken from wound granulation tissue Mol Cell Biochem 2020-11-23 [PMID: 33230787] (FLOW)

Miyasaka Y, Murakami K, Ito K et al. Condensed desmin and actin cytoskeletal communication in lipid droplets Cytoskeleton (Hoboken) 2019-10-18 [PMID: 31626391] (WB)



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-53383-100ug

HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB720-B	Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]
NBP1-43319-0.5mg	Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1)
H00001674-Q01-10ug	Recombinant Human Desmin GST (N-Term) Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-53383

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

