Product Datasheet

ACTH Antibody (CLIP/1449) - C-terminus - Azide and BSA Free NBP2-54324-100ug

Unit Size: 100 ug

Store at -20 to -80C. Avoid freeze-thaw cycles.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-54324

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-54324

NBP2-54324-100ug


ACTH Antibody (CLIP/1449) - C-terminus - Azide and BSA Free

WITH BSA & azide (NBP2-53161). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Mouse Gene ID 5443 Gene Symbol POMC Species Human Specificity/Sensitivity ACTH (same as Corticotropin) is a 39 amino acid active peptide produced by anterior pituitary. This monoclonal antibody is specific to CLIP (aa25-39 of ACTH); does not react with Synacthen (aa1-24 of ACTH). POMC (proopiomelanocortin or corticotropin-lipotropin) is a 267 amino acid polypeptide hormone precursor that goes through extensive, tissue-specific posttranslatic processing by convertases. POMC is cleaved into ten hormone chains name MSH, CLIP (corticotropin-like intermediary peptide), Lipotropin-beta, Lipotrop gamma, beta-endorphin and Met-enkephalin. ACTH is also produced by cells immune system (T-cells, B-cells, and macrophages) in response to stimuli associated with stress. Anti-ACTH is a useful marker in classification of pituit tumors and the study of pituitary disease. It reacts with ACTH-producing cells (corticotrophs). It also may react with other tumors (e.g. some small cell	, , , , , , , , , , , , , , , , , , ,		
Concentration 1 mg/ml Storage Store at -20 to -80C. Avoid freeze-thaw cycles. Clonality Monoclonal Clone CLIP/1449 Preservative No Preservative Isotype IgG1 Kappa Purity Protein A or G purified Buffer 10 mM PBS Product Description To mg/ml of antibody purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WiTH BSA & azide (NBP2-53161). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Mouse Gene ID 5443 Gene Symbol POMC Specificity/Sensitivity ACTH (same as Corticotropin) is a 39 amino acid active peptide produced by anterior pituitary. This monoclonal antibody is specific to CLIP (aa25-39 of ACTH); does not react with Synacthen (aa1-24 of ACTH); POMC (proopiomelanocottin or corticotropin-lipotropin) is a 267 amino acid polypeptide hormone precursor that goes through extensive, tissue-specific posttranslatic processing by convertases. POMC is cleaved into ten hormone chains name NPP, ACTH, alpha-MSH (Melanocyte Stimulating Hormone), beta-MSH, gar MSH, CLIP (corticotroph-like intermediary peptide), Lipotroph-beta, Lipotrop gamma, beta-endorphin and Met-enkephalin. ACTH is also produced by cells immune system (T-cells. B-cells, and macrophages) in response to stimuli associated with stress. Anti-ACTH is a u	Product Information		
Storage Store at -20 to -80C. Avoid freeze-thaw cycles. Clonality Monoclonal Clone CLIP/1449 Preservative No Preservative Isotype IgG1 Kappa Purity Protein A or G purified Buffer 10 mM PBS Product Description 1.0 mg/ml of antibody purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WITH BSA & azide (NBP2-53161). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. -80C. Host Mouse Gene ID 5443 Gene Symbol POMC Species Human Specificity/Sensitivity ACTH (same as Corticotropin) is a 39 amino acid active peptide produced by anterior pituitary. This monoclonal antibody is specific to CLIP (aa25-39 of ACTH); does not react with Synacthen (aa1-24 of ACTH). POMC (pro-opiomelanocortin or corticotropin-lipotropin) is a 267 amino acid polypeptide hormone precursor that goes through extensive, itssue-specific posttranslatic processing by convertases. POMC is cleaved into ten hormone chains name NPP, ACTH, alpha-MSH (Melanocyte Stimulating Hormone), beta-MSH, gam MSH, CLIP (corticotropin-like intermediary peptide). Lipotropin-beta,	Unit Size	100 ug	
Clonality Monoclonal Clone CLIP/1449 Preservative No Preservative Isotype IgG1 Kappa Purity Protein A or G purified Buffer 10 mM PBS Product Description Description Description 1.0 mg/ml of antibody purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WITH BSA & azide (NBP2-53161). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Mouse Gene ID 5443 Gene Symbol POMC Species Human Specificity/Sensitivity ACTH (same as Corticotropin) is a 39 amino acid active peptide produced by anterior pituitary. This monoclonal antibody is specific to CLIP (aa25-39 of ACTH); does not react with Synacthen (aa1-24 of ACTH). POMC (pro- opiomelancortin or corticotropin-liptorpin) is a 267 amino acid polypeptide hormone precursor that goes through extensive, tissue-specific posttranslatic processing by convertases. POMC is clawed into ten hormone chains name MPP, ACTH, alpha-MSH (Melancy peptide). Lipotropin-beta, Lipotropi gramma, beta-endorphin and Met-enkephalin. ACTH is also produced by cells immune system (T-cells, B-cells, and macrophage) in response to stimuli associated with stress. Anti-ACTH is a useful marker in classification of pituit turrors and the study of pituitary disease. It reacts with ACTH-producing cell	Concentration	1 mg/ml	
Clone CLIP/1449 Preservative No Preservative Isotype IgG1 Kappa Purity Protein A or G purified Buffer 10 mM PBS Product Description 1.0 mg/ml of antibody purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WITH BSA & azide (NBP2-53161). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Mouse Gene ID 5443 Gene Symbol POMC Specificity/Sensitivity ACTH (same as Corticotropin) is a 39 amino acid active peptide produced by anterior pituitary. This monoclonal antibody is specific to CLIP (aa25-39 of ACTH); does not react with Synacthen (aa1-24 of ACTH). POMC (pro- opiomelanocortin or corticotropin-lipotropin). Specificity/Sensitivity ACTH (same as Corticotropin-lipotropin). Short H, alpha-MSH (Melanocyte Stimulating Hormone), beta-MSH, gam MSH, CLIP (corticorphin). Superide hormone, precursor that goes through extensive, tissue-specific posttranslatic processing by convertases. POMC is cleaved into the normone chains name NSH, CLIP (corticotrophin ind Met-enkephalin. ACTH) is also produced by cells immune system (T-cells, B-cells, and macrophages) in response to stimuli associated with stress. Anti-ACTH is a useful marker in classification of pituit turnors and the study of pituitary disease. It reacts with ACTH-producing cells (corricotrophs).	Storage	Store at -20 to -80C. Avoid freeze-thaw cycles.	
Preservative No Preservative Isotype IgG1 Kappa Purity Protein A or G purified Buffer 10 mM PBS Product Description 1.0 mg/ml of antibody purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WITH BSA & azide (NBP2-53161). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Mouse Gene ID 5443 Species Human Specificity/Sensitivity ACTH (same as Corticotropin) is a 39 amino acid active peptide produced by anterior pituitary. This monoclonal antibody is specific to CLIP (aa25-39 of ACTH); does not react with Synacthen (aa1-24 of ACTH). POMC (pro- opiomelanocortin or corticotropin-lipotropin) is a 267 amino acid polypeptide hormone precursor that goes through extensive, tissue-specific posttranslatic processing by convertases. POMC is cleaved into ten hormone chains name NPP, ACTH, alpha-MSH (Melanocyte Stimulating Hormone), beta-MSH, gar MSH, CLIP (corticotropin-like intermediary peptide), Lipotropin-beta, Lipotrop gamma, beta-endorphin and Met-enkephalin. ACTH is also produced by cells immune system (T-cells, B-cells, and macrophages) in response to stimuli associated with stress. Anti-ACTH is a useful marker in classification of pitulit tumors and the study of pituitary disease. It reacts with ACTH-producing cells (corticotrophs). It also may react with other tumors (e.g. some small cell	Clonality	Monoclonal	
Isotype IgG1 Kappa Purity Protein A or G purified Buffer 10 mM PBS Product Description 1.0 mg/ml of antibody purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WITH BSA & azide (NBP2-53161). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Mouse Gene ID 5443 Species Human Specificity/Sensitivity ACTH (same as Corticotropin) is a 39 amino acid active peptide produced by anterior pituitary. This monoclonal antibody is specific to CLIP (aa25-39 of ACTH); does not react with Synacthen (aa1-24 of ACTH). POMC (pro- opiomelanocortin or corticotropin-lipotropin) is a 267 amino acid polypeptide hormone precursor that goes through extensive, tissue-specific posttranslatic processing by convertases. POMC is cleaved into ten hormone chains name NPP, ACTH, alpha-MSH (Melanocyte Stimulating Hormone), beta-MSH, gar MSH, CLIP (corticotropin-like intermediary peptide), Lipotropin-beta, Lipotrop garma, beta-endorphin and Met-enkephalin. ACTH is also produced by cells immune system (T-cells, B-cells, and macrophages) in response to stimuli associated with stress. Anti-ACTH is a useful marker in classification of pitulit tumors and the study of pituitary disease. It reacts with ACTH-producing cells (corticotrophs).It also may react with other tumors (e.g. some small cell	Clone	CLIP/1449	
Purity Protein A or G purified Buffer 10 mM PBS Product Description 1.0 mg/ml of antibody purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WITH BSA & azide (NBP2-53161). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Mouse Gene ID 5443 Gene Symbol POMC Species Human Specificity/Sensitivity ACTH (same as Corticotropin) is a 39 amino acid active peptide produced by anterior pituitary. This monoclonal antibody is specific to CLIP (aa25-39 of ACTH); does not react with Synacthen (aa1-24 of ACTH). POMC (pro- opiomelanocortin or corticotropin) is a 267 amino acid polypeptide hormone precursor that goes through extensive, tissue-specific posttranslatic processing by convertases. POMC is cleaved into ten hormone chains name NPP, ACTH, alpha-MSH (Melanocyte Stimulating Hormone), beta-MSH, gan MSH, CLIP (corticotropin-like intermediary peptide), Lipotropin-beta, Lipotrop gamma, beta-endorphin and Met-enkephalin. ACTH is also produced by cells immune system (T-cells, B-cells, and macrophages) in response to stimuli associated with stress. Anti-ACTH is a useful marker in classification of pituli tumors and the study of pituitary disease. It reacts with ACTH-producing cells (corticotrophs). It also may react with other tumors (e.g. some small cell	Preservative	No Preservative	
Buffer 10 mM PBS Product Description 1.0 mg/ml of antibody purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WiTH BSA & azide (NBP2-53161). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Mouse Gene ID 5443 Gene Symbol POMC Species Human Specificity/Sensitivity ACTH (same as Corticotropin) is a 39 amino acid active peptide produced by anterior pituitary. This monoclonal antibody is specific to CLIP (aa25-39 of ACTH); does not react with Synacthen (aa1-24 of ACTH). POMC (pro-opiomelanccortin or corticotropin-lipotropin) is a 267 amino acid polypeptide hormone precursor that goes through extensive, tissue-specific postranslatic processing by convertases. POMC is cleaved into ten hormone chains name NPP, ACTH, alpha-MSH (Melanocyte Stimulating Hormone), beta-MSH, gam MSH, CLIP (corticotropin-like intermediary peptide), Lipotropin-beta, Lipotrop gamma, beta-endorphin and Met-enkephalin. ACTH is also produced by cells immune system (T-cells, B-cells, and macrophages) in response to stimuli associated with stress. Anti-ACTH is a useful marker in classification of pituit tumors and the study of pituitary disease. It reacts with ACTH-producing cells (corticotrophs). It also may react with other tumors (e.g. some small cell	Isotype	IgG1 Kappa	
Product Description 1.0 mg/ml of antibody purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WITH BSA & azide (NBP2-53161). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Mouse Gene ID 5443 Gene Symbol POMC Species Human Specificity/Sensitivity ACTH (same as Corticotropin) is a 39 amino acid active peptide produced by anterior pituitary. This monoclonal antibody is specific to CLIP (aa25-39 of ACTH); does not react with Synacthen (aa1-24 of ACTH). POMC (pro- opiomelanocortin or corticotropin-lipotropin) is a 267 amino acid polypeptide hormone precursor that goes through extensive, tissue-specific posttranslatic processing by convertases. POMC is cleaved into ten hormone chains name NPP, ACTH, alpha-MSH (Melanocyte Stimulating Hormone), beta-MSH, gar MSH, CLIP (corticotropin-like intermediary peptide), Lipotropin-beta, Lipotrop garma, beta-endorphin and Met-enkephalin. ACTH is also produced by cells immune system (T-cells, B-cells, and macrophages) in response to stimuli associated with stress. Anti-ACTH is a useful marker in classification of pituit tumors and the study of pituitary disease. It reacts with ACTH-producing cells (corticotrophs).It also may react with other tumors (e.g. some small cell	Purity	Protein A or G purified	
Description 1.0 mg/ml of antibody purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WITH BSA & azide (NBP2-53161). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Mouse Gene ID 5443 Gene Symbol POMC Species Human Specificity/Sensitivity ACTH (same as Corticotropin) is a 39 amino acid active peptide produced by anterior pituitary. This monoclonal antibody is specific to CLIP (aa25-39 of ACTH); does not react with Synacthen (aa1-24 of ACTH). POMC (pro- opiomelanocortin or corticotropin-lipotropin) is a 267 amino acid polypeptide hormone precursor that goes through extensive, tissue-specific posttranslatic processing by convertases. POMC is cleaved into ten hormone chains name NPP, ACTH, alpha-MSH (Melanocyte Stimulating Hormone), beta-MSH, gam MSH, CLIP (corticotropin-like intermediary peptide), Lipotropin-beta, Lipotrop gamma, beta-endorphin and Met-enkephalin. ACTH is also produced by cells immune system (T-cells, B-cells, and macrophages) in response to stimuli associated with stress. Anti-ACTH is a useful marker in classification of pituit turnors and the study of pituitary disease. It reacts with ACTH-producing cells (corticotrophs).It also may react with other turnors (e.g. some small cell	Buffer	10 mM PBS	
Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WITH BSA & azide (NBP2-53161). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Mouse Gene ID 5443 Gene Symbol POMC Species Human Specificity/Sensitivity ACTH (same as Corticotropin) is a 39 amino acid active peptide produced by anterior pituitary. This monoclonal antibody is specific to CLIP (aa25-39 of ACTH); does not react with Synacthen (aa1-24 of ACTH). POMC (pro-opiomelanocortin or corticotropin-lipotropin) is a 267 amino acid polypeptide hormone precursor that goes through extensive, tissue-specific posttranslatic processing by convertases. POMC is cleaved into ten hormone chains name NPP, ACTH, alpha-MSH (Melanocyte Stimulating Hormone), beta-MSH, gam MSH, CLIP (corticotropin-like intermediary peptide). Lipotropin-beta, Lipotrop gamma, beta-endorphin and Met-enkephalin. ACTH is also produced by cells immune system (T-cells, B-cells, and macrophages) in response to stimuli associated with stress. Anti-ACTH is a useful marker in classification of pituit tumors and the study of pituitary disease. It reacts with ACTH-producing cells (corticotrophs).It also may react with other tumors (e.g. some small cell	Product Description		
Gene ID5443Gene SymbolPOMCSpeciesHumanSpecificity/SensitivityACTH (same as Corticotropin) is a 39 amino acid active peptide produced by anterior pituitary. This monoclonal antibody is specific to CLIP (aa25-39 of ACTH); does not react with Synacthen (aa1-24 of ACTH). POMC (pro- opiomelanocortin or corticotropin-lipotropin) is a 267 amino acid polypeptide hormone precursor that goes through extensive, tissue-specific posttranslatic processing by convertases. POMC is cleaved into ten hormone chains name NPP, ACTH, alpha-MSH (Melanocyte Stimulating Hormone), beta-MSH, gam MSH, CLIP (corticotropin-like intermediary peptide), Lipotropin-beta, Lipotrop gamma, beta-endorphin and Met-enkephalin. ACTH is also produced by cells immune system (T-cells, B-cells, and macrophages) in response to stimuli associated with stress. Anti-ACTH is a useful marker in classification of pituit tumors and the study of pituitary disease. It reacts with ACTH-producing cells (corticotrophs).It also may react with other tumors (e.g. some small cell		Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WITH BSA & azide (NBP2-53161). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to	
Gene Symbol POMC Species Human Specificity/Sensitivity ACTH (same as Corticotropin) is a 39 amino acid active peptide produced by anterior pituitary. This monoclonal antibody is specific to CLIP (aa25-39 of ACTH); does not react with Synacthen (aa1-24 of ACTH). POMC (pro-opiomelanocortin or corticotropin-lipotropin) is a 267 amino acid polypeptide hormone precursor that goes through extensive, tissue-specific posttranslation processing by convertases. POMC is cleaved into ten hormone, beta-MSH, gam MSH, CLIP (corticotropin-like intermediary peptide), Lipotropin-beta, Lipotrop gamma, beta-endorphin and Met-enkephalin. ACTH is also produced by cells immune system (T-cells, B-cells, and macrophages) in response to stimuli associated with stress. Anti-ACTH is a useful marker in classification of pituitat tumors and the study of pituitary disease. It reacts with ACTH-producing cells (corticotrophs).It also may react with other tumors (e.g. some small cell	Host	Mouse	
SpeciesHumanSpecificity/SensitivityACTH (same as Corticotropin) is a 39 amino acid active peptide produced by anterior pituitary. This monoclonal antibody is specific to CLIP (aa25-39 of ACTH); does not react with Synacthen (aa1-24 of ACTH). POMC (pro- opiomelanocortin or corticotropin-lipotropin) is a 267 amino acid polypeptide hormone precursor that goes through extensive, tissue-specific posttranslation processing by convertases. POMC is cleaved into ten hormone chains name NPP, ACTH, alpha-MSH (Melanocyte Stimulating Hormone), beta-MSH, gam MSH, CLIP (corticotropin-like intermediary peptide), Lipotropin-beta, Lipotrop gamma, beta-endorphin and Met-enkephalin. ACTH is also produced by cells immune system (T-cells, B-cells, and macrophages) in response to stimuli associated with stress. Anti-ACTH is a useful marker in classification of pituitat tumors and the study of pituitary disease. It reacts with ACTH-producing cells (corticotrophs).It also may react with other tumors (e.g. some small cell	Gene ID	5443	
Specificity/Sensitivity ACTH (same as Corticotropin) is a 39 amino acid active peptide produced by anterior pituitary. This monoclonal antibody is specific to CLIP (aa25-39 of ACTH); does not react with Synacthen (aa1-24 of ACTH). POMC (pro-opiomelanocortin or corticotropin-lipotropin) is a 267 amino acid polypeptide hormone precursor that goes through extensive, tissue-specific posttranslatic processing by convertases. POMC is cleaved into ten hormone chains name NPP, ACTH, alpha-MSH (Melanocyte Stimulating Hormone), beta-MSH, gam MSH, CLIP (corticotropin-like intermediary peptide), Lipotropin-beta, Lipotrop gamma, beta-endorphin and Met-enkephalin. ACTH is also produced by cells immune system (T-cells, B-cells, and macrophages) in response to stimuli associated with stress. Anti-ACTH is a useful marker in classification of pituitat tumors and the study of pituitary disease. It reacts with ACTH-producing cells (corticotrophs). It also may react with other tumors (e.g. some small cell	Gene Symbol	POMC	
anterior pituitary. This monoclonal antibody is specific to CLIP (aa25-39 of ACTH); does not react with Synacthen (aa1-24 of ACTH). POMC (pro- opiomelanocortin or corticotropin-lipotropin) is a 267 amino acid polypeptide hormone precursor that goes through extensive, tissue-specific posttranslatic processing by convertases. POMC is cleaved into ten hormone chains name NPP, ACTH, alpha-MSH (Melanocyte Stimulating Hormone), beta-MSH, gam MSH, CLIP (corticotropin-like intermediary peptide), Lipotropin-beta, Lipotrop gamma, beta-endorphin and Met-enkephalin. ACTH is also produced by cells immune system (T-cells, B-cells, and macrophages) in response to stimuli associated with stress. Anti-ACTH is a useful marker in classification of pituitat tumors and the study of pituitary disease. It reacts with ACTH-producing cells (corticotrophs).It also may react with other tumors (e.g. some small cell	Species	Human	
		ACTH); does not react with Synacthen (aa1-24 of ACTH). POMC (pro- opiomelanocortin or corticotropin-lipotropin) is a 267 amino acid polypeptide hormone precursor that goes through extensive, tissue-specific posttranslational processing by convertases. POMC is cleaved into ten hormone chains named NPP, ACTH, alpha-MSH (Melanocyte Stimulating Hormone), beta-MSH, gamma- MSH, CLIP (corticotropin-like intermediary peptide), Lipotropin-beta, Lipotropin- gamma, beta-endorphin and Met-enkephalin. ACTH is also produced by cells of immune system (T-cells, B-cells, and macrophages) in response to stimuli associated with stress. Anti-ACTH is a useful marker in classification of pituitary tumors and the study of pituitary disease. It reacts with ACTH-producing cells (corticotrophs).It also may react with other tumors (e.g. some small cell carcinomas of the lung) causing paraneoplastic syndromes by secreting ACTH. '	
ImmunogenSynthetic peptide corresponding to aa25-39 of human ACTH (NGAEDESAEAFPLEF) (Uniprot: P01189)			
Product Application Details	Product Application Details		
Applications ELISA, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-read	Applications	ELISA, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready	
Recommended Dilutions ELISA 1:100 - 1:2000, Immunohistochemistry, Immunohistochemistry-Paraffi 0.5 - 1.0 ug/ml, CyTOF-ready 0.5 - 1.0 ug/ml, CyTOF-ready		ELISA 1:100 - 1:2000, Immunohistochemistry, Immunohistochemistry-Paraffin 0.5 - 1.0 ug/ml, CyTOF-ready	
		Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	



Images

Immunohistochemistry-Paraffin: ACTH Antibody (CLIP/1449) [NBP2-54324] - Formalin-fixed, paraffin-embedded Human Pituitary stained with ACTH Monoclonal Antibody (CLIP/1449).



www.novusbio.com



technical@novusbio.com



Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-54324-100ug

210-TA-005	TNF-alpha [Unconjugated]
NBP1-43319-0.5mg	Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1)
NB720-B	Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]
HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-54324

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

