Product Datasheet

CD45RA Antibody (PTPRC/1131) [PE/Atto594] NBP2-54336PEATT594

Unit Size: 0.1 ml

Store at 4C in the dark. Do not freeze.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-54336PEATT594

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-54336PEATT594



NBP2-54336PEATT594

CD45RA Antibody (PTPRC/1131) [PE/Atto594]

services. Storage Store at 4C in the dark. Do not freeze. Clonality Monoclonal Clone PTPRC/1131 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate PE/Atto594 Purity Protein A or G purified Buffer PBS Product Description House Gene ID 5788 Gene Symbol PTPRC Species Human Marker Leucocyte Marker Specificity/Sensitivity Recognizes a protein of 205kDa-220kDa, identified as CD45RA. CD45RA is soforms of the human leukocyte common antigen (CD45). Human CD45 contains three exons which encode peptide segments designated A, B and C tochsRA is soforms. ABC, AB, BC, B and O. This antibody reacts with ABC and BC isoforms. CD45RA is sopressing CD45RA are memory coldsRA is protessing CD45RA are memory coldsRA are new or vign 1 r-cells. T-cells. T-cells. T-cells. T-cells expressing CD45RO are memory coldsRA are new or vign 1 r-cells. Tocells are prossing CD45RO are memory coldsRA are naive or vign 1 r-cells. T-cells are prossing CD45RO are memory coldsRA are naive or vign 1 r-cells. T-cells. Trecells expressing CD45RO are memory coldsRA are naive or vign 1 r-cells. T-cells. T-cells. T-cells expressing CD45RO are memory coldsRA are naive or vign 1 r-cells. T-cells are sories with ABC and BC isoforms. CD45RA are use or vign 1 r-cells. T-cells are sories not altotor core sories not alto tor core sories not the dare c			
Concentration Please see the vial label for concentration. If unlisted please contact technica services. Storage Store at 4C in the dark. Do not freeze. Clonality Monoclonal Clone PTPRC/1131 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate PE/Atto594 Purity Protein A or G purified Buffer PBS Product Description Mouse Gene ID 5788 Gene Symbol PTPRC Species Human Marker Leucocyte Marker Species Human Marker Leucocyte Marker Specificity/Sensitivity Recognizes a protein of 205kDa-220kDa, identified as CD45RA. CD45RA is isoforms. ABC, AB, BC, B and O. This antibody reacts with ABC and BC isoforms. ABC, AB, BC, B and O. This antibody reacts with ABC and BC isoforms. ABC, AB, BC, B and O. This antibody reacts with ABC and BC isoforms. CD45RA are naive or trign T-cells. Trells expressing CD45RA are naive or trign T-cells. This monoclonal antiboc useful in study on the subpopulation of CD44FR - croB4 T-cells, 50% peripheral CD4F - t-cells. A codefine complementary. predominantly non- overlapping populations of resting peripheral T-cells This monoclonal antiboc useful in study on the subopopulation of che mistry/ Immunofluorescence, Immun	Product Information		
services. Storage Store at 4C in the dark. Do not freeze. Clonality Monoclonal Clone PTPRC/1131 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate PE/Atto594 Purity Protein A or G purified Buffer PBS Product Description Mouse Gene ID 5788 Gene Symbol PTPRC Species Human Marker Leucocyte Marker Species Human Marker Leucocyte Marker Specificity/Sensitivity Recognizes a protein of 205KDa-220kDa, identified as CD45RA, CD45RA is isoforms of the human leukocyte common antigen (CD45), Human CD45 contains three exons which encode peptide segments designated A, B and C respectively. The differential splicing of the exons generates at least five isoforms. ABC, AB, BC, B and O. This antibody reacts with AB cand BC cisoforms. ABC, AB, BC, B and O. This antibody reacts with AB cand BC isoforms. ABC, AB, BC, B and CH ercoells. T-cells. Storesence, Immunofuce to differentiate T-cells True appeciation or the subpopulation of CD45RO 4 ver (D45RA 4 ver). Immunogen Stimulated human leukocytes Product Application Details Meestem Blot, Flow Cyt	Unit Size	0.1 ml	
Clonality Monoclonal Clone PTPRC/1131 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate PE/Atto594 Purity Protein A or G purified Buffer PES Product Description Host Host Mouse Gene Symbol PTPRC Species Human Marker Leucocyte Marker Species Human Marker Leucocyte Marker Specificity/Sensitivity Recognizes a protein of 205kDa-220kDa, identified as CD45RA. CD45RA is isoforms of the human leukocyte common antigen (CD45). Human CD45 contains three exons which encode peptide segments designated A, B and C respectively. The differential splicing of the exons generates at least five isoforms. CD45RA is expressed on 40-50% of peripheral CD4+T-cells, 50% of peripher	Concentration	Please see the vial label for concentration. If unlisted please contact technical services.	
Clone PTPRC/1131 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate PE/Atto594 Purity Protein A or G purified Buffer PBS Product Description Froduct Description Host Mouse Gene ID 5788 Gene Symbol PTPRC Species Human Marker Leucocyte Marker Specificity/Sensitivity Recognizes a protein of 205kDa-220kDa, identified as CD45RA. CD45RA is isoforms of the human leukocyte common antigen (CD45). Human CD45 contains three exons which encode peptide segments designated A, B and C respectively. The differential splicing of the exons generates at least five isoforms. CD45RA is expressing CD45RA are memory cells. CD45RA are is expressing CD45RO are memory cells. CD45RA are or sing peripheral T-cells. T-cells expressing CD45RO are memory cells. CD45RA are of sing peripheral T-cells. T-cells expressing CD45RO are nemory cells. CD45RA are of sing peripheral T-cells. T-cells expressing CD45RO are encory crapping populations of resing peripheral T-cells. T-cells expressing CD45RO are nemory cells. CD45RA are of the subpopulation of CD4+ or CD8+ T-cells. It can especial be used to differentiate T-cell lymphomas (CD45RO +we) from B cell lymphor (D45RA +we). Immunogen Stimulated human leukocytes Product Application Details Mestern Blot, Flow Cytometry, Im	Storage	Store at 4C in the dark. Do not freeze.	
Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate PE/Atto594 Purity Protein A or G purified Buffer PBS Product Description Host Host Mouse Gene ID 5788 Gene Symbol PTPRC Species Human Marker Leucocyte Marker Specificity/Sensitivity Recognizes a protein of 205kDa-220kDa, identified as CD45RA. CD45RA is isoforms of the human leukocyte common antigen (CD45). Human CD45 contains three exons which encode peptide segments designated A, B and C respectively. The differential splicing of the exons generates at least five isoforms. CD45RA cod5RA are naive or virgin T-cells. T-cells softwares. T-cells softwares. CD45RA are naive or virgin T-cells. T-cells spressing CD45RO are memory cells. CD45RA are naive or virgin T-cells. Tr-cells spressing CD45RO are memory cells. CD45RA are naive or virgin T-cells. Tr-cells spressing CD45RO are memory cells. CD45RA are naive or virgin T-cells. Tr-cells spressing CD45RO are memory cells. CD45RA are naive or virgin T-cells. Tr-cells Software are approximately non-overlapping populations of resting peripheral T-cells. This monoclonal antibor useful in study on the subpopulation of CD4+ or CD8+ T-cells. It can especial used to differentiate T-cell lymphomas (CD45RO +ve) from B cell lymphor (CD45RA +ve). Immunogen Stimulated human leukocytes Product Application Details	Clonality	Monoclonal	
Isotype IgG1 Kappa Conjugate PE/Atto594 Purity Protein A or G purified Buffer PBS Product Description Mouse Gene ID 5788 Gene Symbol PTPRC Species Human Marker Leucocyte Marker Specificity/Sensitivity Recognizes a protein of 205kDa-220kDa, identified as CD45RA, CD45RA is isoforms of the human leukocyte common antigen (CD45). Human CD45 contains three exons which encode peptide segments designated A, B and C respectively. The differential splicing of the exons generates at least five isoforms, ABC, AB, BC, B and O. This antibody reacts with ABC and BC contains three exons which encode peptide segments designated A, B and C C respectively. The differential splicing of the exons generates at least five isoforms, ABC, AB, BC, B and O. This antibody reacts with ABC and BC contains three exons which encode peptide segments designated A, B and C C desRA are naive or virgin T-cells. T-cells expressing CD45RO are memory cells. CD45RA are naive or virgin T-cells. T-cells expressing CD45RO are memory cells. CD45RA are naive or virgin T-cells. T-cells expressing CD45RO are memory cells. CD45RA are naive or virgin T-cells. T-cells the cells the septide are special be used to differentiate T-cell lymphomas (CD45RO +ve) from B cell lymphor (CD45RA +ve). Immunogen Stimulated human leukocytes Product Application Details Western Blot, Flow Cytometry, Immunohistochemistry/ Immunofluorescence, Immunohistochemistry. Immunofluorescence, Immunohistochemistry.	Clone	PTPRC/1131	
Conjugate PE/Atto594 Purity Protein A or G purified Buffer PBS Product Description Mouse Gene ID 5788 Gene Symbol PTPRC Species Human Marker Leucocyte Marker Specificity/Sensitivity Recognizes a protein of 205kDa-220kDa, identified as CD45RA. CD45RA is isoforms of the human leukocyte common antigen (CD45). Human CD45 contains three exons which encode peptide segments designated A, B and C respectively. The differential splicing of the exons generates at least five isoforms, ABC, AB, BC, B and O. This antibody reacts with ABC and BC isoforms, ABC, AB, BC, B and O. This antibody reacts with ABC and BC isoforms, CD45RA is respression qD45RO derine complementary, predominantly non-overlapping populations of resting peripheral CD4+ r-cells. This monoclonal antiboc useful in study on the subpopulation of CD4+ or CD45R r-cells. This monoclonal antiboc useful in study on the subpopulation of CD4+ or CD45R r-cells. This monoclonal antiboc useful in study on the subpopulation of CD4+ or CD45R r-cells. This monoclonal antiboc useful in study on the subpopulation of CD4+ or CD45R r-cells. It can especial be used to differentiate T-cell lymphomas (CD45RO +ve) from B cell lymphor (CD45RA +ve). Immunogen Stimulated human leukocytes Product Application Details Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, CyTOF-ready Application Notes Optimal dilution of this antibody should be experimentally deter	Preservative	0.05% Sodium Azide	
Purity Protein A or G purified Buffer PBS Product Description Mouse Gene ID 5788 Gene Symbol PTPRC Species Human Marker Leucocyte Marker Specificity/Sensitivity Recognizes a protein of 205kDa-220kDa, identified as CD45RA. CD45RA is isoforms of the human leukocyte common antigen (CD45). Human CD45 contains three exons which encode peptide segments designated A, B and C respectively. The differential splicing of the exons generates at least five isoforms. CD45R A is expressed on 40-50% of peripheral CD4+ T-cells, 50% peripheral CD8+ T-cells. R-cells and leukemic B-cell lines. T-cells expressing CD45RO are memory cells. CD45RA are naive or virgin T-cells. T-cells expressing CD45RO are memory cells. CD45RA are naive or virgin T-cells. T-cells expressing CD45RO are memory cells. CD45RA are naive or virgin T-cells. T-cells expressing CD45RO are memory cells. CD45RA are naive or virgin T-cells. T-cells expressing CD45RO are memory cells. CD45RA are naive or virgin T-cells. T-cells expressing CD45RO are memory cells. CD45RA are naive or virgin T-cells. T-cells expressing CD45RO are memory cells. UD45RA +ve). Immunogen Stimulated human leukocytes Product Application Details Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, CyTOF-ready Application Notes Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, pelase avoid prolonged exposure to or extreme	Isotype	IgG1 Kappa	
Buffer PBS Product Description Mouse Gene ID 5788 Gene Symbol PTPRC Species Human Marker Leucocyte Marker Specificity/Sensitivity Recognizes a protein of 205kDa-220kDa, identified as CD45RA. CD45RA is isoforms of the human leukocyte common antigen (CD45). Human CD45 contains three exons which encode peptide segments designated A, B and C respectively. The differential splicing of the exons generates at least five isoforms. ABC, AB, BC, B and O. This antibody reacts with ABC and BC isoforms. CD45RA are naive or virgin T-cells. T-cells expressing CD45RO are memory cells. CD45RA are naive or virgin T-cells. T-cells expressing CD45RO are memory cells. CD45RA are naive or virgin T-cells. The monoclonal antibor useful in study on the subpopulation of CD4+ r-cells. This monoclonal antibor useful in study on the subpopulation of CD4+ r-cells. This monoclonal antibor useful in study on the subpopulation of CD4+ r-cells. The monoclonal antibor useful in study on the subpopulation of CD4+ r-cells. The monoclonal antibor useful in study on the subpopulation of CD4+ r-cells. The monoclonal antibor useful in study on the subpopulation of CD4+ r-cells. The monoclonal antibor useful in study on the subpopulation of CD4+ r-cells. The monoclonal antibor useful in study on the subpopulation of CD4+ r-cells. The monoclonal antibor useful in study on the subpopulation of CD4+ r-cells. The monoclonal antibor useful in study on the subpopulation of CD4+ r-cells. The monoclonal antibor useful in study on the subpopulation of the submertity. Immunofluorescence, Immunohistochemistry-Paraffin, CyTOF-ready Applications Western Blot, Fl	Conjugate	PE/Atto594	
Product Description Host Mouse Gene ID 5788 Gene Symbol PTPRC Species Human Marker Leucocyte Marker Specificity/Sensitivity Recognizes a protein of 205kDa-220kDa, identified as CD45RA. CD45RA is isoforms of the human leukocyte common antigen (CD45). Human CD45 contains three exons which encode peptide segments designated A, B and C respectively. The differential splicing of the exons generates at least five isoforms. CD45RA As, BC, B and O. This antibody reacts with ABC and BC isoforms. CD45RA are naive or virgin T-cells. and leukemic B-cell lines. T-cells expressing CD45RA are memory cells. CD45RA are OL45RO dependent of 2045RO are memory cells. CD45RA are to CD45RO dependent of 2045RO are memory cells. CD45RA are to CD45RO dependent of cD45RO exons generates at least five isoforms. CD45RA and CD45RO define complementary, predominantly non-overlapping populations of resting peripheral T-cells. It can especial be used to differentiate T-cell VMphomas (CD45RO +ve) from B cell VMphor (CD45RA +ve). Immunogen Stimulated human leukocytes Product Application Details Mestern Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, CyTOF-ready Application Notes Optimal diultion of this antibody should be experimentally determined. For or externe temperature fluctuations. These can lead to irreversible degradati or decoupling. When staining intracellular targets, specific attention to the fix and permeabilization steps in your flow protocol may be required. Please con our technical support team at technical@novusbio.com	Purity	Protein A or G purified	
Host Mouse Gene ID 5788 Gene Symbol PTPRC Species Human Marker Leucocyte Marker Specificity/Sensitivity Recognizes a protein of 205kDa-220kDa, identified as CD45RA. CD45RA is isoforms of the human leukocyte common antigen (CD45). Human CD45 contains three exons which encode peptide segments designated A, B and C respectively. The differential splicing of the exons generates at least five isoforms. CD45RA is is expressed on 40-50% of peripheral CD4+ T-cells, 50% peripheral CD4+ T-cells, specific antibody reacts with ABC and BC isoforms. CD45RA are D45RA or or virgin T-cells antibody reacts with ABC are memory cells. CD45RA and CD45RO define complementary. predominantly non-overlapping populations of resting peripheral T-cells. This monoclonal antibody useful in study on the subpopulation of CD4+ or CD8+ T-cells. It can especial be used to differentiate T-cell lymphomas (CD45RO +ve) from B cell lymphor (CD45RA +ve). Immunogen Stimulated human leukocytes Product Application Details Western Blot, Flow Cytometry, Immunohistochemistry/ Immunofluorescence, Immunohistochemistry. Immunohistochemistry-Paraffin, CyTOF-ready Application Notes Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to lor exterme temperature fluctuations. These can lead to irreversible degradatio or decoupling, When staining intracellual targets, specific attention to the fix and permeabilization steps in your flow protocol may be required. Please con on ur technical support team at technical @nou	Buffer	PBS	
Gene ID 5788 Gene Symbol PTPRC Species Human Marker Leucocyte Marker Specificity/Sensitivity Recognizes a protein of 205kDa-220kDa, identified as CD45RA. CD45RA is isoforms of the human leukocyte common antigen (CD45). Human CD45 contains three exons which encode peptide segments designated A, B and C respectively. The differential splcing of the exons generates at least five isoforms. ABC, AB, BC, B and O. This antibody reacts with ABC and BC isoforms. CD45RA is expressed on 40-50% of peripheral CD4+ T-cells, 50% peripheral CD8+ T-cells, and leukemic B-cell lines. T-cells expressing CD45RO are memory cells. CD45RA are naive or virgin T-cells. T-cells expressing CD45RO are memory cells. CD45RA and CD45RO define complementary, predominantly non-overlapping populations of resting peripheral T-cells. It can especial be used to differentiate T-cell ymphomas (CD4+ or CD+ T-cells. It can especial be used to differentiate T-cell ymphomas (CD45RO +ve) from B cell lymphor (CD45RA +ve). Immunogen Stimulated human leukocytes Product Application Details Mestern Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry. Immunohistochemistry-Paraffin, CyTOF-ready Recommended Dilutions Western Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry Application Notes Optimal results using our Tandem dyes, please avoid prolonged exposure to lor extreme temperature fluctuations. These can lead to irreversible degradatio or decoupling, When staining intracellual targets, specific attention to the fix and permeabilization steps in your flow protocol may be required. P	Product Description		
Gene Symbol PTPRC Species Human Marker Leucocyte Marker Specificity/Sensitivity Recognizes a protein of 205kDa-220kDa, identified as CD45RA. CD45RA is isoforms of the human leukocyte common antigen (CD45). Human CD45 contains three exons which encode peptide segments designated A, B and C respectively. The differential splicing of the exons generates at least five isoforms. ABC, AB, BC, B and O. This antibody reacts with ABC and BC isoforms. CD45RA is expressed on 40-50% of peripheral CD8+ T-cells, 50% peripheral CD8+ T-cells, T-cells, and leukemic B-cell lines. T-cells expressing CD45RO are memory cells. CD45RA are naive or virgin T-cells. T-cells. This monoclonal antibot useful in study on the subpopulation of CD4+ or CD8+ T-cells. It can especial be used to differentiate T-cell lymphomas (CD45RO +ve) from B cell lymphor (CD45RA +ve). Immunogen Stimulated human leukocytes Product Application Details Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry. Immunocytochemistry, Immunofluorescence, Immunohistochemistry-Paraffin, CyTOF-ready Recommended Dilutions Western Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry Application Notes Optimal dilution of this antibody should be experimentally determined. For optimal esults using our Tandem dyes, please avoid prolonged exposure to or extreme temperature fluctuations. These can lead to irreversible degradati or decoupling, When staining intracellular targets, specific attention to the fix and permeabilization steps in your flow protocol may be required. Pleasecon our technical support team at technical@novusbi	Host	Mouse	
Species Human Marker Leucocyte Marker Specificity/Sensitivity Recognizes a protein of 205kDa-220kDa, identified as CD45RA. CD45RA is isoforms of the human leukocyte common antigen (CD45). Human CD45 contains three exons which encode peptide segments designated A, B and C respectively. The differential splicing of the exons generates at least five isoforms. ABC, AB, BC, B and O. This antibody reacts with ABC and BC isoforms. CD45RA is expressed on 40-50% of peripheral CD4+ T-cells, 50% peripheral CD4+ T-cells, B-cells, and leukemic B-cell lines. T-cells expressing CD45RA are memory cells. CD45RA and CD45RO define complementary, predominantly non-overlapping populations of resting peripheral T-cells. This monoclonal antiboc useful in study on the subpopulation of CD4+ or CD8+ T-cells. It can especial be used to differentiate T-cell lymphomas (CD45RO +ve) from B cell lymphor (CD45RA +ve). Immunogen Stimulated human leukocytes Product Application Details Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry. Immunohistochemistry-Paraffin, CyTOF-ready Application Notes Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to lor extreme temperature fluctuations. These can lead to irreversible degradation decoupling. When staining intracellular targets, specific attention to the fixe and permeabilization steps in your flow protocol may be required. Please con our technical support team at technical@novusbio.com if you have any	Gene ID	5788	
Marker Leucocyte Marker Specificity/Sensitivity Recognizes a protein of 205kDa-220kDa, identified as CD45RA. CD45RA is isoforms of the human leukocyte common antigen (CD45). Human CD45 contains three exons which encode peptide segments designated A, B and C respectively. The differential splicing of the exons generates at least five isoforms, ABC, AB, BC, B and O. This antibody reacts with ABC and BC isoforms. CD45RA is expressed on 40-50% of peripheral CD4+ T-cells, 50% peripheral CD4+ T-cells, B-cells, and leukemic B-cell lines. T-cells expressing CD45RA are naive or virgin T-cells. T-cells expressing CD45RO are memory cells. CD45RA and CD45RO define complementary, predominantly non-overlapping populations of resting peripheral T-cells. It can especial be used to differentiate T-cell lymphomas (CD45RO +ve) from B cell lymphor (CD45RA +ve). Immunogen Stimulated human leukocytes Product Application Details Applications Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunofluorescence, Immunohistochemistry, CyTOF-ready Application Notes Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to lor extreme temperature fluctuations. These can lead to irreversible degradation of query on protocol may be required. Please com our technical support team at technical@novusbio.com if you have any	Gene Symbol	PTPRC	
Specificity/Sensitivity Recognizes a protein of 205kDa-220kDa, identified as CD45RA. CD45RA is isoforms of the human leukocyte common antigen (CD45). Human CD45 contains three exons which encode peptide segments designated A, B and C respectively. The differential splicing of the exons generates at least five isoforms, ABC, AB, BC, B and O. This antibody reacts with ABC and BC isoforms. CD45RA is expressed on 40-50% of peripheral CD4+ T-cells, 50% peripheral CD8+ T-cells, B-cells, and leukemic B-cell lines. T-cells expressing CD45RA are naive or virgin T-cells. T-cells expressing CD45RA are naive or virgin T-cells. This monoclonal antibody cells. CD45RA are naive or virgin T-cells. This monoclonal antibody useful in study on the subpopulation of CD4+ or CD8+ T-cells. It can especial be used to differentiate T-cell lymphomas (CD45RO +ve) from B cell lymphor (CD45RA +ve). Immunogen Stimulated human leukocytes Product Application Details Applications Mestern Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunobistochemistry-Paraffin, CyTOF-ready Application Notes Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to I or extreme temperature fluctuations. These can lead to irreversible degradati or decoupling. When staining intracellular targets, specific attention to the fix and permeabilization steps in your flow protocol may be required. Please com our technical support team at technical@novusbio.com if you have any	Species	Human	
isoforms of the human leukocyte common antigen (CD45). Human CD45 contains three exons which encode peptide segments designated A, B and C respectively. The differential splicing of the exons generates at least five isoforms, ABC, AB, BC, B and O. This antibody reacts with ABC and BC isoforms. CD45RA is expressed on 40-50% of peripheral CD4+ T-cells, 50% peripheral CD8+ T-cells, B-cells, and leukemic B-cell lines. T-cells expressing CD45RA are naive or virgin T-cells. The cells expressing CD45RO are memory cells. CD45RA and CD45RO define complementary, predominantly non- overlapping populations of resting peripheral T-cells. This monoclonal antiboo useful in study on the subpopulation of CD4+ or CD8+ T-cells. It can especial be used to differentiate T-cell lymphomas (CD45RO +ve) from B cell lymphor (CD45RA +ve).ImmunogenStimulated human leukocytesProduct Application DetailsApplicationsWestern Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry. Immunohistochemistry. Paraffin, CyTOF-readyApplication NotesOptimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to I or extreme temperature fluctuations. These can lead to irreversible degradati or decoupling. When staining intracellular targets, specific attention to the fix and permeabilization steps in your flow protocol may be required. Please con our technical support team at technical@novusbio.com if you have any	Marker	Leucocyte Marker	
Product Application DetailsApplicationsWestern Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-readyRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry Immunofluorescence, Immunohistochemistry-Paraffin, CyTOF-readyApplication NotesOptimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to I or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixe and permeabilization steps in your flow protocol may be required. Please com our technical support team at technical@novusbio.com if you have any	Specificity/Sensitivity	isoforms of the human leukocyte common antigen (CD45). Human CD45 contains three exons which encode peptide segments designated A, B and C, respectively. The differential splicing of the exons generates at least five isoforms, ABC, AB, BC, B and O. This antibody reacts with ABC and BC isoforms. CD45RA is expressed on 40-50% of peripheral CD4+ T-cells, 50% of peripheral CD8+ T-cells, B-cells, and leukemic B-cell lines. T-cells expressing CD45RA are naive or virgin T-cells. T-cells expressing CD45RO are memory T- cells. CD45RA and CD45RO define complementary, predominantly non- overlapping populations of resting peripheral T-cells. This monoclonal antibody is useful in study on the subpopulation of CD4+ or CD8+ T-cells. It can especially be used to differentiate T-cell lymphomas (CD45RO +ve) from B cell lymphomas	
ApplicationsWestern Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-readyRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry Immunofluorescence, Immunohistochemistry-Paraffin, CyTOF-readyApplication NotesOptimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to I or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixe and permeabilization steps in your flow protocol may be required. Please com our technical support team at technical@novusbio.com if you have any	Immunogen	Stimulated human leukocytes	
Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-readyRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry Immunofluorescence, Immunohistochemistry-Paraffin, CyTOF-readyApplication NotesOptimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to I or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixe and permeabilization steps in your flow protocol may be required. Please com our technical support team at technical@novusbio.com if you have any	Product Application Details		
Application NotesImmunofluorescence, Immunohistochemistry-Paraffin, CyTOF-readyApplication NotesOptimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to I or extreme temperature fluctuations. These can lead to irreversible degradati or decoupling. When staining intracellular targets, specific attention to the fixa and permeabilization steps in your flow protocol may be required. Please con our technical support team at technical@novusbio.com if you have any	Applications		
optimal results using our Tandem dyes, please avoid prolonged exposure to l or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixed and permeabilization steps in your flow protocol may be required. Please con our technical support team at technical@novusbio.com if you have any	Recommended Dilutions	Western Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, CyTOF-ready	
	Application Notes	optimal results using our Tandem dyes, please avoid prolonged exposure to light or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixation and permeabilization steps in your flow protocol may be required. Please contact our technical support team at technical@novusbio.com if you have any	





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-54336PEATT594

210-TA-005	TNF-alpha [Unconjugated]
M6000B-1	IL-6 [HRP]
6507-IL-010/CF	IL-4 [Unconjugated]
285-IF-100	IFN-gamma [Unconjugated]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-54336PEATT594

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

