Product Datasheet

CD3 epsilon Antibody (C3e/1308) [DyLight 755] NBP2-54392IR

Unit Size: 100 ul

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-54392IR

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-54392IR



NBP2-54392IR

CD3 epsilon Antibody (C3e/1308) [DyLight 755]

Product Information Unit Size 100 ul Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark. Clonality Monoclonal Clonality Monoclonal Clone C3e/1308 Preservative 0.05% Sodium Azide Isotype IgG2b Kappa Conjugate DyLight 755 Purity Protein A or G purified Buffer SomM Sodium Borate Product Description Journe of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 916 Gene Symbol CD3E Species Human, Mouse Reactivity Notes Mouse reactivity reported in scientific literature (PMID: 3422661). Marker T-Cell Marker Specificity/Sensitivity Recognizes the epsilon-chain of CD3, which consists of five different polypetide chain (designated as gamma, delta, epsilon, zeta, and eta) with MV ranging from 16-28KDa. The CD3 complex is closely associated at the tymphocyte cell surface with the T cell antigen recopridely, CD3 complex is closely associated at the tymphocyte cell surdace. CD3 antige	, , , , , , , , , , , , , , , , , , , ,	
Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark. Clonality Monoclonal Clone C3e/1308 Preservative 0.05% Sodium Azide Isotype IgG2b Kappa Conjugate DyLight 755 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description This conjugate is made on demand. Actual recovery may vary from the stated or the datasheet. Host Mouse Gene ID 916 Gene Symbol CD3E Specificity/Notes Mouse Reactivity Notes Mouse reactivity reported in scientific literature (PMID: 33428661). Marker T-Cell Marker Specificity/Sensitivity Recognizes the epsilon-chain of CD3, which consists of five different polypeptide chains (designated as gamma, delta, epsilon, zetu, and eta) with MW ranging from 16-28K0a. The CD3 complex is closely associated at the lymphocyte cell surface with the T cell antigen receptor (TCR), Reportedly, CD3 complex is involved in signal transduction to the T cell integer. In cortical trymcytes, at papers on the T cell integer. In cortical trymcytes, coD3 is predominantly intra-cytoplasmic. How Probaly is closely associated at the lymphocyte cell sur	Product Information	
services. Storage Store at 4C in the dark. Clonality Monoclonal Clone C3a/1308 Preservative 0.05% Sodium Azide Isotype IgG2b Kappa Conjugate DyLight 755 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 916 Gene Symbol CD3E Species Human, Mouse Reactivity Notes Mouse reactivity reported in scientific literature (PMID: 33428661). Marker T-Cell Marker Specificity/Sensitivity Recognizes the epsilon-chain of CD3, which consists of five different polypeptide chains (designated as gamma, delta, epsilon, zeta, and eta) with MW ranging from 16-2840a. The CD3 complex is lors ordening in involved in signal transduction to the T cell interior following antigen recognition. The CO3 antigen is first detectable in early thymocytes and probably represents one of the earliest signs of commitment to the T cell interior following antigen recognition. The CO3 antigen is first detectable in early thymocytes and probably represents one of the earliest signs of commitrement to the T cell intere in thereport (C	Unit Size	100 ul
Clonality Monoclonal Clone C3e/1308 Preservative 0.05% Sodium Azide Isotype IgG2b Kappa Conjugate DyLight 755 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene D 916 Gene D 916 Gene Symbol CD3E Species Human, Mouse Reactivity Notes Mouse reactivity reported in scientific literature (PMID: 33428661). Marker T-Cell Marker Specificity/Sensitivity Reactivity reported in scientific literature (PMID: 33428661). Marker T-Cell Marker Specificity/Sensitivity Reactivity reported in scientific literature (PMID: 33428661). Marker T-Cell Marker Specificity/Sensitivity Reactivity note the epsilon-chain of CD3, which consists of five different polypeptide chains (designated as gamma, delta, epsilon, zeta, and eta) with MW ranging from 16-28k0a. The CD3 acomplex is lossely associ	Concentration	•
Cione C3e/1308 Preservative 0.05% Sodium Azide Isotype IgG2b Kappa Conjugate DyLight 755 Purity Protein A or G purified Buffer 50m Sodium Borate Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 916 Gene Symbol CD3E Species Human, Mouse Reactivity Notes Mouse reactivity reported in scientific literature (PMID: 33428661). Marker T-Cell Marker Specificity/Sensitivity Recognizes the epsilon-chain of CD3, which consists of five different polypeptide chains (designated as gamma, della, epsilon, zeta, and eta) with MW ranging for 16-28KDa. Specificity/Sensitivity Recognizes the epsilon-the to to the T cell interior following antigen recognition. The CD3 antigen is first detectable in early thymocytes and probably represents on the 32 sets on mitternet to the T cell interior following antigen recognition. The CD3 antigen is sins of commitment to the T cell interior following antigen recognition. The CD3 antigen is a highly specific marker for T cells, and is present in majority of T cell nearker. Specificity/Sensitivity Recombinant human CD3 epsilon fragment (aa 23-119) (Uniprot: P077	Storage	Store at 4C in the dark.
Preservative 0.05% Sodium Azide Isotype IgG2b Kappa Conjugate DyLight 755 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 916 Gene Symbol CD3E Species Human, Mouse Reactivity Notes Mouse reactivity reported in scientific literature (PMID: 33428661). Marker T-Cell Marker Specificity/Sensitivity Recognizes the epsilon-chain of CD3, which consists of five different polypeptide chains (designated as gamma, delta, epsion, zeta, and eta) with MW ranging from 16-28kDa. The CD3 complex is closely associated at the lymphocyte cell survives and probabily represents one of the earliest signs of commitment to the T cell interior following antigen recognition. The CD3 antigen is first detectable in early thymocytes, and highly specific marker for T cells, and is present in majority of T cell meeplasms. Immunogen Recombinant human CD3 epsilon fragment (aa 23-119) (Uniprot: P07766) Notes DyLight (R) is a trademark of Thermo Fisher Scientific line. and its subsidiaries. Product Application Detaiis Western Blot, Flow Cytometry, Flow (In	Clonality	Monoclonal
isotype IgG2b Kappa Conjugate DyLight 755 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 916 Gene Symbol CD3E Species Human, Mouse Reactivity Notes Mouse reactivity reported in scientific literature (PMID: 33428661). Marker T-Cell Marker Specificity/Sensitivity Recognizes the epsilon-chain of CD3, which consists of five different polypeptide chains (designated as gamma, delta, epsilon, zeta, and eta) with MW ranging from 16-28kDa. The CD3 complex is closely associated at the lymphocyte cell surface with the T cell anterior following antigen recognition. The CD3 antigen is first detectable in early thymocytes, and eta) with MW ranging from 16-28kDa. The CD3 complex is closely associated at the lymphocyte cell surface with the T cell anterior following antigen recognition. The CD3 antigen is first detectable in early thymocytes, and ot the modulary thymocytes. CD3 is predominantly intra-cytoplasmic. However, in medullary thymocytes, CD3 is predominantly intra-cytoplasmic. However, in medullary thymocytes, CD3 is predominantly intra-cytoplasmic. However, in medullary thymocytes, CD3 is predominantly intra-cytoplasmic. Howevere, in medullary thymocytes, CD3 is predominantly int	Clone	C3e/1308
Conjugate DyLight 755 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 916 Gene Symbol CD3E Species Human, Mouse Reactivity Notes Mouse reactivity reported in scientific literature (PMID: 33428661). Marker T-Cell Marker Specificity/Sensitivity Recognizes the epsilon-chain of CD3, which consists of five different polypeptide chains (designated as gamma, delta, epsilon, zeta, and eta) with MW ranging from 16-28K0a. The CD3 complex is closely associated at the lymphocyte cell surface with the T cell antigen receptor (TCR). Reportedly, CD3 complex is involved in signal transduction to the T cell lineage. In cortical thymocytes, CD3 is predominantly intra-cytoplasmic. However, in medullary thymocytes, it appears on the T cell surface. CD3 antigen is a highly specific marker for T cells, and is present in majority of T cell neoplasms. Immunogen Recombinant human CD3 epsilon fragment (aa 23-119) (Uniprot: P07766) Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries. Product Application Details Western Blot, Flow Cytometry, Flow (Intracellular),	Preservative	0.05% Sodium Azide
Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 916 Gene Symbol CD3E Species Human, Mouse Reactivity Notes Mouse reactivity reported in scientific literature (PMID: 33428661). Marker T-Cell Marker Specificity/Sensitivity Recognizes the epsilon-chain of CD3, which consists of five different polypeptide chains (designated as gamma, delta, epsilon, zeta, and eta) with MW ranging from 16-28k Da. The CD3 acomplex is closely associated at the lymphocyte cell surface with the T cell antigen receptor (TCR). Reportedly, CD3 complex is involved in signal transduction to the T cell interior following antigen recognition. The CD3 antigen is first detectable in early thymocytes, CD3 complex is one of the earliest signs of commitment to the T cell integae. In cortical thymocytes, CD3 is predominantly intra-cytoplasmic. However, in medullary thymocytes, it appears on the T cell surface. CD3 antigen is a highly specific marker for T cells, and is present in majority of T cell neoplasms. Immunogen Recombinant human CD3 epsilon fragment (aa 23-119) (Uniprot: P07766) Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries. Prod	Isotype	IgG2b Kappa
Buffer 50mM Sodium Borate Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 916 Gene Symbol CD3E Species Human, Mouse Reactivity Notes Mouse reactivity reported in scientific literature (PMID: 33428661). Marker T-Cell Marker Specificity/Sensitivity Recognizes the epsilon-chain of CD3, which consists of five different polypeptide chains (designated as gamma, delta, epsilon, zeta, and eta) with MW ranging from 16-28kDa. The CD3 complex is closely associated at the lymphocyte cell surface with the T cell antigen recognition. The CD3 antigen is first detectable in early thymocytes and probably represents one of the earliest signs of commitment to the T cell liner following antigen recognition. The CD3 antigen is first detectable in early thymocytes, in medullary thymocytes, it appears on the T cell surface. CD3 antigen is a highly specific marker for T cells, and is present in majority of T cell neoplasms. Immunogen Recombinant human CD3 epsilon fragment (aa 23-119) (Uniprot: P07766) Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries. Product Application Details Western Blot, Flow Cytometry, Flow (Intracellular), Immunohistochemistry, Immunohistochemistry, Paraffin, Flow (Intracellular), CyTOF-ready,	Conjugate	DyLight 755
Product Description Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 916 Gene Symbol CD3E Species Human, Mouse Reactivity Notes Mouse reactivity reported in scientific literature (PMID: 33428661). Marker T-Cell Marker Specificity/Sensitivity Recognizes the epsilon-chain of CD3, which consists of five different polypeptide chains (designated as gamma, delta, epsilon, zeta, and eta) with MW ranging from 16-28kDa. The CD3 complex is closely associated at the lymphocyte cell surface with the T cell antigen receptor (TCR). Reportedly, CD3 complex is involved in signal transduction to the T cell linterior following antigen recognition. The CD3 antigen is first detectable in early thymocytes and probably represents one of the earliest signs of commitment to the T cell intege. In cortical thymocytes, it appears on the T cell surface. CD3 antigen is a highly specific marker for T cells, and is present in majority of T cell neoplasms. Immunogen Recombinant human CD3 epsilon fragment (aa 23-119) (Uniprot: P07766) Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries. Product Application Details Western Blot, Flow Cytometry, Flow (Intracellular), Immunohistochemistry, Immunohistochemistry, Paraffin, CyTOF-ready, Multiplex Immunoassay </th <th>Purity</th> <th>Protein A or G purified</th>	Purity	Protein A or G purified
DescriptionThis conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.HostMouseGene ID916Gene SymbolCD3ESpeciesHuman, MouseReactivity NotesMouse reactivity reported in scientific literature (PMID: 33428661).MarkerT-Cell MarkerSpecificity/SensitivityRecognizes the epsilon-chain of CD3, which consists of five different polypeptide chains (designated as gamma, delta, epsilon, zeta, and eta) with MW ranging from 16-28KDa. The CD3 complex is closely associated at the lymphocyte cell surface with the T cell antigen receptor (TCR). Reportedly, CD3 complex is involved in signal transduction to the T cell lineage. In cortical thymocytes, CD3 is predominantly intra-cytoplasmic. However, in medullary thymocytes, it appears on the T cell surface. CD3 antigen is a highly specific marker for T cells, and is present in majority of T cell neoplasms.ImmunogenRecombinant human CD3 epsilon fragment (aa 23-119) (Uniprot: P07766)NotesDylight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.Product Application DetailsWestern Blot, Flow Cytometry, Flow (Intracellular), Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Multiplex ImmunoassayRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry. Paraffin, Flow (Intracellular), CyTOF-ready, Multiplex ImmunoassayApplication NotesUse in Multiplex Immunoassay reported in scientific literature (PMID: 31942075).	Buffer	50mM Sodium Borate
volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.HostMouseGene ID916Gene SymbolCD3ESpeciesHuman, MouseReactivity NotesMouse reactivity reported in scientific literature (PMID: 33428661).MarkerT-Cell MarkerSpecificity/SensitivityRecognizes the epsilon-chain of CD3, which consists of five different polypeptide chains (designated as gamma, delta, epsilon, zeta, and eta) with MW ranging from 16-28kDa. The CD3 complex is closely associated at the lymphocyte cell surface with the T cell antigen receptor (TCR). Reportedly, CD3 complex is involved in signal transduction to the T cell linterior following antigen recognition. The CD3 antigen is first detectable in early thymocytes and probably represents one of the earliest signs of commitment to the T cell lineage. In cortical thymocytes, it appears on the T cell surface. CD3 antigen is a highly specific marker for T cells, and is present in majority of T cell neoplasms.ImmunogenRecombinant human CD3 epsilon fragment (aa 23-119) (Uniprot: P07766)NotesDyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.Product Application DetailsWestern Blot, Flow Cytometry, Flow (Intracellular), Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Multiplex ImmunoassayRecommended DilutionsWestern Blot, Flow Zytometry, Immunohistochemistry- Paraffin, Flow (Intracellular), CyTOF-ready, Multiplex ImmunoassayApplication NotesUse in Multiplex Immunoassay reported in scientific literature (PMID: 31942075).	Product Description	
Gene ID 916 Gene Symbol CD3E Species Human, Mouse Reactivity Notes Mouse reactivity reported in scientific literature (PMID: 33428661). Marker T-Cell Marker Specificity/Sensitivity Recognizes the epsilon-chain of CD3, which consists of five different polypeptide chains (designated as gamma, delta, epsilon, zeta, and eta) with MW ranging from 16-28kDa. The CD3 complex is closely associated at the lymphocyte cell surface with the T cell antigen receptor (TCR). Reportedly, CD3 complex is involved in signal transduction to the T cell interior following antigen recognition. The CD3 antigen is first detectable in early thymocytes and probably represents one of the earliest signs of commitment to the T cell inceage. In cortical thymocytes, it appears on the T cell surface. CD3 antigen is a highly specific marker for T cells, and is present in majority of T cell neoplasms. Immunogen Recombinant human CD3 epsilon fragment (aa 23-119) (Uniprot: P07766) Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries. Product Application Details Western Blot, Flow Cytometry, Flow (Intracellular), Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Multiplex Immunoassay Recommended Dilutions Western Blot, Flow Cytometry, Immunohistochemistry-Paraffin, Flow (Intracellular), Limunohistochemistry-Paraffin, Flow (Intracellular), Limunohistochemistry-Paraffin, Flow (Intracellular), Setter Immunoassay	Description	volume of this product. The volume will be greater than or equal to the unit size
Gene SymbolCD3ESpeciesHuman, MouseReactivity NotesMouse reactivity reported in scientific literature (PMID: 33428661).MarkerT-Cell MarkerSpecificity/SensitivityRecognizes the epsilon-chain of CD3, which consists of five different polypeptide chains (designated as gamma, delta, epsilon, zeta, and eta) with MW ranging from 16-28kDa. The CD3 complex is closely associated at the lymphocyte cell surface with the T cell antigen receptor (TCR). Reportedly, CD3 complex is involved in signal transduction to the T cell interior following antigen recognition. The CD3 antigen is first detectable in early thymocytes and probably represents one of the earliest signs of commitment to the T cell lineage. In cortical thymocytes, CD3 is predominantly intra-cytoplasmic. However, in medullary thymocytes, it appears on the T cell surface. CD3 antigen is a highly specific marker for T cells, and is present in majority of T cell neoplasms.ImmunogenRecombinant human CD3 epsilon fragment (aa 23-119) (Uniprot: P07766)NotesDyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.Product Application DetailsWestern Blot, Flow Cytometry, Flow (Intracellular), Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Multiplex ImmunoassayRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry. Paraffin, Flow (Intracellular), Vultiplex ImmunoassayApplication NotesUse in Multiplex Immunoassay reported in scientific literature (PMID: 31942075).	Host	Mouse
SpeciesHuman, MouseReactivity NotesMouse reactivity reported in scientific literature (PMID: 33428661).MarkerT-Cell MarkerSpecificity/SensitivityRecognizes the epsilon-chain of CD3, which consists of five different polypeptide chains (designated as gamma, delta, epsilon, zeta, and eta) with MW ranging from 16-28kDa. The CD3 complex is closely associated at the lymphocyte cell surface with the T cell antigen receptor (TCR). Reportedly, CD3 complex is involved in signal transduction to the T cell interior following antigen recognition. The CD3 antigen is first detectable in early thymocytes and probably represents one of the earliest signs of commitment to the T cell lineage. In cortical thymocytes, CD3 is predominantly intra-cytoplasmic. However, in medullary thymocytes, CD3 is predominantly intra-cytoplasmic. However, in medullary thymocytes, it appears on the T cell surface. CD3 antigen is a highly specific marker for T cells, and is present in majority of T cell neoplasms.ImmunogenRecombinant human CD3 epsilon fragment (aa 23-119) (Uniprot: P07766)NotesDyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.Product Application DetailsWestern Blot, Flow Cytometry, Flow (Intracellular), Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Multiplex ImmunoassayRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry- Paraffin, Flow (Intracellular), CyTOF-ready, Multiplex ImmunoassayApplication NotesUse in Multiplex Immunoassay reported in scientific literature (PMID: 31942075).	Gene ID	916
Reactivity NotesMouse reactivity reported in scientific literature (PMID: 33428661).MarkerT-Cell MarkerSpecificity/SensitivityRecognizes the epsilon-chain of CD3, which consists of five different polypeptide chains (designated as gamma, delta, epsilon, zeta, and eta) with MW ranging from 16-28kDa. The CD3 complex is closely associated at the lymphocyte cell surface with the T cell antigen receptor (TCR). Reportedly, CD3 complex is involved in signal transduction to the T cell interior following antigen recognition. The CD3 antigen is first detectable in early thymocytes and probably represents one of the earliest signs of commitment to the T cell lineage. In cortical thymocytes, CD3 is predominantly intra-cytoplasmic. However, in medullary thymocytes, it appears on the T cell surface. CD3 antigen is a highly specific marker for T cells, and is present in majority of T cell neoplasms.ImmunogenRecombinant human CD3 epsilon fragment (aa 23-119) (Uniprot: P07766)NotesDyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.Product Application DetailsWestern Blot, Flow Cytometry, Flow (Intracellular), Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Multiplex ImmunoassayRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry- Paraffin, Flow (Intracellular), CyTOF-ready, Multiplex ImmunoassayApplication NotesUse in Multiplex Immunoassay reported in scientific literature (PMID: 31942075).	Gene Symbol	CD3E
Marker T-Cell Marker Specificity/Sensitivity Recognizes the epsilon-chain of CD3, which consists of five different polypeptide chains (designated as gamma, delta, epsilon, zeta, and eta) with MW ranging from 16-28kDa. The CD3 complex is closely associated at the lymphocyte cell surface with the T cell antigen receptor (TCR). Reportedly, CD3 complex is involved in signal transduction to the T cell interior following antigen recognition. The CD3 antigen is first detectable in early thymocytes and probably represents one of the earliest signs of commitment to the T cell intergion. However, in medullary thymocytes, CD3 is predominantly intra-cytoplasmic. However, in medullary thymocytes, it appears on the T cell surface. CD3 antigen is a highly specific marker for T cells, and is present in majority of T cell neoplasms. Immunogen Recombinant human CD3 epsilon fragment (aa 23-119) (Uniprot: P07766) Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries. Product Application Details Western Blot, Flow Cytometry, Flow (Intracellular), Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Multiplex Immunoassay Recommended Dilutions Western Blot, Flow Cytometry, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Multiplex Immunoassay Application Notes Use in Multiplex Immunoassay reported in scientific literature (PMID: 31942075).	Species	Human, Mouse
Specificity/SensitivityRecognizes the epsilon-chain of CD3, which consists of five different polypeptide chains (designated as gamma, delta, epsilon, zeta, and eta) with MW ranging from 16-28kDa. The CD3 complex is closely associated at the lymphocyte cell surface with the T cell antigen receptor (TCR). Reportedly, CD3 complex is involved in signal transduction to the T cell interior following antigen recognition. The CD3 antigen is first detectable in early thymocytes and probably represents one of the earliest signs of commitment to the T cell lineage. In cortical thymocytes, CD3 is predominantly intra-cytoplasmic. However, in medullary thymocytes, it appears on the T cell surface. CD3 antigen is a highly specific marker for T cells, and is present in majority of T cell neoplasms.ImmunogenRecombinant human CD3 epsilon fragment (aa 23-119) (Uniprot: P07766)NotesDyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.Product Application DetailsWestern Blot, Flow Cytometry, Flow (Intracellular), Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Multiplex ImmunoassayRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry, Immunohistochemistry, Paraffin, Flow (Intracellular), CyTOF-ready, Multiplex ImmunoassayApplication NotesUse in Multiplex Immunoassay reported in scientific literature (PMID: 31942075).	Reactivity Notes	Mouse reactivity reported in scientific literature (PMID: 33428661).
chains (designated as gamma, delta, epsilon, zeta, and eta) with MW ranging from 16-28kDa. The CD3 complex is closely associated at the lymphocyte cell surface with the T cell antigen receptor (TCR). Reportedly, CD3 complex is involved in signal transduction to the T cell interior following antigen recognition. The CD3 antigen is first detectable in early thymocytes and probably represents one of the earliest signs of commitment to the T cell lineage. In cortical thymocytes, CD3 is predominantly intra-cytoplasmic. However, in medullary thymocytes, it appears on the T cell surface. CD3 antigen is a highly specific marker for T cells, and is present in majority of T cell neoplasms.ImmunogenRecombinant human CD3 epsilon fragment (aa 23-119) (Uniprot: P07766)NotesDyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.Product Application DetailsWestern Blot, Flow Cytometry, Flow (Intracellular), Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Multiplex ImmunoassayRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry, Paraffin, Flow (Intracellular), CyTOF-ready, Multiplex ImmunoassayApplication NotesUse in Multiplex Immunoassay reported in scientific literature (PMID: 31942075).	Marker	T-Cell Marker
NotesDyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.Product Application DetailsApplicationsWestern Blot, Flow Cytometry, Flow (Intracellular), Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Multiplex ImmunoassayRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry, Immunohistochemistry- Paraffin, Flow (Intracellular), CyTOF-ready, Multiplex ImmunoassayApplication NotesUse in Multiplex Immunoassay reported in scientific literature (PMID: 31942075).	Specificity/Sensitivity	chains (designated as gamma, delta, epsilon, zeta, and eta) with MW ranging from 16-28kDa. The CD3 complex is closely associated at the lymphocyte cell surface with the T cell antigen receptor (TCR). Reportedly, CD3 complex is involved in signal transduction to the T cell interior following antigen recognition. The CD3 antigen is first detectable in early thymocytes and probably represents one of the earliest signs of commitment to the T cell lineage. In cortical thymocytes, CD3 is predominantly intra-cytoplasmic. However, in medullary thymocytes, it appears on the T cell surface. CD3 antigen is a highly specific
DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.Product Application DetailsApplicationsWestern Blot, Flow Cytometry, Flow (Intracellular), Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Multiplex ImmunoassayRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry, Immunohistochemistry- Paraffin, Flow (Intracellular), CyTOF-ready, Multiplex ImmunoassayApplication NotesUse in Multiplex Immunoassay reported in scientific literature (PMID: 31942075).	Immunogen	Recombinant human CD3 epsilon fragment (aa 23-119) (Uniprot: P07766)
ApplicationsWestern Blot, Flow Cytometry, Flow (Intracellular), Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Multiplex ImmunoassayRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry, Immunohistochemistry- Paraffin, Flow (Intracellular), CyTOF-ready, Multiplex ImmunoassayApplication NotesUse in Multiplex Immunoassay reported in scientific literature (PMID: 31942075).	Notes	DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.
Immunohistochemistry-Paraffin, CyTOF-ready, Multiplex ImmunoassayRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry, Immunohistochemistry- Paraffin, Flow (Intracellular), CyTOF-ready, Multiplex ImmunoassayApplication NotesUse in Multiplex Immunoassay reported in scientific literature (PMID: 31942075).	Product Application Details	
Paraffin, Flow (Intracellular), CyTOF-ready, Multiplex ImmunoassayApplication NotesUse in Multiplex Immunoassay reported in scientific literature (PMID: 31942075).	Applications	
	Recommended Dilutions	
	Application Notes	





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-54392IR

NBP1-43317IR	Mouse IgG2b Kappa Light Chain Isotype Control (MG2b) [DyLight 755]
NBP2-22752	Recombinant Human CD3 epsilon His Protein
210-TA-005	TNF-alpha [Unconjugated]
9850-CD-050	CD3 epsilon [Unconjugated]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-54392IR

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

