# **Product Datasheet**

# CD3 epsilon Antibody (C3e/1308) [PE/Cy5.5] NBP2-54392PECY55

Unit Size: 0.1 ml

Store at 4C in the dark. Do not freeze.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-54392PECY55

Updated 10/23/2024 v.20.1







# NBP2-54392PECY55

CD3 epsilon Antibody (C3e/1308) [PE/Cy5.5]

0.1 ml	
Please see the vial label for concentration. If unlisted please contact technical services.	
Store at 4C in the dark. Do not freeze.	
Monoclonal	
C3e/1308	
0.05% Sodium Azide	
IgG2b Kappa	
PE/Cy5.5	
Protein A or G purified	
PBS	
Mouse	
916	
CD3E	
Human, Mouse	
Mouse reactivity reported in scientific literature (PMID: 33428661).	
T-Cell Marker	
Recognizes the epsilon-chain of CD3, which consists of five different polypeptide chains (designated as gamma, delta, epsilon, zeta, and eta) with MW ranging from 16-28kDa. The CD3 complex is closely associated at the lymphocyte cell surface with the T cell antigen receptor (TCR). Reportedly, CD3 complex is involved in signal transduction to the T cell interior following antigen recognition. The CD3 antigen is first detectable in early thymocytes and probably represents one of the earliest signs of commitment to the T cell lineage. In cortical thymocytes, CD3 is predominantly intra-cytoplasmic. However, in medullary thymocytes, it appears on the T cell surface. CD3 antigen is a highly specific marker for T cells, and is present in majority of T cell neoplasms.	
Recombinant human CD3 epsilon fragment (aa 23-119) (Uniprot: P07766)	
Product Application Details	
Western Blot, Flow Cytometry, Flow (Intracellular), Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Multiplex Immunoassay	
Western Blot, Flow Cytometry, Immunohistochemistry, Immunohistochemistry- Paraffin, Flow (Intracellular), CyTOF-ready, Multiplex Immunoassay	
Use in Multiplex Immunoassay reported in scientific literature (PMID: 31942075). Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to light or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixation and permeabilization steps in your flow protocol may be required. Please contact our technical support team at technical@novusbio.com if you have any questions.	





#### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

# **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

# **General Contact Information**

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

# Products Related to NBP2-54392PECY55

NBP1-43317PECY55	Mouse IgG2b Kappa Light Chain Isotype Control (MG2b) [PE/Cy5.5]
NBP2-22752	Recombinant Human CD3 epsilon His Protein
210-TA-005	TNF-alpha [Unconjugated]
9850-CD-050	CD3 epsilon [Unconjugated]

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-54392PECY55

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

