## **Product Datasheet**

### TIA1 Antibody (TIA1/1313) [Alexa Fluor® 405] NBP2-54406AF405

Unit Size: 100 ul

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-54406AF405

Updated 10/23/2024 v.20.1

# Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-54406AF405



#### NBP2-54406AF405

TIA1 Antibody (TIA1/1313) [Alexa Fluor® 405]

Unit Size         100 ul           Concentration         Please see the vial label for concentration. If unlisted please contact technical services.           Storage         Store at 4C in the dark.           Clonality         Monoclonal           Clone         TIA1/1313           Preservative         0.05% Sodium Azide           Isotype         IgG2b Kappa           Conjugate         Alexa Fluor 405           Purity         Protein A purified           Buffer         50mM Sodium Borate           Product Description         Mouse           Gene ID         7072           Gene Symbol         TIA1           Species         Human           Specificity/Sensitivity         TIA-1 (T-cell intracytoplasmic antigen) is a cytoplasmic granule-associated protein as it preferentially recognizes poly(A) homopolymers and induces DNA fragmentation in CTL targets: The major granule-associated species is a 15kDa protein that this protein may be involved in the induction of apoptosis as it preferentially recognizes poly(A) homopolymers and induces DNA fragmentation in CTL targets:. The major granule-associated species is a 15kDa protein thy to be drived in the induction of apoptosis as is preferentially recognizes poly(A) homopolymers and induces DNA fragmentation in CTL targets:. The major granule-associated species is a 15kDa protein thy to be drived in the induction of apoptosis as a to the contal the carbox/ terminal with cells and natural killer cells (NK cells). It is also expressed in T-cell magnazies (low expr		
Concentration         Please see the vial label for concentration. If unlisted please contact technical services.           Storage         Store at 4C in the dark.           Clonality         Monoclonal           Clone         TIA1/1313           Preservative         0.05% Sodium Azide           Isotype         IgG2b Kappa           Conjugate         Alexa Fluor 405           Purity         Protein A purified           Buffer         50mM Sodium Borate           Product Description         Mouse           Gene ID         7072           Gene Symbol         TIA1           Species         Human           Specificity/Sensitivity         TIA-1 (T-cell intracytoplasmic antigen) is a cytoplasmic granule-associated protein, expressed in lymphocytes processing cytolytic potential. TIA-1 is a member of an RNA-binding protein framily and possessen sucleolytic activity against cytotoxic lymphocyte (CTL) target cells. It has been suggested that this protein may be involved in the induction of apoptosis as its preferentially recognizes poly(A) homopolymers and induces DNA fragmentation in CTL targets. The major granule-associated species is a 15kDa protein thought to be derived from the carboxyl terminus of the 40kDa product by proteolytic processing. TIA antibody labels cytoxic T cells and natural killer cells (NK cells). It is also expressed in T-cell lymphocyte (LGL) leukemia and hairy cell leukemia (high expression) in T-cell malignancies.           Inmounogen         Recombinant human TIA1 fragment of 102 </th <th>Product Information</th> <th></th>	Product Information	
services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneTIA1/1313Preservative0.05% Sodium AzideIsotypeIgG2b KappaConjugateAlexa Fluor 405PurityProtein A purifiedBuffer50mM Sodium BorateProduct DescriptionMouseGene ID7072Gene SymbolTIA1SpeciesHumanSpeciesTIA1 (T-cell intracytoplasmic antigen) is a cytoplasmic granule-associated protein, expressed in lymphocyte (CTL) target cells. It has been suggested that this protein may be involved in the induction of apoptosis as it preferentially recognizes. The major granule-associated sprotes in ducts to the 40KDa protein by terkingt of the 1-cell sodial sociated sprotes in the induction of the 40KDa protein by recognizes. The major granule-associated sprotesions in a subovy of the 40KDa protein by recognizes. The major granule-associated sprotesions in the induction of the 40KDa protein by recognizes. The major granule-associated species is a 15kDa protein hought to be derived from the carboxy/ terminus of the 40KDa product by proteolytic processing. TIA1 antiody labels cytotoxic T cell and natural killer cells (NK cells). It is also expressed in T-cell lymphocytes in the study of immune response to nalignancies.ImmunogenRecombinant human TIA1 fragment of 102 amino acid residues (aa279-380)	Unit Size	100 ul
Clonality       Monoclonal         Clone       TIA1/1313         Preservative       0.05% Sodium Azide         Isotype       IgG2b Kappa         Conjugate       Alexa Fluor 405         Purity       Protein A purified         Buffer       50mM Sodium Borate         Product Description       Mouse         Gene ID       7072         Gene Symbol       TIA1         Species       Human         Specificity/Sensitivity       TIA-1 (T-cell intracytoplasmic antigen) is a cytoplasmic granule-associated protein expressed in lymphocytes processing cytolytic potential. TIA-1 is a member of an RNA-binding protein family and possesses nucleolytic activity against cytotoxic lymphocyte (CLL) target cells. It has been suggested that this protein may be involved in the induction of apoptosis as it preferentially recognizes poly(A) homopolymers and induces DNA fragmentation in CTL targets. The major granule-associated species is a 15kDa protein thought to be derived from the carboxyl terminus of the 40kDa product by proteolytic processing. TIA1 antibody labels cytotoxic T cells and natural killer cells (NK cells). It is also expressed in T-cell lymphoma, large granular lymphocyte (ICL).         Intervention       Eukemia and hairy cell Leukemia. TIA1 expression in T-cell lymphocytosis and other T-cell diseases (low expression). ThA1 may also be used to label turor-infiltrating lymphocytes in the study of immune response to malignancies.	Concentration	
CloneTIA1/1313Preservative0.05% Sodium AzideIsotypeIgG2b KappaConjugateAlexa Fluor 405PurityProtein A purifiedBuffer50mM Sodium BorateProduct Description4000HostMouseGene ID7072Gene SymbolTIA1SpeciesHumanSpeciesTIA-1 (T-cell intracytoplasmic antigen) is a cytoplasmic granule-associated protein, expressed in lymphocytes processing cytolytic potential. TIA-1 is a member of an RNA-binding protein family and possesses nucleolytic activity against cytotoxic lymphocyte (CTL) target cells. It has been suggested that this protein may be involved in the induction of apoptosis as it preferentially recognizes poly(A) homopolymers and induces DNA fragmentation in CTL targets. The major granule-associated species is a 15kDa protein thought to be derived from the carboxyl terminus of the 40kDa product by proteolytic processing. TIA1 antibody labels cytotoxic T cells and natural killer cells (NK cells). It is also expressed in T-cell lymphocytes in T-cell malignancies may help in differentiating LGL leukemia. TIA1 expression in T-cell malignancies.ImmunogenRecombinant human TIA1 fragment of 102 amino acid residues (aa279-380)	Storage	Store at 4C in the dark.
Preservative       0.05% Sodium Azide         Isotype       IgG2b Kappa         Conjugate       Alexa Fluor 405         Purity       Protein A purified         Buffer       50mM Sodium Borate         Product Description       Mouse         Gene ID       7072         Gene Symbol       TIA1         Species       Human         Specificity/Sensitivity       TIA-1 (T-cell intracytoplasmic antigen) is a cytoplasmic granule-associated protein, expressed in lymphocytes processing cytolytic potential. TIA-1 is a member of an RNA-binding protein family and possesses nucleolytic activity against cytotoxic lymphocyte (CTL) target cells. It has been suggested that this protein may be involved in the induction of apoptosis as it preferentially recognizes poly(A) homopolymers and induces DNA fragmentation in CTL targets. The major granule-associated species is a 15kDa protein thought to be derived from the carboxyl terminus of the 40kDa product by proteolytic processing. TIA1 antibody labels cytotoxic T cells and natural killer cells (NK cells). It is also expressed in T-cell lymphoma, large granular lymphocyte (LGL) leukemia and hairy cell leukemia. TIA1 expression) from T-cell malignancies may help in differentiating LGL leukemia (high expression) from T-cell malignancies.         Immunogen       Recombinant human TIA1 fragment of 102 amino acid residues (aa279-380)	Clonality	Monoclonal
Isotype       IgG2b Kappa         Conjugate       Alexa Fluor 405         Purity       Protein A purified         Buffer       50mM Sodium Borate         Product Description       Mouse         Gene ID       7072         Gene Symbol       TIA1         Species       Human         Specificity/Sensitivity       TIA-1 (T-cell intracytoplasmic antigen) is a cytoplasmic granule-associated protein, expressed in lymphocytes processing cytolytic potential. TIA-1 is a member of an RNA-binding protein family and possesses nucleolytic activity against cytotoxic lymphocyte (CTL) target cells. It has been suggested that this protein may be involved in the induction of apoptosis as it preferentially recognizes poly(A) homopolymers and induces DNA fragmentation in CTL targets. The major granule-associated species is a 15kDa protein thought to be derived from the carboxyl terminus of the 40kDa product by proteolytic processing. TIA1 antibody labels cytotoxic T cells and natural killer cells (NK cells). It is also expressed in T-cell lymphoma, large granular lymphocytes (LGL) leukemia and hairy cell leukemia. TIA1 expression in T-cell malignancies. may help in differentiating LGL leukemia (high expression) from T-cell lymphocytosis and other T-cell diseases (low expression). TIA1 may also be used to label tumor-infiltrating lymphocytes in the study of immune response to malignancies.	Clone	TIA1/1313
ConjugateAlexa Fluor 405PurityProtein A purifiedBuffer50mM Sodium BorateProduct DescriptionMouseHostMouseGene ID7072Gene SymbolTIA1SpeciesHumanSpecificity/SensitivityTIA-1 (T-cell intracytoplasmic antigen) is a cytoplasmic granule-associated protein, expressed in lymphocytes processing cytolytic potential. TIA-1 is a member of an RNA-binding protein family and posesses nucleolytic activity against cytotoxic lymphocyte (CTL) target cells. It has been suggested that this protein may be involved in the induction of apoptosis as it preferentially recognizes poly(A) homopolymers and induces DNA fragmentation in CTL targets. The major granule-associated species is a 15kDa protein thought to be derived from the carboxyl terminus of the 40kDa product by proteolytic processing. TIA-1 attibody labels cytotoxic T cells and natural killer cells (NK cells). It is also expressed in T-cell lymphoma, large granular lymphocyte (LGL) leukemia and hairy cell leukemia. TIA1 expression in T-cell lymphocytosis and other T-cell diseases (low expression). TIA1 may also be used to label tumor-infiltrating lymphocytes in the study of immune response to malignancies.ImmunogenRecombinant human TIA1 fragment of 102 amino acid residues (aa279-380)	Preservative	0.05% Sodium Azide
Purity       Protein A purified         Buffer       50mM Sodium Borate         Product Description       Mouse         Gene ID       7072         Gene Symbol       TIA1         Species       Human         Specificity/Sensitivity       TIA-1 (T-cell intracytoplasmic antigen) is a cytoplasmic granule-associated protein, expressed in lymphocytes processing cytolytic potential. TIA-1 is a member of an RNA-binding protein family and possesses nucleolytic activity against cytotoxic lymphocyte (CTL) target cells. It has been suggested that this protein may be involved in the induction of apoptosis as it preferentially recognizes poly(A) homopolymers and induces DNA fragmentation in CTL targets. The major granule-associated species is a 15kDa protein thought to be derived from the carboxyl terminus of the 40kDa product by proteolytic processing. TIA 1 antibody labels cytotoxic T cells and natural killer cells (NK cells). It is also expressed in T-cell lymphoma, large granular lymphocytes (LGL) leukemia and hairy cell leukemia. TIA1 expression in T-cell mynphocytes is and other T-cell diseases (low expression). TIA1 may also be used to label tumor-infiltrating lymphocytes in the study of immune response to malignancies.         Immunogen       Recombinant human TIA1 fragment of 102 amino acid residues (aa279-380)	Isotype	IgG2b Kappa
Buffer       50mM Sodium Borate         Product Description       Mouse         Gene ID       7072         Gene Symbol       TIA1         Species       Human         Specificity/Sensitivity       TIA-1 (T-cell intracytoplasmic antigen) is a cytoplasmic granule-associated protein, expressed in lymphocytes processing cytolytic potential. TIA-1 is a member of an RNA-binding protein family and possesses nucleolytic activity against cytotoxic lymphocyte (CTL) target cells. It has been suggested that this protein may be involved in the induction of apoptosis as it preferentially recognizes poly(A) homopolymers and induces DNA fragmentation in CTL targets. The major granule-associated species is a 15kDa protein thought to be derived from the carboxyl terminus of the 40kDa product by protelytic processing, TIA1 antibody labels cytotoxic T cells and natural killer cells (NK cells). It is also expressed in T-cell lymphoma, large granular lymphocyte (LGL) leukemia and hairy cell leukemia. TIA1 expression in T-cell malignancies may help in differentiating LGL leukemia the expression). TIA1 may also be used to label tumor-infiltrating lymphocytes in the study of immune response to malignancies.         Immunogen       Recombinant human TIA1 fragment of 102 amino acid residues (aa279-380)	Conjugate	Alexa Fluor 405
Product Description         Host       Mouse         Gene ID       7072         Gene Symbol       TIA1         Species       Human         Specificity/Sensitivity       TIA-1 (T-cell intracytoplasmic antigen) is a cytoplasmic granule-associated protein, expressed in lymphocytes processing cytolytic potential. TIA-1 is a member of an RNA-binding protein family and possesses nucleolytic activity against cytotoxic lymphocyte (CTL) target cells. It has been suggested that this protein may be involved in the induction of apoptosis as it preferentially recognizes poly(A) homopolymers and induces DNA fragmentation in CTL targets. The major granule-associated species is a 15kDa protein thought to be derived from the carboxyl terminus of the 40kDa product by proteolytic processing. TIA1 antibody labels cytotoxic T cells and natural killer cells (NK cells). It is also expressed in T-cell lymphoma, large granular lymphocyte (LGL) leukemia and hairy cell leukemia. TIA1 expression in T-cell malignancies may help in differentiating LGL leukemia (high expression) from T-cell lymphocytosis and other T-cell diseases (low expression). TIA1 may also be used to label tumor-infiltrating lymphocytes in the study of immune response to malignancies.	Purity	Protein A purified
HostMouseGene ID7072Gene SymbolTIA1SpeciesHumanSpecificity/SensitivityTIA-1 (T-cell intracytoplasmic antigen) is a cytoplasmic granule-associated protein, expressed in lymphocytes processing cytolytic potential. TIA-1 is a member of an RNA-binding protein family and possesses nucleolytic activity against cytotoxic lymphocyte (CTL) target cells. It has been suggested that this protein may be involved in the induction of apoptosis as it preferentially recognizes poly(A) homopolymers and induces DNA fragmentation in CTL targets. The major granule-associated species is a 15kDa protein thought to be derived from the carboxyl terminus of the 40kDa product by proteolytic processing. TIA1 antibody labels cytotoxic T cells and natural killer cells (NK cells). It is also expressed in T-cell lymphoma, large granular lymphocyte (LGL) leukemia and hairy cell leukemia (high expression) from T-cell lymphocytosis and other T-cell diseases (low expression). TIA1 may also be used to label tumor-infiltrating lymphocytes in the study of immune response to malignancies.ImmunogenRecombinant human TIA1 fragment of 102 amino acid residues (aa279-380)	Buffer	50mM Sodium Borate
Gene ID7072Gene SymbolTIA1SpeciesHumanSpecificity/SensitivityTIA-1 (T-cell intracytoplasmic antigen) is a cytoplasmic granule-associated protein, expressed in lymphocytes processing cytolytic potential. TIA-1 is a member of an RNA-binding protein family and possesses nucleolytic activity against cytotoxic lymphocyte (CTL) target cells. It has been suggested that this protein may be involved in the induction of apoptosis as it preferentially recognizes poly(A) homopolymers and induces DNA fragmentation in CTL targets. The major granule-associated species is a 15kDa protein thought to be derived from the carboxyl terminus of the 40kDa product by proteolytic processing. TIA1 antibody labels cytotoxic T cells and natural killer cells (NK cells). It is also expressed in T-cell lymphoma, large granular lymphocyte (LGL) leukemia and hairy cell leukemia (high expression) from T-cell malignancies may help in differentiating LGL leukemia (high expression) from T-cell lymphocytosis and other T-cell diseases (low expression). TIA1 may also be used to label tumor-infiltrating lymphocytes in the study of immune response to malignancies.ImmunogenRecombinant human TIA1 fragment of 102 amino acid residues (aa279-380)	Product Description	
Gene SymbolTIA1SpeciesHumanSpecificity/SensitivityTIA-1 (T-cell intracytoplasmic antigen) is a cytoplasmic granule-associated protein, expressed in lymphocytes processing cytolytic potential. TIA-1 is a member of an RNA-binding protein family and possesses nucleolytic activity against cytotoxic lymphocyte (CTL) target cells. It has been suggested that this protein may be involved in the induction of apoptosis as it preferentially recognizes poly(A) homopolymers and induces DNA fragmentation in CTL targets. The major granule-associated species is a 15kDa protein thought to be derived from the carboxyl terminus of the 40kDa product by proteolytic processing. TIA1 antibody labels cytotoxic T cells and natural killer cells (NK cells). It is also expressed in T-cell lymphoma, large granular lymphocyte (LGL) leukemia and hairy cell leukemia (high expression) from T-cell lymphocytosis and other T-cell diseases (low expression). TIA1 may also be used to label tumor-infiltrating lymphocytes in the study of immune response to malignancies.ImmunogenRecombinant human TIA1 fragment of 102 amino acid residues (aa279-380)	Host	Mouse
SpeciesHumanSpecificity/SensitivityTIA-1 (T-cell intracytoplasmic antigen) is a cytoplasmic granule-associated protein, expressed in lymphocytes processing cytolytic potential. TIA-1 is a member of an RNA-binding protein family and possesses nucleolytic activity against cytotoxic lymphocyte (CTL) target cells. It has been suggested that this protein may be involved in the induction of apoptosis as it preferentially recognizes poly(A) homopolymers and induces DNA fragmentation in CTL targets. The major granule-associated species is a 15kDa protein thought to be derived from the carboxyl terminus of the 40kDa product by proteolytic processing. TIA1 antibody labels cytotoxic T cells and natural killer cells (NK cells). It is also expressed in T-cell lymphoma, large granular lymphocyte (LGL) leukemia and hairy cell leukemia. TIA1 expression in T-cell malignancies may help in differentiating LGL leukemia (high expression) from T-cell lymphocytosis and other T-cell diseases (low expression). TIA1 may also be used to label tumor-infiltrating lymphocytes in the study of immune response to malignancies.ImmunogenRecombinant human TIA1 fragment of 102 amino acid residues (aa279-380)	Gene ID	7072
Specificity/SensitivityTIA-1 (T-cell intracytoplasmic antigen) is a cytoplasmic granule-associated protein, expressed in lymphocytes processing cytolytic potential. TIA-1 is a member of an RNA-binding protein family and possesses nucleolytic activity against cytotoxic lymphocyte (CTL) target cells. It has been suggested that this protein may be involved in the induction of apoptosis as it preferentially recognizes poly(A) homopolymers and induces DNA fragmentation in CTL targets. The major granule-associated species is a 15kDa protein thought to be derived from the carboxyl terminus of the 40kDa product by proteolytic processing. TIA1 antibody labels cytotoxic T cells and natural killer cells (NK cells). It is also expressed in T-cell lymphoma, large granular lymphocyte (LGL) leukemia and hairy cell leukemia. TIA1 expression in T-cell malignancies may help in differentiating LGL leukemia (high expression) from T-cell lymphocytosis and other T-cell diseases (low expression). TIA1 may also be used to label tumor-infiltrating lymphocytes in the study of immune response to malignancies.ImmunogenRecombinant human TIA1 fragment of 102 amino acid residues (aa279-380)	Gene Symbol	TIA1
<ul> <li>protein, expressed in lymphocytes processing cytolytic potential. TIA-1 is a member of an RNA-binding protein family and possesses nucleolytic activity against cytotoxic lymphocyte (CTL) target cells. It has been suggested that this protein may be involved in the induction of apoptosis as it preferentially recognizes poly(A) homopolymers and induces DNA fragmentation in CTL targets. The major granule-associated species is a 15kDa protein thought to be derived from the carboxyl terminus of the 40kDa product by proteolytic processing. TIA1 antibody labels cytotoxic T cells and natural killer cells (NK cells). It is also expressed in T-cell lymphoma, large granular lymphocyte (LGL) leukemia and hairy cell leukemia. TIA1 expression in T-cell malignancies may help in differentiating LGL leukemia (high expression) from T-cell lymphocytosis and other T-cell diseases (low expression). TIA1 may also be used to label tumor-infiltrating lymphocytes in the study of immune response to malignancies.</li> <li>Immunogen</li> </ul>	Species	Human
	Specificity/Sensitivity	protein, expressed in lymphocytes processing cytolytic potential. TIA-1 is a member of an RNA-binding protein family and possesses nucleolytic activity against cytotoxic lymphocyte (CTL) target cells. It has been suggested that this protein may be involved in the induction of apoptosis as it preferentially recognizes poly(A) homopolymers and induces DNA fragmentation in CTL targets. The major granule-associated species is a 15kDa protein thought to be derived from the carboxyl terminus of the 40kDa product by proteolytic processing. TIA1 antibody labels cytotoxic T cells and natural killer cells (NK cells). It is also expressed in T-cell lymphoma, large granular lymphocyte (LGL) leukemia and hairy cell leukemia. TIA1 expression in T-cell malignancies may help in differentiating LGL leukemia (high expression) from T-cell lymphocytosis and other T-cell diseases (low expression). TIA1 may also be used to label
	Immunogen	



	C I
Notes	Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.
Product Application Details	
Applications	Western Blot, Flow Cytometry, Flow (Intracellular), Immunocytochemistry/ Immunofluorescence, CyTOF-ready
Recommended Dilutions	Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Flow (Intracellular), CyTOF-ready
Application Notes	Optimal dilution of this antibody should be experimentally determined.
4	





#### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

#### Products Related to NBP2-54406AF405

NBP1-43317AF405	Mouse IgG2b Kappa Light Chain Isotype Control (MG2b) [Alexa Fluor® 405]
H00007072-P01-10ug	Recombinant Human TIA1 GST (N-Term) Protein
210-TA-005	TNF-alpha [Unconjugated]
NBP2-06452	TIA1 Overexpression Lysate

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-54406AF405

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

