Product Datasheet

TIA1 Antibody (TIA1/1313) [Alexa Fluor® 647] NBP2-54406AF647

Unit Size: 100 ul

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-54406AF647

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-54406AF647



NBP2-54406AF647

TIA1 Antibody (TIA1/1313) [Alexa Fluor® 647]

Product Information Unit Size 100 ul Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark. Clonality Monoclonal Clone TIA1/1313 Preservative 0.05% Sodium Azide Isotype IgG2b Kappa Conjugate Alexa Fluor 647 Purity Protein A purified Buffer 50mM Sodium Borate Product Description House Gene ID 7072 Gene Symbol TIA1 Species Human Specificity/Sensitivity TIA-1 (r-cell intracytoplasmic antigen) is a cytoplasmic granule-associated protein, expressed in lymphocytes processing cytolytic potential. TIA-1 is a member of an RNA-binding protein family and possesses nucleolytic activity against cytotoxic hymphocyte (CTL) target cells. It has been suggested that this procein may be involved in the induction of apoptosis as it preferentially recognizes poly(A) homopolymers and induces DNA fragmentation in CTL targets. The major granule-associated protein (SLB) protein hought to be derived from the carboxyl terminus of the 40kDa product by proteolytic processing. TIA-1 antibody labels cytotoxic T cells and natural killer cells (NK cells). It is allow expression). TIA-1 may also be used to label tumor-infiltrating lymphocytes in T-cell maligna			
ConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalClonalityMonoclonalClonality0.05% Sodium AzideIsotypeIgG2b KappaConjugateAlexa Fluor 647PurityProtein A purifiedBuffer50mM Sodium BorateProduct DescriptionMouseGene ID7072Gene SymbolTIA1Specificity/SensitivityTIA-1 (T-cell intracytoplasmic antigen) is a cytoplasmic granule-associated protein, expressed in hymphocytes processing cytolytic potential. TIA-1 is a member of an RNA-binding protein family and possessen sucleolytic activity against cytotoxic lymphocyte (CTL) target cells. It has been suggested that this protein may be involved in the induction of aportosis as its preferentially recognizes poly(A) homopolymers and induces DNA fragmentation in CTL targets. The major granule-associated species is a 15kDa protein thought to be derived from the carboxyl terminus of the 40kDa product by protolytic processing. TIA 1 asloe expressed in T-cell lymphocyte (ICL) leukemia and hairy cell leukemia (high expression) nr T-cell lymphocyte (LGL) leukemia and hairy cell leukemia (high expression) nr T-cell lymphocyte (LGL) leukemia and hairy cell leukemia (high expression) nr T-cell lymphocyte (LGL) leukemia and hairy cell leukemia (high expression) nr T-cell seganser (we expression) nr T-cell lymphocyte (LGL) leukemia and hairy cell leukemia (high expression) nr T-cell lymphocytes in help in differentiating LGL leukemia (high expression) from T-cell lymphocytes (az279-380)	Product Information		
services. Storage Store at 4C in the dark. Clonality Monoclonal Clone TIA1/1313 Preservative 0.05% Sodium Azide Isotype IgG2b Kappa Conjugate Alexa Fluor 647 Purity Protein A purified Buffer 50mM Sodium Borate Product Description Mouse Gene ID 7072 Gene ID 7072 Gene Symbol TIA1 Species Human Specificity/Sensitivity TIA-1 (T-cell intracytoplasmic antigen) is a cytoplasmic granule-associated protein, expressed in lymphocytes processing cytolytic potential. TIA-1 is a member of an RNA-binding protein family and possesses nucleolytic activity against cytotoxic lymphocyte (CTL) target cells. It has been suggested that this protein may be involved in the induction of apoptosis as it preferentially recognizes poly(A) homopolymers and induces DNA fragmentation in CTL targets. The major granule-associated species is a 15kDa protein thought to be derived from the carboxyl terminus of the 40kDa product by proteolytic processing. TIA1 antibody labels cytotoxic T cell and natural killer cells (NK cells). It is also expressed in T-cell lymphocyte for CL leukemia and hairy cell leukemia. TIA1 expression in T-cell manginancies may help in differentiating LGL leukemia. INA1 expression from T-cell manginancies and help in certain figh expression) from T-cell manginancies and help of antimure response to malignancies.	Unit Size	100 ul	
ClonalityMonoclonalCloneTIA1/1313Preservative0.05% Sodium AzideIsotypeIgG2b KappaConjugateAlexa Fluor 647PurityProtein A purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID7072Gene SymbolTIA1SpeciesHumanSpeciesHumanSpeciesHumanSpeciesSite of a RNA-binding protein family and possesses nucleolytic activity against cytotoxic lymphocyte (CTL) targenetial. TIA-1 is a member of an RNA-binding protein family and possesses nucleolytic activity against cytotoxic lymphocyte (CTL) targenetiation in CTL targets. The major granule-associated protein may be involved in the induction of apoptosis as it preferentially recognizes poly(A) homopolymers and induces DNA fragmentation in CTL targets. The major granule-associated sprotes is a 15kDa protein thought to be derived from the carboxyl terminus of the 40kDa protein thought to be derived from the carboxyl terminus of the 40kDa protein langipancies may help in differentialing LGL leukemia (high expression) T-cell lymphocytes (CLL) largets on T-cell lymphocytes (CLL)ImmunogenRecombinant human TIA1 fargment of 102 amino acid residues (aa279-380)	Concentration		
CloneTIA1/1313Preservative0.05% Sodium AzideIsotypeIgG2b KappaConjugateAlexa Fluor 647PurityProtein A purifiedBuffer50mM Sodium BorateProduct Description4000000000000000000000000000000000000	Storage	Store at 4C in the dark.	
Preservative0.05% Sodium AzideIsotypeIgG2b KappaConjugateAlexa Fluor 647PurityProtein A purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID7072Gene SymbolTIA1SpeciesHumanSpecificity/SensitivityTIA-1 (T-cell intracytoplasmic antigen) is a cytoplasmic granule-associated protein may be involved in the induction of apoptosis as it preferentially recognizes poly(A) homopolymers and induces DNA fragmentation in CTL targets. The major granule-associated species is a 15kDa protein thought to be derived from the carboxyl terminus of the 40kDa product by proteolytic processing. TIA1 antibody labels cytotoxic T cell sand natural killer cells (NK cells). It is also expressed in T-cell lymphoma, large granular lymphocytosis and other T-cell diseases (low expression). TIA1 may also be used to label tumor-infiltrating lymphocytes in the study of immure response to malignancies.ImmunogenRecombinant human TIA1 fragment of 102 amino acid residues (aa279-380)	Clonality	Monoclonal	
IsotypeIgG2b KappaConjugateAlexa Fluor 647PurityProtein A purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID7072Gene SymbolTIA1SpeciesHumanSpeciesHumanSpecificity/SensitivityTIA-1 (T-cell intracytoplasmic antigen) is a cytoplasmic granule-associated protein, expressed in lymphocytes processing cytolytic potential. TIA-1 is a member of an RNA-binding protein family and possesses nucleolytic activity against cytotoxic lymphocyte (CTL) target cells. It has been suggested that this protein may be involved in the induction of apoptosis as it preferentially recognizes poly(A) homopolymers and induces DNA fragmentation in CTL targets. The major granule-associated species is a 15kDa protein thought to be derived from the carboxyl terminus of the 40kDa product by proteolytic processing. TIA1 antibody labels cytotoxic T cells and natural killer cells (NK cells). It is also expressed in T-cell diseases (low expression) from T-cell lymphocytes (LGL) leukemia and hairy cell leukemia. TIA1 expression in T-cell malignancies.ImmunogenRecombinant human TIA1 fragment of 102 amino acid residues (aa279-380)	Clone	TIA1/1313	
ConjugateAlexa Fluor 647PurityProtein A purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID7072Gene SymbolTIA1SpeciesHumanSpecificity/SensitivityTIA-1 (T-cell intracytoplasmic antigen) is a cytoplasmic granule-associated protein, expressed in lymphocytes processing cytolytic potential. TIA-1 is a member of an RNA-binding protein family and possesses nucleolytic activity against cytotoxic lymphocyte (CTL) target cells. It has been suggested that this protein may be involved in the induction of apoptosis as it preferentially recognizes poly(A) homopolymers and induces DNA fragmentation in CTL targets. The major granule-associated species is a 15kDa protein thought to be derived from the carboxyl terminus of the 40kDa product by proteolytic processing. TIA1 antibody labels cytotoxic cells and natural killer cells (NK cells). It is also expressed in T-cell lymphoma, large granular lymphocyte (LGL) leukemia and hairy cell leukemia. TIA1 expression in T-cell malignancies may help in differentiating LGL leukemia (high expression), TIA1 may also be used to label tumor-infiltrating lymphocytes in the study of immune response to malignancies.ImmunogenRecombinant human TIA1 fragment of 102 amino acid residues (aa279-380)	Preservative	0.05% Sodium Azide	
Purity Protein A purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 7072 Gene Symbol TIA1 Species Human Specificity/Sensitivity TIA-1 (T-cell intracytoplasmic antigen) is a cytoplasmic granule-associated protein, expressed in lymphocytes processing cytolytic potential. TIA-1 is a member of an RNA-binding protein family and possesses nucleolytic activity against cytotoxic lymphocyte (CTL) target cells. It has been suggested that this protein may be involved in the induction of apoptosis as it preferentially recognizes poly(A) homopolymers and induces DNA fragmentation in CTL targets. The major granule-associated species is a 15kDa protein thought to be derived from the carboxyl terminus of the 40kDa product by proteolytic processing. TIA1 antibody labels cytotoxic T cells and natural killer cells (NK cells). It is also expressed in T-cell lymphoma, large granular lymphocyte (LGL) leukemia and hairy cell leukemia. TIA1 expression in T-cell malignancies may help in differentiating LGL leukemia (high expression) from T-cell lymphocytosis and other T-cell diseases (low expression). TIA1 may also be used to label tumor-infiltrating lymphocytes in the study of immune response to malignancies. Immunogen Recombinant human TIA1 fragment of 102 amino acid residues (aa279-380)	Isotype	IgG2b Kappa	
Buffer 50mM Sodium Borate Product Description Host Host Mouse Gene ID 7072 Gene Symbol TIA1 Species Human Specificity/Sensitivity TIA-1 (T-cell intracytoplasmic antigen) is a cytoplasmic granule-associated protein, expressed in lymphocytes processing cytolytic potential. TIA-1 is a member of an RNA-binding protein family and possesses nucleolytic activity against cytotoxic lymphocyte (CTL) target cells. It has been suggested that this protein may be involved in the induction of apoptosis as it preferentially recognizes poly(A) homopolymers and induces DNA fragmentation in CTL targets. The major granule-associated species is a 15kDa protein thought to be derived from the carboxyl terminus of the 40kDa product by proteolytic processing. TIA1 antibody labels cytotoxic T cells and natural killer cells (NK cells). It is also expressed in T-cell lymphoma, large granular lymphocytes (LGL) leukemia and hairy cell leukemia. TIA1 expression) from T-cell lymphocytosis and other T-cell diseases (low expression). TIA1 may also be used to label tumor-infiltrating lymphocytes in the study of immune response to malignancies. Immunogen Recombinant human TIA1 fragment of 102 amino acid residues (aa279-380)	Conjugate	Alexa Fluor 647	
Product Description Host Mouse Gene ID 7072 Gene Symbol TIA1 Species Human Specificity/Sensitivity TIA-1 (T-cell intracytoplasmic antigen) is a cytoplasmic granule-associated protein, expressed in lymphocytes processing cytolytic potential. TIA-1 is a member of an RNA-binding protein family and possesses nucleolytic activity against cytotoxic lymphocyte (CTL) target cells. It has been suggested that this protein may be involved in the induction of apoptosis as it preferentially recognizes poly(A) homopolymers and induces DNA fragmentation in CTL targets. The major granule-associated species is a 15kDa protein thought to be derived from the carboxyl terminus of the 40kDa product by proteolytic processing. TIA1 antibody labels cytotoxic T cells and natural killer cells (NK cells). It is also expressed in T-cell lymphoma, large granular lymphocyte (LGL) leukemia and hairy cell leukemia. TIA1 expression in T-cell malignancies may help in differentiating LGL leukemia (high expression). TIA1 may also be used to label tumor-infiltrating lymphocytes in the study of immune response to malignancies. Immunogen Recombinant human TIA1 fragment of 102 amino acid residues (aa279-380)	Purity	Protein A purified	
HostMouseGene ID7072Gene SymbolTIA1SpeciesHumanSpecificity/SensitivityTIA-1 (T-cell intracytoplasmic antigen) is a cytoplasmic granule-associated protein, expressed in lymphocytes processing cytolytic potential. TIA-1 is a member of an RNA-binding protein family and possesses nucleolytic activity against cytotoxic lymphocyte (CTL) target cells. It has been suggested that this protein may be involved in the induction of apoptosis as it preferentially recognizes poly(A) homopolymers and induces DNA fragmentation in CTL targets. The major granule-associated species is a 15kDa protein thought to be derived from the carboxyl terminus of the 40kDa product by proteolytic processing. TIA1 antibody labels cytotoxic T cells and natural killer cells (NK cells). It is also expressed in T-cell lymphoma, large granular lymphocyte (LGL) leukemia and hairy cell leukemia. TIA1 expression) from T-cell lymphocytosis and other T-cell diseases (low expression). TIA1 may also be used to label tumor-infiltrating lymphocytes in the study of immune response to malignancies.ImmunogenRecombinant human TIA1 fragment of 102 amino acid residues (aa279-380)	Buffer	50mM Sodium Borate	
Gene ID7072Gene SymbolTIA1SpeciesHumanSpecificity/SensitivityTIA-1 (T-cell intracytoplasmic antigen) is a cytoplasmic granule-associated protein, expressed in lymphocytes processing cytolytic potential. TIA-1 is a member of an RNA-binding protein family and possesses nucleolytic activity against cytotoxic lymphocyte (CTL) target cells. It has been suggested that this protein may be involved in the induction of apoptosis as it preferentially recognizes poly(A) homopolymers and induces DNA fragmentation in CTL targets. The major granule-associated species is a 15kDa protein thought to be derived from the carboxyl terminus of the 40kDa product by proteolytic processing. TIA1 antibody labels cytotoxic T cells and natural killer cells (NK cells). It is also expressed in T-cell lymphoma, large granular lymphocyte (LGL) leukemia and hairy cell leukemia (high expression) from T-cell lymphocytosis and other T-cell diseases (low expression). TIA1 may also be used to label tumor-infiltrating lymphocytes in the study of immune response to malignancies.ImmunogenRecombinant human TIA1 fragment of 102 amino acid residues (aa279-380)	Product Description		
Gene SymbolTIA1SpeciesHumanSpecificity/SensitivityTIA-1 (T-cell intracytoplasmic antigen) is a cytoplasmic granule-associated protein, expressed in lymphocytes processing cytolytic potential. TIA-1 is a member of an RNA-binding protein family and possesses nucleolytic activity against cytotoxic lymphocyte (CTL) target cells. It has been suggested that this protein may be involved in the induction of apoptosis as it preferentially recognizes poly(A) homopolymers and induces DNA fragmentation in CTL targets. The major granule-associated species is a 15kDa protein thought to be derived from the carboxyl terminus of the 40kDa product by proteolytic processing. TIA1 antibody labels cytotoxic T cells and natural killer cells (NK cells). It is also expressed in T-cell lymphoma, large granular lymphocyte (LGL) leukemia and hairy cell leukemia. TIA1 expression in T-cell malignancies may help in differentiating LGL leukemia (high expression) from T-cell lymphocytosis and other T-cell diseases (low expression). TIA1 may also be used to label tumor-infiltrating lymphocytes in the study of immune response to malignancies.ImmunogenRecombinant human TIA1 fragment of 102 amino acid residues (aa279-380)	Host	Mouse	
SpeciesHumanSpecificity/SensitivityTIA-1 (T-cell intracytoplasmic antigen) is a cytoplasmic granule-associated protein, expressed in lymphocytes processing cytolytic potential. TIA-1 is a member of an RNA-binding protein family and possesses nucleolytic activity against cytotoxic lymphocyte (CTL) target cells. It has been suggested that this protein may be involved in the induction of apoptosis as it preferentially recognizes poly(A) homopolymers and induces DNA fragmentation in CTL targets. The major granule-associated species is a 15kDa protein thought to be derived from the carboxyl terminus of the 40kDa product by proteolytic processing. TIA1 antibody labels cytotoxic T cells and natural killer cells (NK cells). It is also expressed in T-cell lymphoma, large granular lymphocyte (LGL) leukemia and hairy cell leukemia. TIA1 expression in T-cell malignancies may help in differentiating LGL leukemia (high expression) from T-cell lymphocytosis and other T-cell diseases (low expression). TIA1 may also be used to label tumor-infiltrating lymphocytes in the study of immune response to malignancies.ImmunogenRecombinant human TIA1 fragment of 102 amino acid residues (aa279-380)	Gene ID	7072	
Specificity/SensitivityTIA-1 (T-cell intracytoplasmic antigen) is a cytoplasmic granule-associated protein, expressed in lymphocytes processing cytolytic potential. TIA-1 is a member of an RNA-binding protein family and possesses nucleolytic activity against cytotoxic lymphocyte (CTL) target cells. It has been suggested that this protein may be involved in the induction of apoptosis as it preferentially recognizes poly(A) homopolymers and induces DNA fragmentation in CTL targets. The major granule-associated species is a 15kDa protein thought to be derived from the carboxyl terminus of the 40kDa product by proteolytic processing. TIA1 antibody labels cytotoxic T cells and natural killer cells (NK cells). It is also expressed in T-cell lymphoma, large granular lymphocyte (LGL) leukemia and hairy cell leukemia. TIA1 expression in T-cell malignancies may help in differentiating LGL leukemia (high expression) from T-cell lymphocytosis and other T-cell diseases (low expression). TIA1 may also be used to label tumor-infiltrating lymphocytes in the study of immune response to malignancies.ImmunogenRecombinant human TIA1 fragment of 102 amino acid residues (aa279-380)	Gene Symbol	TIA1	
protein, expressed in lymphocytes processing cytolytic potential. TIA-1 is a member of an RNA-binding protein family and possesses nucleolytic activity against cytotoxic lymphocyte (CTL) target cells. It has been suggested that this protein may be involved in the induction of apoptosis as it preferentially recognizes poly(A) homopolymers and induces DNA fragmentation in CTL targets. The major granule-associated species is a 15kDa protein thought to be derived from the carboxyl terminus of the 40kDa product by proteolytic processing. TIA1 antibody labels cytotoxic T cells and natural killer cells (NK cells). It is also expressed in T-cell lymphoma, large granular lymphocyte (LGL) leukemia and hairy cell leukemia. TIA1 expression in T-cell malignancies may help in differentiating LGL leukemia (high expression) from T-cell lymphocytosis and other T-cell diseases (low expression). TIA1 may also be used to label tumor-infiltrating lymphocytes in the study of immune response to malignancies.ImmunogenRecombinant human TIA1 fragment of 102 amino acid residues (aa279-380)	Species	Human	
	Specificity/Sensitivity	protein, expressed in lymphocytes processing cytolytic potential. TIA-1 is a member of an RNA-binding protein family and possesses nucleolytic activity against cytotoxic lymphocyte (CTL) target cells. It has been suggested that this protein may be involved in the induction of apoptosis as it preferentially recognizes poly(A) homopolymers and induces DNA fragmentation in CTL targets. The major granule-associated species is a 15kDa protein thought to be derived from the carboxyl terminus of the 40kDa product by proteolytic processing. TIA1 antibody labels cytotoxic T cells and natural killer cells (NK cells). It is also expressed in T-cell lymphoma, large granular lymphocyte (LGL) leukemia and hairy cell leukemia. TIA1 expression in T-cell malignancies may help in differentiating LGL leukemia (high expression) from T-cell lymphocytosis and other T-cell diseases (low expression). TIA1 may also be used to label	
	Immunogen		



	C I
Notes	Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.
Product Application Details	
Applications	Western Blot, Flow Cytometry, Flow (Intracellular), Immunocytochemistry/ Immunofluorescence, CyTOF-ready
Recommended Dilutions	Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Flow (Intracellular), CyTOF-ready
Application Notes	Optimal dilution of this antibody should be experimentally determined.
4	





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-54406AF647

Mouse IgG2b Kappa Light Chain Isotype Control (MG2b) [Alexa Fluor® 647]
Recombinant Human TIA1 GST (N-Term) Protein
TNF-alpha [Unconjugated]
TIA1 Overexpression Lysate

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-54406AF647

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

