Product Datasheet

EpCAM/TROP1 Antibody (EGP40/1556R) [Alexa Fluor® 750] - (ECD) NBP2-54425AF750

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-54425AF750

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.



Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-54425AF750

NBP2-54425AF750

EpCAM/TROP1 Antibody (EGP40/1556R) [Alexa Fluor® 750] - (ECD)

Product InformationUnit Size0.1 mlConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneEGP40/1556RPreservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 750PurityProtein A purifiedBufferSomM Sodium BorateProduct Description4072Gene ID4072Gene IDEPCAMSpeciesHuman, Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse, Canine, Feline or Rat.MarkerEpithelial MarkerSpecificity/SensitivityEGP40 is a 40-43KDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (EpCAM). It is expressed on baso-lateral cell sufface in most simple epithelia and a vast majority of carcinomas from pleural mesothelionar and hepatocellular arcarcinoma. This antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelioma.ImmunogenRecombinant human full-length EpCAM/TROP1 protein (Uniprot: P16422)			
ConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneEGP40/1556RPreservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 750PurityProtein A purifiedBuffer50mM Sodium BorateProduct DescriptionHostRabbitGene ID4072Gene SymbolEPCAMSpeciesHuman, Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse, Canine, Feline or Rat.MarkerEpithelial MarkerSpecificity/SensitivityEGP40 is a 40-43XDa transmembrane epithelial cellular adhesion molecule (EpCAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majority of carcinomas with the exception of adult squamous epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (EpCAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majority of carcinomas with the exception of adult squamous epithelial specific antigen (ESA), or epithelial cellus This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. This antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelioma.	Product Information		
services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneEGP40/1556RPreservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 750PurityProtein A purifiedBuffer50mM Sodium BorateProduct Description4072Gene ID4072Gene SymbolEPCAMSpeciesHuman, Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse, Canine, Feline or Rat.MarkerEGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific artigen (ESA), or epithelia cellular adhesion molecule (EpCAM). It is expressed on baso-lateral cell surface in most simple epithelial and a vast majority of carcinomas with the exception of adult squamous epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (EpCAM). It is expressed on baso-lateral cell surface in most simple epithelial and a vast majority of carcinomas with the exception of adult squamous epithelian and a vast majority of carcinomas with the exception of adult squamous epithelian denocarcinoma from pleural mesothelioma and hepatocellular carcinoma. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma.	Unit Size	0.1 ml	
ClonalityMonoclonalCloneEGP40/1556RPreservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 750PurityProtein A purifiedBuffer50mM Sodium BorateProduct DescriptionHostRabbitGene ID4072Gene SymbolEPCAMSpeciesHuman, Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse, Canine, Feline or Rat.MarkerEpithelial MarkerSpecificity/SensitivityEGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (EpCAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majority of carcinomas with the exception of adult squamous epithelian and a vast majority of salso useful in distinguishing serous carcinomas of the ovary from mesothelioma.	Concentration		
CloneEGP40/1556RPreservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 750PurityProtein A purifiedBuffer50mM Sodium BorateProduct DescriptionHostRabbitGene ID4072Gene SymbolEPCAMSpeciesHuman, Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse, Canine, Feline or Rat.MarkerEGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (EpCAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majority of carcinomas after epithelial cells. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinomas of the ovary from mesothelioma.	Storage	Store at 4C in the dark.	
Preservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 750PurityProtein A purifiedBuffer50mM Sodium BorateProduct Description4072HostRabbitGene ID4072Gene SymbolEPCAMSpeciesHuman, Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse, Canine, Feline or Rat.MarkerEpT40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific atigen (ESA), or epithelial cellular adhesion molecule (EpCAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majority of carcinomas with the exception of adult squamous epithelial adenocarcinoma from pleural mesothelioma and hepatocellular carcinomas of the ovary from mesothelioma.	Clonality	Monoclonal	
isotypeIgGConjugateAlexa Fluor 750PurityProtein A purifiedBuffer50mM Sodium BorateProduct Description4072HostRabbitGene ID4072Gene SymbolEPCAMSpeciesHuman, Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse, Canine, Feline or Rat.MarkerEpithelial MarkerSpecificity/SensitivityEGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (EpCAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majority of carcinomas with the exception of adult squamous epithelium, hepatocytes and gastric epithelial cells. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. This antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelioma.	Clone	EGP40/1556R	
ConjugateAlexa Fluor 750PurityProtein A purifiedBuffer50mM Sodium BorateProduct DescriptionHostRabbitGene ID4072Gene SymbolEPCAMSpeciesHuman, Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse, Canine, Feline or Rat.MarkerEgP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (EpCAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majority of carcinomas with the exception of adult squamous epithelium, hepatocytes and gastric epithelial cells. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. This antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelioma.	Preservative	0.05% Sodium Azide	
PurityProtein A purifiedBuffer50mM Sodium BorateProduct DescriptionHostRabbitGene ID4072Gene SymbolEPCAMSpeciesHuman, Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse, Canine, Feline or Rat.MarkerEpithelial MarkerSpecificity/SensitivityEGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (EpCAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majority of carcinomas with the exception of adult squamous epithelium, hepatocytes and gastric epithelial cells. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. This antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelioma.	Isotype	IgG	
Buffer50mM Sodium BorateProduct DescriptionHostRabbitGene ID4072Gene SymbolEPCAMSpeciesHuman, Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse, Canine, Feline or Rat.MarkerEpithelial MarkerSpecificity/SensitivityEGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (EpCAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majority of carcinomas with the exception of adult squamous epithelium, hepatocytes and gastric epithelial cells. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinomas. This antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelioma.	Conjugate	Alexa Fluor 750	
Product DescriptionHostRabbitGene ID4072Gene SymbolEPCAMSpeciesHuman, Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse, Canine, Feline or Rat.MarkerEpithelial MarkerSpecificity/SensitivityEGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (EpCAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majority of carcinomas with the exception of adult squamous epitheliam, hepatocytes and gastric epithelial cells. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. This antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelioma.	Purity	Protein A purified	
HostRabbitGene ID4072Gene SymbolEPCAMSpeciesHuman, Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse, Canine, Feline or Rat.MarkerEpithelial MarkerSpecificity/SensitivityEGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (EpCAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majority of carcinomas with the exception of adult squamous epithelium, hepatocytes and gastric epithelial cells. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. This antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelioma.	Buffer	50mM Sodium Borate	
Gene ID4072Gene SymbolEPCAMSpeciesHuman, Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse, Canine, Feline or Rat.MarkerEpithelial MarkerSpecificity/SensitivityEGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (EpCAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majority of carcinomas with the exception of adult squamous epithelium, hepatocytes and gastric epithelial cells. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. This antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelioma.	Product Description		
Gene SymbolEPCAMSpeciesHuman, Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse, Canine, Feline or Rat.MarkerEpithelial MarkerSpecificity/SensitivityEGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (EpCAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majority of carcinomas with the exception of adult squamous epithelium, hepatocytes and gastric epithelial cells. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. This antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelioma.	Host	Rabbit	
SpeciesHuman, Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse, Canine, Feline or Rat.MarkerEpithelial MarkerSpecificity/SensitivityEGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (EpCAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majority of carcinomas with the exception of adult squamous epithelium, hepatocytes and gastric epithelial cells. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. This antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelioma.	Gene ID	4072	
Reactivity NotesDoes not react with Mouse, Canine, Feline or Rat.MarkerEpithelial MarkerSpecificity/SensitivityEGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (EpCAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majority of carcinomas with the exception of adult squamous epithelium, hepatocytes and gastric epithelial cells. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. This antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelioma.	Gene Symbol	EPCAM	
MarkerEpithelial MarkerSpecificity/SensitivityEGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (EpCAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majority of carcinomas with the exception of adult squamous epithelium, hepatocytes and gastric epithelial cells. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. This antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelioma.	Species	Human, Mouse (Negative), Rat (Negative)	
Specificity/Sensitivity EGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (EpCAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majority of carcinomas with the exception of adult squamous epithelium, hepatocytes and gastric epithelial cells. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinomas. This antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelioma.	Reactivity Notes	Does not react with Mouse, Canine, Feline or Rat.	
epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (EpCAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majority of carcinomas with the exception of adult squamous epithelium, hepatocytes and gastric epithelial cells. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. This antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelioma.	Marker	Epithelial Marker	
ImmunogenRecombinant human full-length EpCAM/TROP1 protein (Uniprot: P16422)	Specificity/Sensitivity	epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (EpCAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majority of carcinomas with the exception of adult squamous epithelium, hepatocytes and gastric epithelial cells. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. This antibody is also useful in distinguishing serous carcinomas of	
	Immunogen	Recombinant human full-length EpCAM/TROP1 protein (Uniprot: P16422)	



	Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.
Product Application Details	
Applications	Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready
Recommended Dilutions	Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, CyTOF-ready
Application Notes	Optimal dilution of this antibody should be experimentally determined.

Notes





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-54425AF750

960-EP-050	EpCAM/TROP1 [Unconjugated]
202-IL-010	IL-2 [Unconjugated]
NBP2-52190-0.05mg	Recombinant Human EpCAM/TROP1 His Protein
NBP2-24891AF750	Rabbit IgG Isotype Control [Alexa Fluor® 750]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-54425AF750

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

