# **Product Datasheet**

# Von Willebrand Factor Antibody (VWF/1465) [Alexa Fluor® 532] NBP2-54430AF532

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-54430AF532

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP2-54430AF532



# NBP2-54430AF532

Von Willebrand Factor Antibody (VWF/1465) [Alexa Fluor® 532]

Product Information	Voli Willebrahd Factor Antibody (VWF/1403) [Alexa Fidol® 532]		
Concentration Please see the vial label for concentration. If unlisted please contact technical services.  Storage Store at 4C in the dark.  Clonality Monoclonal Clone VWF/1465 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate Alexa Fluor 532 Purity Protein A or G purified Buffer 50mM Sodium Borate  Product Description Host Mouse Gene ID 7450 Gene Symbol VWF Species Human Marker Endothelial Marker Specificity/Sensitivity Von Willebrand Factor (VWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. WWF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposis sarcoma acradiac mysoma. It is widely used for differentiating vascular markers although not all tumors of endothelial differentiation contain this antigen.  Immunogen Recombinant fragment of human Von Willebrand Factor protein (aa1815-1939)	Product Information		
Storage Store at 4C in the dark.  Clonality Monoclonal Clone VWF/1465 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate Alexa Fluor 532 Purity Protein A or G purified Buffer 50mM Sodium Borate  Product Description Host Mouse Gene ID 7450 Gene Symbol VWF Species Huma Marker Endothelial Marker  Specificity/Sensitivity von Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslational modifications that influence haffinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposis sarcoma and cardiac myxoma. It is widely used for differentiating vascular lesions from those of other tissue differentiation within a panel of other vascular markers although not all tumors of endothelial differentiation contain this antilgen.  Immunogen Recombinant fragment of human Von Willebrand Factor protein (aa1815-1939)	Unit Size	0.1 ml	
Clone VWF/1465 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate Alexa Fluor 532 Purity Protein A or G purified Buffer 50mM Sodium Borate  Product Description Host Mouse Gene ID 7450 Gene Symbol VWF Species Human Marker Endothelial Marker Specificity/Sensitivity von Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. wWF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposis sarcoma and cardiac myxoma. It is widely used for differentiating vascular lesions from those of other tissue differentiation within a panel of other vascular markers although not all tumors of endothelial differentiation contain this antigen.  Immunogen Recombinant fragment of human Von Willebrand Factor protein (aa1815-1939)	Concentration	'	
Clone VWF/1465 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate Alexa Fluor 532 Purity Protein A or G purified Buffer 50mM Sodium Borate  Product Description Host Mouse Gene ID 7450 Gene Symbol VWF Species Human Marker Endothelial Marker Specificity/Sensitivity von Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposis sarcoma and cardiac myxoma. It is widely used for differentiation yascular lesions from those of other tissue differentiation within a panel of other vascular markers although not all tumors of endothelial differentiation contain this antigen.  Immunogen Recombinant fragment of human Von Willebrand Factor protein (aa1815-1939)	Storage	Store at 4C in the dark.	
Preservative 0.05% Sodium Azide  Isotype	Clonality	Monoclonal	
Isotype	Clone	VWF/1465	
Conjugate Alexa Fluor 532  Purity Protein A or G purified  Buffer 50mM Sodium Borate  Product Description  Host Mouse  Gene ID 7450  Gene Symbol VWF  Species Human  Marker Endothelial Marker  Specificity/Sensitivity von Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposis sarcoma and cardiac myxoma. It is widely used for differentiating vascular markers although not all tumors of endothelial differentiation contain this antigen.  Immunogen Recombinant fragment of human Von Willebrand Factor protein (aa1815-1939)	Preservative	0.05% Sodium Azide	
Purity Protein A or G purified  Buffer 50mM Sodium Borate  Product Description  Host Mouse  Gene ID 7450  Gene Symbol VWF  Species Human  Marker Endothelial Marker  Specificity/Sensitivity von Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposis sarcoma and cardiac myxoma. It is widely used for differentiating vascular lesions from those of other tissue differentiation within a panel of other vascular markers although not all tumors of endothelial differentiation contain this antigen.  Immunogen Recombinant fragment of human Von Willebrand Factor protein (aa1815-1939)	Isotype	IgG1 Kappa	
Buffer 50mM Sodium Borate  Product Description  Host Mouse  Gene ID 7450  Gene Symbol VWF  Species Human  Marker Endothelial Marker  Specificity/Sensitivity von Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposis sarcoma and cardiac myxoma. It is widely used for differentiating vascular lesions from those of other tissue differentiation within a panel of other vascular markers although not all tumors of endothelial differentiation contain this antigen.  Immunogen Recombinant fragment of human Von Willebrand Factor protein (aa1815-1939)	Conjugate	Alexa Fluor 532	
Product Description  Host Mouse  Gene ID 7450  Gene Symbol VWF  Species Human  Marker Endothelial Marker  Specificity/Sensitivity von Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposis sarcoma and cardiac myxoma. It is widely used for differentiating vascular lesions from those of other tissue differentiation within a panel of other vascular markers although not all tumors of endothelial differentiation contain this antigen.  Immunogen Recombinant fragment of human Von Willebrand Factor protein (aa1815-1939)	Purity	Protein A or G purified	
Host Gene ID 7450 Gene Symbol VWF Species Human Endothelial Marker Specificity/Sensitivity von Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposis sarcoma and cardiac myxoma. It is widely used for differentiating vascular lesions from those of other tissue differentiation within a panel of other vascular markers although not all tumors of endothelial differentiation contain this antigen.  Immunogen Recombinant fragment of human Von Willebrand Factor protein (aa1815-1939)	Buffer	50mM Sodium Borate	
Gene Symbol  VWF  Species  Human  Endothelial Marker  Specificity/Sensitivity  von Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposis sarcoma and cardiac myxoma. It is widely used for differentiating vascular lesions from those of other tissue differentiation within a panel of other vascular markers although not all tumors of endothelial differentiation contain this antigen.  Immunogen  Recombinant fragment of human Von Willebrand Factor protein (aa1815-1939)	Product Description		
Gene Symbol  VWF  Species  Human  Endothelial Marker  Specificity/Sensitivity  von Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposis sarcoma and cardiac myxoma. It is widely used for differentiating vascular lesions from those of other tissue differentiation within a panel of other vascular markers although not all tumors of endothelial differentiation contain this antigen.  Immunogen  Recombinant fragment of human Von Willebrand Factor protein (aa1815-1939)	Host	Mouse	
Species  Human  Endothelial Marker  Specificity/Sensitivity  von Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposis sarcoma and cardiac myxoma. It is widely used for differentiating vascular lesions from those of other tissue differentiation within a panel of other vascular markers although not all tumors of endothelial differentiation contain this antigen.  Immunogen  Recombinant fragment of human Von Willebrand Factor protein (aa1815-1939)	Gene ID	7450	
Marker  Specificity/Sensitivity  von Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposis sarcoma and cardiac myxoma. It is widely used for differentiating vascular lesions from those of other tissue differentiation within a panel of other vascular markers although not all tumors of endothelial differentiation contain this antigen.  Immunogen  Recombinant fragment of human Von Willebrand Factor protein (aa1815-1939)	Gene Symbol	VWF	
Specificity/Sensitivity  von Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposis sarcoma and cardiac myxoma. It is widely used for differentiating vascular lesions from those of other tissue differentiation within a panel of other vascular markers although not all tumors of endothelial differentiation contain this antigen.  Immunogen  Recombinant fragment of human Von Willebrand Factor protein (aa1815-1939)	Species	Human	
endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposis sarcoma and cardiac myxoma. It is widely used for differentiating vascular lesions from those of other tissue differentiation within a panel of other vascular markers although not all tumors of endothelial differentiation contain this antigen.  Immunogen  Recombinant fragment of human Von Willebrand Factor protein (aa1815-1939)	Marker	Endothelial Marker	
		endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposis sarcoma and cardiac myxoma. It is widely used for differentiating vascular lesions from those of other tissue differentiation within a panel of other vascular markers although not all tumors of endothelial differentiation contain this antigen.	
	Immunogen		



	Page 2 of 3 V.20.1 Updated 10/23/2024
	Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.
Product Application Details	
	Western Blot, Flow Cytometry, Immunohistochemistry, Immunohistochemistry-Paraffin, Protein Array, CyTOF-ready

Product Application Details	
Applications	Western Blot, Flow Cytometry, Immunohistochemistry, Immunohistochemistry-Paraffin, Protein Array, CyTOF-ready
Recommended Dilutions	Western Blot, Flow Cytometry, Immunohistochemistry, Immunohistochemistry-Paraffin, Protein Array, CyTOF-ready
Application Notes	Optimal dilution of this antibody should be experimentally determined.





### **Novus Biologicals USA**

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom

Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

## Products Related to NBP2-54430AF532

NBP1-84762PEP Von Willebrand Factor Recombinant Protein Antigen

210-TA-005 TNF-alpha [Unconjugated]

KA0512 Human Von Willebrand Factor ELISA Kit (Colorimetric)

DVE00 VEGF [HRP]

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-54430AF532

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

