# **Product Datasheet**

# Erythropoietin/EPO Antibody (EPO/1367) - Azide and BSA Free NBP2-54447-100ug

Unit Size: 100 ug

Store at -20 to -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-54447

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP2-54447




## NBP2-54447-100ug

| Erythropoietin/EPO Antibody (EPO/1367) - Azide and BSA Free |  |
|---|--|
| Product Information   |  |
| Unit Size   | 100 ug   |
| Concentration   | 1 mg/ml  |
| Storage   | Store at -20 to -80C. Avoid freeze-thaw cycles.  |
| Clonality   | Monoclonal   |
| Clone   | EPO/1367   |
| Preservative  | No Preservative  |
| Isotype   | IgG1 Kappa   |
| Purity  | Protein A or G purified  |
| Buffer  | 10 mM PBS  |
| Target Molecular Weight                                     | 37 kDa   |
| Product Description   |  |
| Description   | 1.0 mg/ml of antibody purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WITH BSA & azide (NBP2-53241).  Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C.  |
| Host  | Mouse  |
| Gene ID   | 2056   |
| Gene Symbol   | EPO  |
| Species   | Human  |
| Marker  | Marker of Placentation Disorders   |
| Immunogen   | Recombinant fragment (around aa 28-162) of human Erythropoietin/EPO protein (exact sequence is proprietary) (Uniprot: P01588)  |
| Product Application Details                                 |  |
| Applications  | ELISA, Immunohistochemistry, Immunohistochemistry-Paraffin, Protein Array  |
| Recommended Dilutions                                       | ELISA 2-4 ug/ml, Immunohistochemistry, Immunohistochemistry-Paraffin, Protein Array  |
| Application Notes   | ELISA: Use Ab at 2-4ug/ml for coating. Order Ab without BSA. Immunohistochemistry (Formalin-fixed): 2-4ug/ml for 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes. Optimal dilution for a specific application should be determined. |




#### **Images**


Immunohistochemistry-Paraffin: Erythropoietin/EPO Antibody (EPO/1367) - Azide and BSA Free [NBP2-54447] - Formalin-fixed, paraffin-embedded Human Colon Carcinoma stained with Erythropoietin (EPO) Mouse Monoclonal Antibody (EPO/1367).



Immunohistochemistry-Paraffin: Erythropoietin/EPO Antibody (EPO/1367) - Azide and BSA Free [NBP2-54447] - Formalin-fixed, paraffin-embedded Human Renal Cell Carcinoma stained with Erythropoietin (EPO) Mouse Monoclonal Antibody (EPO/1367).



Protein Array: Erythropoietin/EPO Antibody (EPO/1367) - Azide and BSA Free [NBP2-54447] - Analysis of Protein Array containing more than 19,000 full-length human proteins using Erythropoietin/EPO Antibody (EPO/1367).Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5.





### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

## Products Related to NBP2-54447-100ug

HAF007 Goat anti-Mouse IgG Secondary Antibody [HRP]

NB720-B Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]

NBP1-43319-0.5mg Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1)
NBP2-62447 Recombinant Porcine Erythropoietin/EPO Protein

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-54447

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

