# **Product Datasheet**

# PMEL17/SILV Antibody (PMEL/1825R) - Azide and BSA Free NBP2-54454-100ug

Unit Size: 100 ug

Store at -20 to -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-54454

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-54454



## NBP2-54454-100ug

PMEL17/SILV Antibody (PMEL/1825R) - Azide and BSA Free

0251() - Azide and BOA i ree	
Product Information	
100 ug	
1 mg/ml	
Store at -20 to -80C. Avoid freeze-thaw cycles.	
Monoclonal	
PMEL/1825R	
No Preservative	
IgG	
Protein A or G purified	
10 mM PBS	
95 kDa	
1.0 mg/ml of antibody purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WITH BSA & azide (NBP2-53180).  Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80 C.	
Rabbit	
6490	
PMEL	
Human	
Does not react with Canine and Rat	
Melanoma Marker	
The gp100 molecule is a 100kDa glycosylated protein that is cleaved into a small (26kDa) carboxy-terminal fragment and a larger amino- terminal section (60 64 kDa), which is subsequently cleaved to generate 26kDa and 34 38kDa fragments. By immunohistochemistry, it specifically recognizes a protein in melanocytes and melanomas. This monoclonal antibody reacts with junctional and blue nevus cells and variably with fetal and neonatal melanocytes. Intradermal nevi, normal adult melanocytes, and non-melanocytic cells are negative. It does not stain tumor cells of epithelial, lymphoid, glial, or mesenchymal origin. Metastatic amelanotic melanoma can often be confused with a variety of poorly differentiated carcinomas, large cell lymphomas, and sarcomas using H E stains alone. It is also difficult to differentiate melanoma from spindle cell carcinomas and various types of mesenchymal neoplasms. This monoclonal antibody stains fetal and neonatal melanocytes, junctional and blue nevus cells, and malignant melanoma. This monoclonal antibody also stains Angiomyolipoma (PEComa).	
Recombinant full-length human PMEL17/SILV protein (Uniprot: P40967)	
Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin	
Western Blot 2-4 ug/ml, Immunohistochemistry, Immunohistochemistry-Paraffin 0.5 - 1.0 ug/ml	




#### **Application Notes**


Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes. Optimal dilution for a specific application should be determined.

#### **Images**

Western Blot: PMEL17/SILV Antibody (PMEL/1825R) - Azide and BSA Free [NBP2-54454] - Western Blot Analysis of COLO-38 cell lysate using PMEL17/SILV Rabbit Recombinant Monoclonal Antibody (PMEL/1825R).



Immunohistochemistry-Paraffin: PMEL17/SILV Antibody (PMEL/1825R) - Azide and BSA Free [NBP2-54454] - Formalin-fixed, paraffin-embedded human Melanoma stained with gp100 Recombinant Rabbit Monoclonal Antibody (PMEL/1825R).





### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

## Products Related to NBP2-54454-100ug

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

NBP2-38185PEP PMEL17/SILV Recombinant Protein Antigen

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-54454

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

