## **Product Datasheet**

### B220/CD45R Antibody (PTPRC/1460) [PE/Atto594] NBP2-54465PEATT594

Unit Size: 0.1 ml

Store at 4C in the dark. Do not freeze.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-54465PEATT594

Updated 10/23/2024 v.20.1

# Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-54465PEATT594



#### NBP2-54465PEATT594

B220/CD45R Antibody (PTPRC/1460) [PE/Atto594]

or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixation		
Concentration         Please see the vial label for concentration. If unlisted please contact technical services.           Storage         Store at 4C in the dark. Do not freeze.           Clonality         Monoclonal           Clone         PTPRC/1460           Preservative         0.05% Sodium Azide           Isotype         IgG1 Kappa           Conjugate         PE/Atto594           Purity         Protein A or G purified           Buffer         PBS           Product Description         Mouse           Gene ID         5788           Gene Symbol         PTPRC           Species         Human           Marker         B-Cell Marker           Species         Human           Marker         B-Cell Marker           Specificity/Sensitivity         CD45R, also designated CD45 and PTPRC, has been identified as a transmembrane givcoprotein, broady expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms anse because of alternative splicing of exora 5, 5, and 6. The corresponding protein domains are characterized by the binding of monochonal antibodies specific for CD45RA (exon 4). CD45RB (exon 0). SD45RB (exon 0). S	Product Information	
services.           Storage         Store at 4C in the dark. Do not freeze.           Clonality         Monoclonal           Clone         PTPRC/1460           Preservative         0.05% Sodium Azide           Isotype         IgG1 Kappa           Conjugate         PE/Atto594           Purity         Protein A or G purified           Buffer         PBS           Product Description         Mouse           Host         Mouse           Gene ID         5788           Gene Symbol         PTPRC           Species         Human           Marker         B-Cell Marker           Specificity/Sensitivity         CD45R, also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms arise because of alternatetized by the binding of monochoral antibodies specific for CD45RA (scon 4). CD45RB (scon 5). CD45RC (scon 6) and CD45RO (scon 4 to 6 spliced out). The variation in these isoforms is localized to the extracellular domain of CD45RB (scon 5). CD45RC (scon 6) and CD45RO (scon 4 to 6 spliced out). The variation in these isoforms is localized to the extracellular domain of CD45RB (scon 5). CD45RC (scon 6) and CD45RO (scon 4 to 6 spliced out). The variation in these isoforms is localized to the extracellular domain of CD45RB (scon 5). CD45RC (scon 6) and CD45RO (scon 4 to 6 spliced out). The variation in these isofor	Unit Size	0.1 ml
Clonality       Monoclonal         Clone       PTPRC/1460         Preservative       0.05% Sodium Azide         Isotype       IgG1 Kappa         Conjugate       PE/Atto594         Purity       Protein A or G purified         Buffer       PBS         Product Description       Mouse         Gene ID       5788         Gene Symbol       PTPRC         Species       Human         Marker       B-Cell Marker         Specificity/Sensitivity       CD45R, also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms arise because of alternative splicing of exons 4, 5, and 6. The corresponding protein domains are characterized by the intracellular domain is conserved. CD45R functions as a phosphort-yrosine phosphatase. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic undifferentiated neoplasms.         Immunogen       Human B-lymphocytes (Uniprot: P08575)         Product Application Details       Flow Cytometry, Immunohistochemistry-Parafin, CyTOF-ready         Application Notes       Flow Cytometry, Immunohistochemistry-Parafin, CyTOF-ready         Application Notes       Flow Cytometry, Immunohistochemistry, Parafin, CyTOF-ready         Application Notes       Flow Cytometry, I	Concentration	•
Clone         PTPRC/1460           Preservative         0.05% Sodium Azide           Isotype         IgG1 Kappa           Conjugate         PE/Atto594           Purity         Protein A or G purified           Buffer         PBS           Product Description            Host         Mouse           Gene ID         5788           Gene Symbol         PTPRC           Specificity/Sensitivity         CO45R, also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R and PTPRC and a propertien domains are characterized by the be intracellular domain is conserved. CD45R Netwite the intracellular domain is conserved. CD45R functions as a phosphortyrosine phosphatase. Antibody to CD45 is useful in differentiated neoplasms.           Immunogen         Human B-lymphocytes (Uniprot: P08575)           Product Application Details         Flow Cytometry, Immunohistochemistry-Paraffin, CyTOF-ready           Recommended Dilutions         Flow Cytometry, Immunohistochemistry-Paraffin, CyTOF-ready           Recommended Dilut	Storage	Store at 4C in the dark. Do not freeze.
Preservative         0.05% Sodium Azide           Isotype         IgG1 Kappa           Conjugate         PE/Atto594           Purity         Protein A or G purified           Buffer         PBS           Product Description         Mouse           Gene ID         5788           Gene Symbol         PTPRC           Species         Human           Marker         B-Cell Marker           Specificity/Sensitivity         CD45R also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms arise because of alternative splicing of exons 4, 5, and 6. The corresponding protein domains are characterized by the binding of monoclonal antibodies specific for CD45R (exon 4), CD45RB (exon 5), CD45RC (exon 6) and CD45R (exon 4), CD45RB (exon 5), CD45RC (exon 6) and CD45R (ustor 4 to 6 spliced out). The variation in these isoforms is localized to the extracellular domain of CD45R, while the intracellular domain is conserved. CD45R functions as a phosphor-tyrosine phosphatase. Antibody to CD45 is usedui in differential diagnosis of lymphoid tumors from non-hematopoietic undifferentiated neoplasms.           Immunogen         Human B-lymphocytes (Uniprot: P08575)           Product Application Details         Flow Cytometry, Immunohistochemistry-Paraffin, CyTOF-ready           Application Notes         Optimal dilution of this antibody should be experimentally determined. For opti	Clonality	Monoclonal
Isotype       IgG1 Kappa         Conjugate       PE/Atto594         Purity       Protein A or G purified         Buffer       PBS         Product Description       Mouse         Gene ID       5788         Gene Symbol       PTPRC         Species       Human         Marker       B-Cell Marker         Specificity/Sensitivity       CD45R, also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms arise because of alternative splicing of exons 4, 5, and 6. The corresponding protein domains are characterized by the binding of monoclonal antibodies specific for CD45RA (scon 4), CD45RB (scon 5), CD45RC (exon 6) and CD45RO (exons 4 to 6 spliced out). The variation in these isoforms is localized to the extracellular domain is conserved. CD45R functions as a phosphor-tyrosine phosphatase. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic undifferential diagnos	Clone	PTPRC/1460
Conjugate         PE/Atto594           Purity         Protein A or G purified           Buffer         PBS           Product Description           Host         Mouse           Gene ID         5788           Gene Symbol         PTPRC           Species         Human           Marker         B-Cell Marker           Specificity/Sensitivity         CD45R, also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms arise because of alternative splicing of exons 4, 5, and 6. The corresponding protein domains are characterized by the binding of monocional antibodies specific for CD45R, dexon 4), CD45RB (exon 5), CD45RC (exon 6) and CD45R (unctions as a phosphor-tyrosine phosphatase. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic undifferentiated neoplasms.           Immunogen         Human B-lymphocytes (Uniprot: P08575)           Product Application Details         Flow Cytometry, Immunohistochemistry-Paraffin, CyTOF-ready           Recommended Dilutions         Flow Cytometry, Immunohistochemistry-Paraffin, CyTOF-ready           Application Notes         Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to light           Gene D         Optimal dilution of this antibody should be experimentally	Preservative	0.05% Sodium Azide
Purity         Protein A or G purified           Buffer         PBS           Product Description         Mouse           Gene ID         5788           Gene Symbol         PTPRC           Species         Human           Marker         B-Cell Marker           Specificity/Sensitivity         CD45R, also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoletic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms arise because of alternative splicing of exons 4, 5, and 6. The corresponding protein domains are characterized by the binding of monoclonal antibodies specific for CD45RA (exon 4), CD45RB (exon 5), CD45RC (exon 6) and CD45RO (exons 4 to 6 spliced out). The variation in these isoforms is localized to the extracellular domain of CD45R, while the intracellular domain is conserved. CD45R functions as a phosphor-tyrosine phosphatase. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic undifferentiated neoplasms.           Immunogen         Human B-lymphocytes (Uniprot: P08575)           Product Application Details         Flow Cytometry, Immunohistochemistry/ Immunofluorescence, Immunohistochemistry, Paraffin, CyTOF-ready           Recommended Dilutions         Flow Cytometry, Immunohistochemistry, Paraffin, CyTOF-ready           Application Notes         Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to light or extreme temperature fluc	Isotype	IgG1 Kappa
Buffer         PBS           Product Description         Mouse           Gene ID         5788           Gene Symbol         PTPRC           Species         Human           Marker         B-Cell Marker           Specificity/Sensitivity         CD45R, also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms arise because of alternative splicing of exores 4, 5, and 6. The corresponding protein domains are characterized by the binding of monoclonal antibodies specific for CD45RA (exon 4), CD45RB (exon 5), CD45RC (exon 6) and CD45RO (exons 4 to 6 spliced out). The variation in these isoforms is localized to the extracellular domain of CD45R, while the intracellular domain is conserved. CD45R functions as a phosphor-tyrosine phosphatase. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic undifferentiated neoplasms.           Immunogen         Human B-lymphocytes (Uniprot: P08575)           Product Application Details         Flow Cytometry, Immunohistochemistry/ Immunofluorescence, Immunohistochemistry, Immunofluorescence, Immunofluorescence, Immunohistochemistry. Paraffin, CyTOF-ready           Application Notes         Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to light or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellula rogets, specific attention to the fixation and permeabilizati	Conjugate	PE/Atto594
Product Description         Host       Mouse         Gene ID       5788         Gene Symbol       PTPRC         Species       Human         Marker       B-Cell Marker         Specificity/Sensitivity       CD45R, also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms arise because of alternative splicing of exons 4, 5, and 6. The corresponding protein domains are characterized by the binding of monoclonal antibodies specific for CD45RA (exon 4), CD45RB (exon 5), CD45RC (exon 6) and CD45RO (exons 4 to 6 spliced out). The variation in these isoforms is localized to the extracellular domain of CD45R, while the intracellular domain is conserved. CD45R functions as a phosphor-tyrosine phosphatase. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic undifferentiated neoplasms.         Immunogen       Human B-lymphocytes (Uniprot: P08575)         Product Application Details       Flow Cytometry, Immunohistochemistry-Paraffin, CyTOF-ready         Recommended Dilutions       Flow Cytometry, Immunohistochemistry-Paraffin, CyTOF-ready         Application Notes       Optimal results using our Tandem dyes, please avoid prolonged exposure to light or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixation and permeabilization steps in your flow protocol may be required. Please contact our technical support team at technic	Purity	Protein A or G purified
Host         Mouse           Gene ID         5788           Gene Symbol         PTPRC           Species         Human           Marker         B-Cell Marker           Specificity/Sensitivity         CD45R, also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms arise because of alternative splicing of exons 4, 5, and 6. The corresponding protein domains are characterized by the binding of monoclonal antibodies specific for CD45R (exon 4), CD45R (exon 5), CD45RC (exon 6) and CD45R (exon 4), CD45R (exon 5), CD45RC (exon 6) and CD45R (exon 4), CD45R (exon 5), CD45R (exon 5), CD45R (con 5), CD45R (con 5), CD45R (con 5) and Spattese. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic undifferentiated neoplasms.           Immunogen         Human B-lymphocytes (Uniprot: P08575)           Product Application Details         Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunofluorescence, Immunohistochemistry, Immunohistochemistry. CyTOF-ready           Recommended Dilutions         Flow Cytometry, Immunohistochemistry. Paraffin, CyTOF-ready           Application Notes         Optimal results using our Tandem dyes, please avoid prolonged exposure to light or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling, When staining intracellular targets, specific attention to the fixation and permeabilization steps in your flow protocol may be required. Please contact our technical suport team at technical@novusbi	Buffer	PBS
Gene ID       5788         Gene Symbol       PTPRC         Species       Human         Marker       B-Cell Marker         Specificity/Sensitivity       CD45R, also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms arise because of alternative splicing of exons 4, 5, and 6. The corresponding protein domains are characterized by the binding of monoclonal antibodies specific for CD45RA (exon 4), CD45RB (exon 5), CD45RC (exon 6) and CD45RO (exons 4 to 6 spliced out). The variation in these isoforms is localized to the extracellular domain of CD45R, while the intracellular domain is conserved. CD45R functions as a phosphor-tyrosine phosphatase. Antibody to CD45 is useful in differentiated neoplasms.         Immunogen       Human B-lymphocytes (Uniprot: P08575)         Product Application Details       Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready         Recommended Dilutions       Flow Cytometry, Immunohistochemistry-Paraffin, CyTOF-ready         Application Notes       Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to light or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixation and permeabilization steps in your flow protocod may be required. Please contact our technical support team at technical@novusbio.com if you have any </th <th colspan="2">Product Description</th>	Product Description	
Gene Symbol       PTPRC         Species       Human         Marker       B-Cell Marker         Specificity/Sensitivity       CD45R, also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms arise because of alternative splicing of exons 4, 5, and 6. The corresponding protein domains are characterized by the binding of monoclonal antibodies specific for CD45RA (exon 4), CD45RB (exon 5), CD45RC (exon 6) and CD45RO (exons 4 to 6 spliced out). The variation in these isoforms is localized to the extracellular domain of CD45R, while the intracellular domain is conserved. CD45R functions as a phosphor-tyrosine phosphatase. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic undifferentiated neoplasms.         Immunogen       Human B-lymphocytes (Uniprot: P08575)         Product Application Details       Flow Cytometry, Immunocytochemistry/ Immunofluorescence, ImmunohistochemistryParaffin, CyTOF-ready         Recommended Dilutions       Flow Cytometry, Immunohistochemistry-Paraffin, CyTOF-ready         Application Notes       Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, pelase avoid prolonged exposure to light or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixation and permeabilization steps in your flow protocol may be required. Please contact our technical support team at technical@novusbio.com if you have any   <	Host	Mouse
Species       Human         Marker       B-Cell Marker         Specificity/Sensitivity       CD45R, also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms arise because of alternative splicing of exons 4, 5, and 6. The corresponding protein domains are characterized by the binding of monoclonal antibodies specific for CD45RA (exon 4), CD45RB (exon 5), CD45RC (exon 6) and CD45RO (exons 4 to 6 spliced out). The variation in these isoforms is localized to the extracellular domain of CD45R, while the intracellular domain is conserved. CD45R functions as a phosphor-tyrosine phosphatase. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic undifferentiated neoplasms.         Immunogen       Human B-lymphocytes (Uniprot: P08575)         Product Application Details       Flow Cytometry, Immunohistochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry, CyTOF-ready         Application Notes       Optimal results using our Tandem dyes, please avoid prolonged exposure to light or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixation and permeabilization steps in your flow protocol may be required. Please contact our technical support team at technical@novusbio.com if you have any	Gene ID	5788
Marker         B-Cell Marker           Specificity/Sensitivity         CD45R, also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms arise because of alternative splicing of exons 4, 5, and 6. The corresponding protein domains are characterized by the binding of monoclonal antibodies specific for CD45RA (exon 4), CD45RB (exon 5), CD45RC (exon 6) and CD45RO (exons 4 to 6 spliced out). The variation in these isoforms is localized to the extracellular domain of CD45R, while the intracellular domain is conserved. CD45R functions as a phosphor-tyrosine phosphatase. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic undifferentiated neoplasms.           Immunogen         Human B-lymphocytes (Uniprot: P08575)           Product Application Details         Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready           Recommended Dilutions         Flow Cytometry, Immunohistochemistry. Paraffin, CyTOF-ready           Application Notes         Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to light or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixation and permeabilization steps in your flow protocol may be required. Please contact our technical support team at technical@novusbio.com if you have any	Gene Symbol	PTPRC
Specificity/Sensitivity       CD45R, also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms arise because of alternative splicing of exons 4, 5, and 6. The corresponding protein domains are characterized by the binding of monoclonal antibodies specific for CD45RA (exon 4), CD45RB (exon 5), CD45RC (exon 6) and CD45RO (exons 4 to 6 spliced out). The variation in these isoforms is localized to the extracellular domain of CD45R, while the intracellular domain is conserved. CD45R functions as a phosphor-tyrosine phosphatase. Antibody to CD45s is useful in differential diagnosis of lymphoid tumors from non-hematopoietic undifferentiated neoplasms.         Immunogen       Human B-lymphocytes (Uniprot: P08575)         Product Application Details       Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready         Recommended Dilutions       Flow Cytometry, Immunohistochemistry-Paraffin, CyTOF-ready         Application Notes       Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to light or externe temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular atragets, specific attention to the fixation and permeabilization steps in your flow protocol may be required. Please contact our technical support team at technical@novusbio.com if you have any	Species	Human
transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms arise because of alternative splicing of exons 4, 5, and 6. The corresponding protein domains are characterized by the 	Marker	B-Cell Marker
Product Application DetailsApplicationsFlow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-readyRecommended DilutionsFlow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, CyTOF-readyApplication NotesOptimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to light or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixation and permeabilization steps in your flow protocol may be required. Please contact our technical support team at technical@novusbio.com if you have any	Specificity/Sensitivity	transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms arise because of alternative splicing of exons 4, 5, and 6. The corresponding protein domains are characterized by the binding of monoclonal antibodies specific for CD45RA (exon 4), CD45RB (exon 5), CD45RC (exon 6) and CD45RO (exons 4 to 6 spliced out). The variation in these isoforms is localized to the extracellular domain of CD45R, while the intracellular domain is conserved. CD45R functions as a phosphor-tyrosine phosphatase. Antibody to CD45 is useful in differential diagnosis of lymphoid
ApplicationsFlow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-readyRecommended DilutionsFlow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, CyTOF-readyApplication NotesOptimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to light or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixation and permeabilization steps in your flow protocol may be required. Please contact our technical support team at technical@novusbio.com if you have any	Immunogen	Human B-lymphocytes (Uniprot: P08575)
Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-readyRecommended DilutionsFlow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, CyTOF-readyApplication NotesOptimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to light or extreme temperature fluctuations. These can lead to irreversible degradation 	Product Application Details	
Application NotesImmunofluorescence, Immunohistochemistry-Paraffin, CyTOF-readyApplication NotesOptimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to light or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixation and permeabilization steps in your flow protocol may be required. Please contact our technical support team at technical@novusbio.com if you have any	Applications	
optimal results using our Tandem dyes, please avoid prolonged exposure to light or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixation and permeabilization steps in your flow protocol may be required. Please contact our technical support team at technical@novusbio.com if you have any	Recommended Dilutions	
	Application Notes	optimal results using our Tandem dyes, please avoid prolonged exposure to light or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixation and permeabilization steps in your flow protocol may be required. Please contact our technical support team at technical@novusbio.com if you have any





#### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

#### Products Related to NBP2-54465PEATT594

210-TA-005	TNF-alpha [Unconjugated]
M6000B-1	IL-6 [HRP]
6507-IL-010/CF	IL-4 [Unconjugated]
285-IF-100	IFN-gamma [Unconjugated]

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-54465PEATT594

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

