## **Product Datasheet**

### CD79A Antibody (IGA/1688R) [Alexa Fluor® 405] NBP2-54469AF405

Unit Size: 100 ul

Store at 4C in the dark.

www.novusbio.com

G

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-54469AF405

Updated 10/23/2024 v.20.1

# Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-54469AF405



#### NBP2-54469AF405

CD79A Antibody (IGA/1688R) [Alexa Fluor® 405]

Product InformationUnit Size100 ulConcentrationPlease see the vial label for concentration. If unisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneIGA/1688RPreservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 405PurityProtein A purifiedBuffer50mM Sodium BorateProduct Description973Gene ID973Gene SymbolCD79ASpecificity/SensitivityA disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of motion in the majority of acute leukemias of precursor Be cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is generally used to complement amit-CD20. Anti-CD79a is of acute leukemias of precursor B cell type, in B cell lines. Anti-CD79a is generally used to complement amit-CD20. Anti-CD79a is of acute leukemias of precursor B cell type. In B cell lines. Anti-CD79a is of acute ieukemias and precisits until meany of the same tymphomas anti-CD20. Anti-CD79a is of acute leukemia than is anti-CD20. Anti-CD79a is of and in some myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is of allow myelomas and in cordically for mature B-cell stape where it is found as an intracellular component. CD79a is of allow and in some myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20. Anti-CD79a is of allow and in some myelomas. It is not present in my	у. , г.	•
Concentration         Please see the vial label for concentration. If unlisted please contact technical services.           Storage         Store at 4C in the dark.           Clonality         Monoclonal           Clone         IGA/1688R           Preservative         0.05% Sodium Azide           Isotype         IgG           Conjugate         Alexa Fluor 405           Purity         Protein A purified           Buffer         50mM Sodium Borate           Product Description         Host           Gene ID         973           Gene Symbol         CD79A           Species         Human, Mouse, Rat           Marker         B-Cell Marker           Specificity/Sensitivity         A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79a) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 opypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 opypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a land B29 (or CD79a) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 opypeptides and immunoglobulins on s cells tage. Aery to mature B-cell Imphomas and is some every to mature B-cell Imphomas and is some cell stage. early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precision B-temphoblastic furphomas and i	Product Information	
services.         Storage       Store at 4C in the dark.         Clonality       Monoclonal         Clone       IGA/1688R         Preservative       0.05% Sodium Azide         Isotype       IgG         Conjugate       Alexa Fluor 405         Purity       Protein A purified         Buffer       50mM Sodium Borate         Product Description       Fooduct Description         Host       Rabbit         Gene ID       973         Gene Symbol       CD79A         Species       Human, Mouse, Rat         Marker       B-Cell Marker         Specificity/Sensitivity       A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre B cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell type, pora sifer treatment with Rituximab (anti-CD20, but also is more likely to stain B-lymphomas after treatment with Rituximab (anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20, but also is more likely to stain B-lymphoblastic lymphomas externed with Rituximab (anti-CD20, but also is more likely to stain B-lymphoblastic lightomas as anti-CD20, but also is more likely to stain B-lymphoblastic	Unit Size	100 ul
Clonality       Monoclonal         Clone       IGA/1688R         Preservative       0.05% Sodium Azide         Isotype       IgG         Conjugate       Alexa Fluor 405         Purity       Protein A purified         Buffer       50mM Sodium Borate         Product Description       Rabbit         Gene ID       973         Gene Symbol       CD79A         Species       Human, Mouse, Rat         Marker       B-Cell Marker         Specificity/Sensitivity       A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 oplypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 oplypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 oplypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 oplypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 oplypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 oplypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 oplypeptides and immunoglobulins on P cells. This complex of mb-1 and B29 oplypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 oplypeptides and immunoglobulins on P cells. This complex of mb-1 and B29 oplypeptides and immunoglobulins on P cells. This complex of mb-1 and B29 oplypeptides and immunoglobulins on P cells. This complex of the and bypeptides and immunoglobulins on the ce	Concentration	
Clone       IGA/1688R         Preservative       0.05% Sodium Azide         Isotype       IgG         Conjugate       Alexa Fluor 405         Purity       Protein A purified         Buffer       50mM Sodium Borate         Product Description       Host         Host       Rabbit         Gene ID       973         Gene Symbol       CD79A         Species       Human, Mouse, Rat         Marker       B-Cell Marker         Specificity/Sensitivity       A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 oplypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 oplypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 oplypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 oplypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 oplypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 oplypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 oplypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 oplypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 oplypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 oplypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 oplypeptides and immunoglobulins on the cell Ag receptor. CD79a is generally used to complement anti-CD20 especially for	Storage	Store at 4C in the dark.
Preservative       0.05% Sodium Azide         Isotype       IgG         Conjugate       Alexa Fluor 405         Purity       Protein A purified         Buffer       50mM Sodium Borate         Product Description       Host         Host       Rabbit         Gene ID       973         Gene Symbol       CD79A         Species       Human, Mouse, Rat         Marker       B-Cell Marker         Specificity/Sensitivity       A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 nolypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 nolypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 nolypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20. Nut also is more likely to stain B-tymphoblastic lymphomas as anti-CD20, but also is more likely to stain B-tymphoblastic lymphomas as anti-CD20, but also is more likely to stai	Clonality	Monoclonal
Isotype       IgG         Conjugate       Alexa Fluor 405         Purity       Protein A purified         Buffer       50mM Sodium Borate         Product Description       Image: Complement of the second sec	Clone	IGA/1688R
ConjugateAlexa Fluor 405PurityProtein A purifiedBuffer50mM Sodium BorateProduct DescriptionHostRabbitGene ID973Gene SymbolCD79ASpeciesHuman, Mouse, RatMarkerB-Cell MarkerSpecificity/SensitivityA disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and sea where it is found as an intracellular component. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the same lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well.ImmunogenA synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of	Preservative	0.05% Sodium Azide
Purity       Protein A purified         Buffer       50mM Sodium Borate         Product Description          Host       Rabbit         Gene ID       973         Gene Symbol       CD79A         Species       Human, Mouse, Rat         Marker       B-Cell Marker         Specificity/Sensitivity       A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins constitute the B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell lype, in B cell lines. Anti-CD79a is generally used to complement anti-CD20. but also is more likely to stain B-lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the same lymphoma/leukemia than is anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD2	Isotype	IgG
Buffer       50mM Sodium Borate         Product Description       Rabbit         Host       Rabbit         Gene ID       973         Gene Symbol       CD79A         Species       Human, Mouse, Rat         Marker       B-Cell Marker         Specificity/Sensitivity       A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines, Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the same lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphomas and end cocasionally some types of endothelial cells as well.         Immunogen       A synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of	Conjugate	Alexa Fluor 405
Product Description         Host       Rabbit         Gene ID       973         Gene Symbol       CD79A         Species       Human, Mouse, Rat         Marker       B-Cell Marker         Specificity/Sensitivity       A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines. B cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the same lymphomal suit. Clymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well.         Immunogen       A synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of	Purity	Protein A purified
HostRabbitGene ID973Gene SymbolCD79ASpeciesHuman, Mouse, RatMarkerB-Cell MarkerSpecificity/SensitivityA disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the same lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well.ImmunogenA synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of	Buffer	50mM Sodium Borate
Gene ID973Gene SymbolCD79ASpeciesHuman, Mouse, RatMarkerB-Cell MarkerSpecificity/SensitivityA disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the same lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well.ImmunogenA synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of	Product Description	
Gene SymbolCD79ASpeciesHuman, Mouse, RatMarkerB-Cell MarkerSpecificity/SensitivityA disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the same lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well.ImmunogenA synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of	Host	Rabbit
SpeciesHuman, Mouse, RatMarkerB-Cell MarkerSpecificity/SensitivityA disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the same lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well.ImmunogenA synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of	Gene ID	973
MarkerB-Cell MarkerSpecificity/SensitivityA disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the same lymphoma as anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well.ImmunogenA synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of	Gene Symbol	CD79A
Specificity/SensitivityA disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the same lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well.ImmunogenA synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of	Species	Human, Mouse, Rat
<ul> <li>CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines. B cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the same lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well.</li> <li>Immunogen</li> </ul>	Marker	B-Cell Marker
	Specificity/Sensitivity	CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the same lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of
	Immunogen	

www.novusbio.com



	5
Notes	Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.
Product Application Details	
Applications	Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready
Recommended Dilutions	Western Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, CyTOF-ready
Application Notes	Optimal dilution of this antibody should be experimentally determined.





#### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

#### Products Related to NBP2-54469AF405

NBP2-24891AF405	Rabbit IgG Isotype Control [Alexa Fluor® 405]
NBP2-60209-50ug	Recombinant Human CD79A His Protein
7268-CT-100	CTLA-4 [Unconjugated]
9685-CD-050	CD79A [Unconjugated]

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-54469AF405

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

