Product Datasheet

PAX8 Antibody (PAX8/1492) [Alexa Fluor® 700] NBP2-54539AF700

Unit Size: 100 ul

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-54539AF700

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-54539AF700



NBP2-54539AF700

PAX8 Antibody (PAX8/1492) [Alexa Fluor® 700]

Product InformationUnit Size100 ulConcentrationPlease see the vial label for concentration. If unisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalClonePAX8/1492Preservative0.05% Sodium AzideIsotypeIgG2b KappaConjugateAlexa Fluor 700PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseGene ID7849Gene SymbolPAX8Specificity/SensitivityRecognizes a protein of 62kDa, identified as PAX8. It is a member of the paired box (PAX) family of transcription factors. This nuclear protein is involved in thryroid follicular carl development and expression of thryroid-specific genes. Murations in this gene have been associated with thryroid sygenesis, thryroid follicular carcinomas, and atypical thryroid adenous. PAX-8 is expressed in the thryroid (and associated carcinomas), non-ciliated mucosal cells of the fallopian tubes, and simple ovarian inclusion cysts, but not normal ovarian sturface epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, epithel		
ConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalClonePAX8/1492Preservative0.05% Sodium AzideIsotypeIgG2b KappaConjugateAlexa Fluor 700PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description7849Gene ID7849Gene SymbolPAX8SpeciesHuman, CanineMarkerRenal Cell MarkerSpecificity/SensitivityRecognizes a protein of 62kDa, identified as PAX8. It is a member of the paired box (PAX) failing in thryoid datomasion at thryoid datomasion at thryoid (and associated carcinomas, and atypical thryoid admorans expression is reported in thryoid (and associated carcinomas, phyloplatom servers), endmany ovarian expression is reported in renal tubules as well as renal cell carcinomas, PAX-8 is expressed in the thryoid (and associated carcinomas, phyloplatom ovarian serves, endometrioid, and clear cell carcinomas, PAX-8 is expressed in the thryoid (and associated carcinomas, phyloplatom ovarian serves, endometrioid, and clear cell carcinomas, PAX-8 is expressed in the thryoid (and essociated carcinomas, phyloplatom, PAX-8 antibody may be used as an additional immunohistochemical marker for renal epithelial tubes, and simple ovarian inclusion cysts, but no tornal ovarian serves, endometrioid, and clear cell carcinomas, and seminoma. PAX-8 antibody may be used as an additional immunohistochemical marker for renal epithelial tumors.ImmunogenRecombinant fragment (around aa 60-261) of human PAX8 protein (exact	Product Information	
services. Storage Store at 4C in the dark. Clonality Monoclonal Clone PAX8/1492 Preservative 0.05% Sodium Azide Isotype IgG2b Kappa Conjugate Alexa Fluor 700 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Mouse Gene ID 7849 Gene Symbol PAX8 Species Human, Canine Marker Renal Cell Marker Specificity/Sensitivity Recognizes a protein of 62kDa, identified as PAX8. It is a member of the paired box (PAX) family of transcription factors. This nuclear protein is involved in thyroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular cell development and expression is ordinan servas, endometricid, and clear cell carcinomas, NAX-8 expression is reported in the thyroid (and associated carcinomas), non-ciliated mucosal cells of the fallopian tubes, and simple ovarian inclusion cysts, but not normal ovarian surface epithelial cells. PAX-8 is expressed in a high percentage of ovarian servas, endometricid, and clear cell carcinomas, PAX-8 expression is reported in renal tubules as well as renal cell carcinoma, nephrobalstoma, and serinoma, PAX-8	Unit Size	100 ul
ClonalClonalClonePAX8/1492Preservative0.05% Sodium AzideIsotypeIgG2b KappaConjugateAlexa Fluor 700PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseHostMouseGene ID7849Gene SymbolPAX8SpeciesHuman, CanineMarkerRenal Cell MarkerSpecificity/SensitivityRecognizes a protein of 62kDa, identified as PAX8. It is a member of the paired box (PAX) family of transcription factors. This nuclear protein is involved in thyroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular carcinomas, and atypical thyroid adenomas. PAX-8 is expressed in the thyroid follicular carcinomas, but on ormal ovarian surfuces epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, endometrioid, and clear cell carcinomas, but on remal ovarian serous, endometrioid, and clear cell carcinomas, hut normal ovarian serous, endometrioid, and clear cell carcinomas, and seminoma. PAX-8 antibody may be used as an additional immunchistochemical marker for renal epithelial tumors.ImmunogenRecombinant fragment (around aa 60-261) of human PAX8 protein (exact	Concentration	
ClonePAX8/1492Preservative0.05% Sodium AzideIsotypeIgG2b KappaConjugateAlexa Fluor 700PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseGene ID7849Gene SymbolPAX8SpeciesHuman, CanineMarkerRecognizes a protein of 62kDa, identified as PAX8. It is a member of the paired box (PAX) family of transcription factors. This nuclear protein is involved in thyroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular carcinomas, and atypical thyroid adenomas. PAX-8 is expressed in the tryoid (and associated carcinomas), but on romal ovarian surface epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, endometrioid, and clear cell carcinomas, but only rarely in primary ovarian mucinous adenocarcinomas. PAX-8 expression is reported in renal tubules as well as renal cell carcinoma, nephroblastoma, and seminoma. PAX-8 antibody may be used as an additional immunohistochemical marker for renal epithelial tumors.	Storage	Store at 4C in the dark.
Preservative0.05% Sodium AzideIsotypeIgG2b KappaConjugateAlexa Fluor 700PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseGene ID7849Gene SymbolPAX8SpeciesHuman, CanineMarkerRenal Cell MarkerSpecificity/SensitivityRecognizes a protein of 62kDa, identified as PAX8. It is a member of the paired box (PAX) family of transcription factors. This nuclear protein is involved in thyroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular cell development and expression of the fallopian tubes, and simple ovarian inclusion cysts, but not normal ovarian surface epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, endometrioid, and clear cell carcinomas, hut only rarely in primary ovarian mucinous adenocarinomas. PAX-8 expression is reported in renal tubules as well as renal cell carcinoma, nephroblastoma, and seminoma. PAX-8 antibody may be used as an additional immunohistochemical marker for renal epithelial tumors.ImmunogenRecombinant fragment (around aa 60-261) of human PAX8 protein (exact	Clonality	Monoclonal
IsotypeIgG2b KappaConjugateAlexa Fluor 700PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseHostMouseGene ID7849Gene SymbolPAX8SpeciesHuman, CanineMarkerRenal Cell MarkerSpecificity/SensitivityRecognizes a protein of 62kDa, identified as PAX8. It is a member of the paired box (PAX) family of transcription factors. This nuclear protein is involved in thyroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular cell carcinomas, and atypical thyroid adenomas. PAX-8 is expressed in the thyroid (and associated carcinomas), non-ciliated mucosal cells of the fallopian tubes, and simple ovarian inclusion cysts, but not normal ovarian surface epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, endometrioid, and clear cell carcinomas, PAX-8 and seminoma. PAX-8 antibody may be used as an additional immunohistochemical marker for renal epithelial tumors.ImmunogenRecombinant fragment (around aa 60-261) of human PAX8 protein (exact	Clone	PAX8/1492
ConjugateAlexa Fluor 700PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseGene ID7849Gene SymbolPAX8SpeciesHuman, CanineMarkerRenal Cell MarkerSpecificity/SensitivityRecognizes a protein of 62kDa, identified as PAX8. It is a member of the paired box (PAX) family of transcription factors. This nuclear protein is involved in thyroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular carcinomas, and atypical thyroid adenomas. PAX-8 is expressed in the thyroid (and associated carcinomas), non-ciliated mucosal cells of the fallopian tubes, and simple ovarian inclusion cysts, but not normal ovarian surface epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, endometrioid, and clear cell carcinomas, and seminoma. PAX-8 antibody may be used as an additional immunohistochemical marker for renal epithelial tumors.ImmunogenRecombinant fragment (around aa 60-261) of human PAX8 protein (exact	Preservative	0.05% Sodium Azide
PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID7849Gene SymbolPAX8SpeciesHuman, CanineMarkerRenal Cell MarkerSpecificity/SensitivityRecognizes a protein of 62kDa, identified as PAX8. It is a member of the paired box (PAX) family of transcription factors. This nuclear protein is involved in thyroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid specific genes. Mutations in this gene have been associated with thyroid specific genes. Mutations in this gene have been associated with thyroid specific genes. Mutations in the gene have been associated with thyroid specific genes. Mutations in the gene have been associated with thyroid specific genes. Mutations in the gene have been associated with thyroid specific genes. Mutations in the gene have been associated with thyroid specific genes. Mutations in the gene have been associated with thyroid specific genes. Mutations in the gene have been associated with thyroid specific genes. Mutations in the gene have been associated with thyroid specific genes. Mutations in the gene have been associated with thyroid adenomas. PAX-8 is expressed in the thyroid (and associated carcinomas), non-ciliated mucosal cells of the fallopian tubes, and simple ovarian inclusion cysts, but not normal ovarian surface epithelial cells. PAX-8 expression is reported in renal tubules as well as renal cell carcinoma, nephroblastoma, and seminoma. PAX-8 antibody may be used as an additional immunohistochemical marker for renal epithelial tumors.ImmunogenRecombinant fragment (around aa 60-261) of human PAX8 protein (exact	Isotype	IgG2b Kappa
Buffer50mM Sodium BorateProduct DescriptionHostMouseGene ID7849Gene SymbolPAX8SpeciesHuman, CanineMarkerRenal Cell MarkerSpecificity/SensitivityRecognizes a protein of 62kDa, identified as PAX8. It is a member of the paired box (PAX) family of transcription factors. This nuclear protein is involved in thyroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular carcinomas, and atypical thyroid adenomas. PAX-8 is expressed in the thyroid (and associated carcinomas), non-ciliated mucosal cells of the fallopian tubes, and simple ovarian in clusion cysts, but not normal ovarian surface epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, endometrioid, and clear cell carcinomas, and seminoma. PAX-8 antibody may be used as an additional immunohistochemical marker for renal epithelial ImmunogenImmunogenRecombinant fragment (around aa 60-261) of human PAX8 protein (exact	Conjugate	Alexa Fluor 700
Product DescriptionHostMouseGene ID7849Gene SymbolPAX8SpeciesHuman, CanineMarkerRenal Cell MarkerSpecificity/SensitivityRecognizes a protein of 62kDa, identified as PAX8. It is a member of the paired box (PAX) family of transcription factors. This nuclear protein is involved in thyroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular carcinomas, and atypical thyroid adenomas. PAX-8 is expressed in the thyroid follicular cell carcinomas, non-ciliated mucosal cells of the fallopian tubes, and simple ovarian inclusion cysts, but not normal ovarian surface epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, endometrioid, and clear cell carcinomas, but only rarely in primary ovarian mucinous adenocarcinomas. PAX-8 expression is reported in renal tubules as well as renal cell carcinoma, nephroblastoma, and seminoma. PAX-8 antibody may be used as an additional immunohistochemical marker for renal epithelial tumors.ImmunogenRecombinant fragment (around aa 60-261) of human PAX8 protein (exact	Purity	Protein A or G purified
HostMouseGene ID7849Gene SymbolPAX8SpeciesHuman, CanineMarkerRenal Cell MarkerSpecificity/SensitivityRecognizes a protein of 62kDa, identified as PAX8. It is a member of the paired box (PAX) family of transcription factors. This nuclear protein is involved in thyroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular carcinomas, and atypical thyroid adenomas. PAX-8 is expressed in the thyroid (and associated carcinomas), non-ciliated mucosal cells of the fallopian tubes, and simple ovarian inclusion cysts, but not normal ovarian surface epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, endometrioid, and clear cell carcinomas, but only rarely in primary ovarian mucinous adenocarcinomas. PAX-8 expression is reported in renal tubules as well as renal cell carcinoma, nephroblastoma, and seminoma. PAX-8 antibody may be used as an additional immunohistochemical marker for renal epithelial tumors.ImmunogenRecombinant fragment (around aa 60-261) of human PAX8 protein (exact	Buffer	50mM Sodium Borate
Gene ID7849Gene SymbolPAX8SpeciesHuman, CanineMarkerRenal Cell MarkerSpecificity/SensitivityRecognizes a protein of 62kDa, identified as PAX8. It is a member of the paired box (PAX) family of transcription factors. This nuclear protein is involved in thyroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular carcinomas, and atypical thyroid adenomas. PAX-8 is expressed in the thyroid (and associated carcinomas), non-ciliated mucosal cells of the fallopian tubes, and simple ovarian inclusion cysts, but not normal ovarian surface epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, endometrioid, and clear cell carcinomas, but only rarely in primary ovarian mucinous adenocarcinomas. PAX-8 expression is reported in renal tubules as well as renal cell carcinoma, nephroblastoma, and seminoma. PAX-8 antibody may be used as an additional immunohistochemical marker for renal epithelial tumors.ImmunogenRecombinant fragment (around aa 60-261) of human PAX8 protein (exact	Product Description	
Gene SymbolPAX8SpeciesHuman, CanineMarkerRenal Cell MarkerSpecificity/SensitivityRecognizes a protein of 62kDa, identified as PAX8. It is a member of the paired box (PAX) family of transcription factors. This nuclear protein is involved in thyroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular carcinomas, and atypical thyroid adenomas. PAX-8 is expressed in the thyroid (and associated carcinomas), non-ciliated mucosal cells of the fallopian tubes, and simple ovarian inclusion cysts, but not normal ovarian surface epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, endometrioid, and clear cell carcinomas, but only rarely in primary ovarian mucinous adenocarcinoma. PAX-8 expression is reported in renal tubules as well as renal cell carcinoma, nephroblastoma, and seminoma. PAX-8 antibody may be used as an additional immunohistochemical marker for renal epithelial tumors.ImmunogenRecombinant fragment (around aa 60-261) of human PAX8 protein (exact	Host	Mouse
SpeciesHuman, CanineMarkerRenal Cell MarkerSpecificity/SensitivityRecognizes a protein of 62kDa, identified as PAX8. It is a member of the paired box (PAX) family of transcription factors. This nuclear protein is involved in thyroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular carcinomas, and atypical thyroid adenomas. PAX-8 is expressed in the thyroid (and associated carcinomas), non-ciliated mucosal cells of the fallopian tubes, and simple ovarian inclusion cysts, but not normal ovarian surface epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, endometrioid, and clear cell carcinomas, but only rarely in primary ovarian mucinous adenocarcinomas. PAX-8 expression is reported in renal tubules as well as renal cell carcinoma, nephroblastoma, and seminoma. PAX-8 antibody may be used as an additional immunohistochemical marker for renal epithelial tumors.ImmunogenRecombinant fragment (around aa 60-261) of human PAX8 protein (exact	Gene ID	7849
MarkerRenal Cell MarkerSpecificity/SensitivityRecognizes a protein of 62kDa, identified as PAX8. It is a member of the paired box (PAX) family of transcription factors. This nuclear protein is involved in thyroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular carcinomas, and atypical thyroid adenomas. PAX-8 is expressed in the thyroid (and associated carcinomas), non-ciliated mucosal cells of the fallopian tubes, and simple ovarian inclusion cysts, but not normal ovarian surface epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, endometrioid, and clear cell carcinomas, but only rarely in primary ovarian mucinous adenocarcinomas. PAX-8 expression is reported in renal tubules as well as renal cell carcinoma, nephroblastoma, and seminoma. PAX-8 antibody may be used as an additional immunohistochemical marker for renal epithelial tumors.ImmunogenRecombinant fragment (around aa 60-261) of human PAX8 protein (exact	Gene Symbol	PAX8
Specificity/SensitivityRecognizes a protein of 62kDa, identified as PAX8. It is a member of the paired box (PAX) family of transcription factors. This nuclear protein is involved in thyroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular carcinomas, and atypical thyroid adenomas. PAX-8 is expressed in the thyroid (and associated carcinomas), non-ciliated mucosal cells of the fallopian tubes, and simple ovarian inclusion cysts, but not normal ovarian surface epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, endometrioid, and clear cell carcinomas, but only rarely in primary ovarian mucinous adenocarcinomas. PAX-8 expression is reported in renal tubules as well as renal cell carcinoma, nephroblastoma, and seminoma. PAX-8 antibody may be used as an additional immunohistochemical marker for renal epithelial tumors.ImmunogenRecombinant fragment (around aa 60-261) of human PAX8 protein (exact	Species	Human, Canine
 box (PAX) family of transcription factors. This nuclear protein is involved in thyroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular carcinomas, and atypical thyroid adenomas. PAX-8 is expressed in the thyroid (and associated carcinomas), non-ciliated mucosal cells of the fallopian tubes, and simple ovarian inclusion cysts, but not normal ovarian surface epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, endometrioid, and clear cell carcinomas, but only rarely in primary ovarian mucinous adenocarcinomas. PAX-8 expression is reported in renal tubules as well as renal cell carcinoma, nephroblastoma, and seminoma. PAX-8 antibody may be used as an additional immunohistochemical marker for renal epithelial tumors. Immunogen 	Marker	Renal Cell Marker
	Specificity/Sensitivity	box (PAX) family of transcription factors. This nuclear protein is involved in thyroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular carcinomas, and atypical thyroid adenomas. PAX-8 is expressed in the thyroid (and associated carcinomas), non-ciliated mucosal cells of the fallopian tubes, and simple ovarian inclusion cysts, but not normal ovarian surface epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, endometrioid, and clear cell carcinomas, but only rarely in primary ovarian mucinous adenocarcinomas. PAX-8 expression is reported in renal tubules as well as renal cell carcinoma, nephroblastoma, and seminoma. PAX-8 antibody may be used as an additional immunohistochemical marker for renal epithelial
	Immunogen	

www.novusbio.com



Notes	Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.
Product Application Details	
Applications	Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Protein Array, CyTOF- ready
Recommended Dilutions	Western Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, Protein Array, CyTOF- ready
Application Notes	Optimal dilution of this antibody should be experimentally determined.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-54539AF700

NBP1-43317AF700Mouse IgG2b Kappa Light Chain Isotype Control (MG2b) [Alexa Fluor®
700]NBP2-51907-0.05mgRecombinant Human PAX8 His Protein288-TPN-025/CFThrombopoietin/THPO [Unconjugated]NBL1-14132PAX8 Overexpression Lysate

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-54539AF700

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

