

Product Datasheet

HDAC1 Antibody - BSA Free

NBP2-54621

Unit Size: 50 ug

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-54621

Updated 9/9/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-54621



NBP2-54621

HDAC1 Antibody - BSA Free

Product Information

Unit Size	50 ug
Concentration	Please see the vial label for concentration. If unlisted please contact technical services.
Storage	Store at -20C. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.05% Sodium Azide and 0.05% ProClin 300
Isotype	IgG
Purity	Affinity purified
Buffer	PBS

Product Description


Description	Novus Biologicals Rabbit HDAC1 Antibody - BSA Free (NBP2-54621) is a polyclonal antibody validated for use in WB, ELISA, ICC/IF and ChIP. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	3065
Gene Symbol	HDAC1
Species	Human, Mouse
Immunogen	HDAC1

Product Application Details


Applications	Western Blot, ELISA, Immunocytochemistry/ Immunofluorescence, Protein Array, Chromatin Immunoprecipitation (ChIP), Chromatin Immunoprecipitation Sequencing, Knockdown Validated
Recommended Dilutions	Western Blot 1:1000, ELISA 1:4000, Immunocytochemistry/ Immunofluorescence 1:500, Protein Array 1:100000, Chromatin Immunoprecipitation (ChIP) 2 ug/IP, Chromatin Immunoprecipitation Sequencing, Knockdown Validated

Images


Western Blot: HDAC1 Antibody [NBP2-54621] - Whole cell extracts (50 ug) from HeLa cells transfected with HDAC1 siRNA (lane 2) and from an untransfected control (lane 1) were analysed by Western blot using the antibody against HDAC1 diluted 1:1,000 in TBS-Tween containing 5% skimmed milk. The position of the protein of interest is indicated on the right (expected size: 55 kDa); the marker (in kDa) is shown on the left.




Immunocytochemistry/Immunofluorescence: HDAC1 Antibody [NBP2-54621] - HeLa cells were stained with the antibody against HDAC1 and with DAPI. Cells were fixed with 4% formaldehyde for 10' and blocked with PBS/TX-100 containing 5% normal goat serum and 1% BSA. The cells were immunofluorescently labelled with the HDAC1 antibody (left) diluted 1:500 in blocking solution followed by an anti-rabbit antibody conjugated to Alexa488. The middle panel shows staining of the nuclei with DAPI. A merge of the two stainings is shown on the right.




Chromatin Immunoprecipitation: HDAC1 Antibody [NBP2-54621] - ChIP was performed with the antibody against HDAC1 on sheared chromatin from 4,000,000 HeLa cells. An antibody titration consisting of 1, 2, 5 and 10 µg per ChIP experiment was analysed. IgG (2 µg/IP) was used as negative IP control. QPCR was performed with primers specific for the EIF4A2 and GAPDH promoters, used as positive controls, and for the MYOD1 gene and Sat2 satellite repeat, used as negative controls. Figure shows the recovery, expressed as a % of input (the relative amount of immunoprecipitated DNA compared to input DNA after qPCR analysis).




Western Blot: HDAC1 Antibody [NBP2-54621] - Whole cell extracts (25 µg, lane 1) and nuclear extracts (25 µg, lane 2) from HeLa cells were analysed by Western blot using the antibody against HDAC1 diluted 1:1,000 in TBS-Tween containing 5% skimmed milk. The position of the protein of interest is indicated on the right (expected size: 55 kDa); the marker (in kDa) is shown on the left.



ELISA: HDAC1 Antibody [NBP2-54621] - To determine the titer of the antibody, an ELISA was performed using a serial dilution of the antibody directed against HDAC1, crude serum and flow through. The plates were coated with the peptide used for immunization of the rabbit. By plotting the absorbance against the antibody dilution, the titer of the antibody was estimated to be 1:75,000.



Protein Array: HDAC1 Antibody [NBP2-54621] - The specificity of the antibody against HDAC1 was demonstrated using a human protein microarray containing more than 19,000 human proteins. The antibody was used at a dilution of 1:100,000. The figure shows the Z-score of the signal intensity (mean value of the duplicate spots on the array). The names of the proteins with 5 highest Z-scores are indicated at the bottom. This figure clearly shows the high specificity of the antibody for HDAC1.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-54621

HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control
H00003065-P01-10ug	Recombinant Human HDAC1 GST (N-Term) Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-54621

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

