## **Product Datasheet**

### Bcl-6 Antibody (BCL6/1475) [Alexa Fluor® 700] NBP2-59597AF700

Unit Size: 100 ul

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-59597AF700

Updated 10/23/2024 v.20.1

# Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-59597AF700



#### NBP2-59597AF700

Bcl-6 Antibody (BCL6/1475) [Alexa Fluor® 700]

Product InformationUnit Size100 ulConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneBCL6/1475Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 700PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description604Gene ID604Gene SymbolBCL6SpeciesHumanMarkerFollicular Lymphoma MarkerSpecificity/SensitivityThe specificity of this monoclonal antibody to its intended target was validated by HuProTM Array, containing more than 19.000, full-length human proteins. Recognizes a protein of 95kDa, which is identified as Bcl-6. Antibody to bcl-6 is helpful in a number of diagnostic settings: (1) In the differential diagnosis of small B-cell lymphoma. Follicular lymphoma will show bcl-6 (and CD10) positivity whereas other small B-cell lymphomas (LBCL), where CD10, bcl-6 and MUM1/IRF4 are used to identify germinal center and activated B-cell phenotypes. (3) Bcl-6 can be valuable in distinguisting classical Hodgkin lymphoma (NLEPHL). The Reed-Stermberg cells of classical Hodgkin lymphomaImmunogenRecominant human bcl-6 protein fragment (faround aa256-389) (exact sequence is proprietary) (longrot: P41162)		-			
ConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneBCL6/1475Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 700PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description604Gene ID604Gene SymbolBCL6SpeciesHumanMarkerFolicular Lymphoma MarkerSpecificity/SensitivityThe specificity of this monoclonal antibody to its intended target was validated by HuProtTM Array, containing more than 19.000, full-length human proteins. Recognizes a protein of 95kDa, which is identified as Bcl-6. Antibody to bcl-6 is helpful in a number of diagnostic settings: (1) In the differential diagnosis of small B-cell lymphoma. Folicular lymphoma will show bcl-6 is helpful in a number of diagnostic settings: (1) In the differential diagnosis of small B-cell phenotypes. (3) Bcl-6 can be valuable in distinguishing classical Hodgkin lymphoma from nodular lymphoma are usually negative. (2) Bcl-6 is an important progenosic marker in diffuse lacel lymphomas (DLBCL), where Colo, bcl-6 and SUL0 / Destivity whereas other small B-cell lymphomas are usually negative. (2) Bcl-6 is an important progenosic marker in diffuse lacel-Bockin lymphoma (NLBPHL). The Reed-Sternberg cells of classical Hodgkin lymphoma are bcl-6 negative whereas the large (L+1) cells of NLPHL are bcl-6 positive. In contrast, anti-Bcl-6 rarely stains mantle-cell lymphoma and MALT lymphoma.ImmunogenRecombinant human bcl-6 protein fragment (around aa256-389) (exac	Product Information				
services.   Storage Store at 4C in the dark.   Clonality Monoclonal   Clone BCL6/1475   Preservative 0.05% Sodium Azide   Isotype IgG1 Kappa   Conjugate Alexa Fluor 700   Purity Protein A or G purified   Buffer SomM Sodium Borate   Product Description Mouse   Gene ID 604   Gene Symbol BCL6   Species Human   Marker Follicular Lymphoma Marker   Specificity/Sensitivity The specificity of this monoclonal antibody to its intended target was validated by HuProtTM Array, containing more than 19,000, full-length human proteins. Recognizes a protein of 95kDa, which is identified as Bcl-6. Antibody to bcl-6 is helpful in a number of diagnostic settings: (1) In the differential diagnosis of small B-cell lymphoma. Folicular lymphomas are usually negative. (2) Bcl-6 is an important prognostic marker in diffuse large B-cell lymphomas (DLBCL), where CD10, bcl-6 and MUM/1/RF4 are used to identify gerinial center and activated B-cell phenotypes. (3) Bcl-6 can be valuable in distinguishing classical Hodgkin lymphoma from nodular lymphocyte predominant Hodgkin lymphoma (NLPHL). The Reed-Stemberg cells of loen of yead which is identify leasiel Hodgkin lymphoma (NLPHL). The Reed-Stemberg cells of and HuM1/1/RF4 are bcl-6 positive. In contrast, anti-Bcl-6 rarely stams mantle-cell lymphoma and MALT lymphoma.	Unit Size	100 ul			
ClonalityMonoclonalCloneBCL6/1475Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 700PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHouseHostMouseGene ID604Gene SymbolBCL6SpeciesHumanMarkerFolicular Lymphoma MarkerSpecificity/SensitivityThe specificity of this monoclonal antibody to its intended target was validated by HuProtTM Array, containing more than 19,000, full-length human proteins. Recognizes a protein of 95kDa, which is identified as Bcl-6. Antibody to bcl-6 is helpful in a number of diagnostic settings: (1) In the differential diagnosis of small B-cell lymphoma. Follicular Imphoma will show bcl-6 (and CD10) positivity whereas other small B-cell lymphoma are usually negative. (2) Bcl-6 is an important prognostic marker in diffuse large B-cell lymphoma (NLPHL). The Reed-Sternberg cells of classical Hodgkin lymphoma are bcl-6 negative whereas the large ('LH') cells of NLPHL are bcl-6 positive. In contrast, anti-Bcl-6 rarely stains mantle-cell lymphoma and MALT lymphoma.ImmunogenRecombinant human bcl-6 protein fragment (around aa256-389) (exact	Concentration				
CloneBCL6/1475Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 700PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description604HostMouseGene ID604Gene SymbolBCL6SpeciesHumanMarkerFollicular Lymphoma MarkerSpecificity/SensitivityThe specificity of this monoclonal antibody to its intended target was validated by HuProtTM Array, containing more than 19.000, fill-length human proteins. Recognizes a protein of 95kDa, which is identifier al Bd-6 is an important prognoma. Follicular lymphoma will show bcl-6 is an important prognoma for modular lymphomas are usually negative. (2) Bcl-6 is an important prognoma for modular lymphomas (DLBCL), where CD10, bcl-6 and MUM1/IRF4 are used to identify germinal center and activated B-cell lymphoma from nodular lymphomyhocyte predominant Hodgkin lymphoma (NLPHL). The Reed-Sternberg cells of classical Hodgkin lymphoma. ImmunogenImmunogenRecombinant human bcl-6 protein fragment (around aa256-389) (exact	Storage	Store at 4C in the dark.			
Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 700PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description604Gene ID604Gene SymbolBCL6SpeciesHumanMarkerFollicular Lymphoma MarkerSpecificity/SensitivityThe specificity of this monoclonal antibody to its intended target was validated by HuProtTM Array, containing more than 19,000, full-length human proteins. Recognizes a protein of 95KDa, which is identified as Bcl-6. Antibody to bcl-6 is helpful in a number of diagnostic settings: (1) In the differential diagnosis of small B-cell lymphoma. Follicular lymphoma sare usually negative. (2) Bcl-6 is an important prognostic marker in diffuse large B-cell lymphomas (DLBCL), where CD10, bcl-6 and MUM1/IRF4 are used to identify germinal center and activated B-cell phenotypes. (3) Bcl-6 can be valuable in distinguishing classical Hodgkin lymphoma are bcl-6 negative whereas the large ('LH') cells of NLPHL are bcl-6 negative whereas the large ('LH') cells of NLPHL are bcl-6 negative whereas the large ('LH') cells of NLPHL are bcl-6 negative whereas the large ('LH') cells of NLPHL are bcl-6 negative whereas the large ('LH') cells of NLPHL are bcl-6 negative whereas the large ('LH') cells of NLPHL are bcl-6 negative whereas the large ('LH') cells of NLPHL are bcl-6 negative whereas the large ('LH') cells of NLPHL are bcl-6 negative whereas the large ('LH') cells of NLPHL are bcl-6 negative whereas the large ('LH') cells of NLPHL are bcl-6 negative whereas the large ('LH') cells of NLPHL are bcl-6 negative whereas the large ('LH') cells of NLPHL are bcl-6 negative whereas the large ('LH') cells of NLPHL are bcl-6 negative whereas the large ('LH') cells of NLPHL are bcl-6 negative whereas the large	Clonality	Monoclonal			
IsotypeIgG1 KappaConjugateAlexa Fluor 700PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID604Gene SymbolBCL6SpeciesHumanMarkerFollicular Lymphoma MarkerSpecificity/SensitivityThe specificity of this monoclonal antibody to its intended target was validated by HuProtTM Array, containing more than 19,000, full-length human proteins. Recognizes a protein of 95KDa, which is identified as Bcl-6. Antibody to bcl-6 is helpful in a number of diagnostic settings: (1) In the differential diagnosis of small B-cell lymphoma. Follicular lymphoma will show bcl-6 (and CD10) positivity whereas other small B-cell lymphomas are usually negative. (2) Bcl-6 is an important prognostic marker in diffuse large B-cell lymphomas (DLBCL), where CD10, bcl-6 and MUM1/IRF4 are used to identify germinal center and activated B-cell phenotypes. (3) Bcl-6 can be valuable in distinguishing classical Hodgkin lymphoma from nodular lymphonzy and bock16 positive. In contrast, anti-Bcl-6 arrely stains mantle-cell lymphoma and MALT lymphoma.ImmunogenRecombinant human bcl-6 protein fragment (around aa256-389) (exact	Clone	BCL6/1475			
ConjugateAlexa Fluor 700PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID604Gene SymbolBCL6SpeciesHumanMarkerFollicular Lymphoma MarkerSpecificity/SensitivityThe specificity of this monoclonal antibody to its intended target was validated by HuProtTM Array, containing more than 19,000, full-length human proteins. Recognizes a protein of 95KDa, which is identified as Bcl-6. Antibody to bcl-6 is helpful in a number of diagnostic settings: (1) In the differential diagnosis of small B-cell lymphoma. Follicular lymphoma will show bcl-6 (and CD10) positivity whereas other small B-cell lymphomas are usually negative. (2) Bcl-6 is an important prognostic marker in diffuse large B-cell lymphomas (DLBCL), where CD10, bcl-6 and MUM1/IRF4 are used to identify germinal center and activated B-cell phenotypes. (3) Bcl-6 can be valuable in distinguishing classical Hodgkin lymphoma from nodular lymphocy predominal Hodgkin lymphoma are bcl-6 negative whereas the large ('LH') cells of NLPHL are bcl-6 positive. In contrast, anti-Bcl-6 rarely stains mantle-cell lymphoma and MALT lymphoma.ImmunogenRecombinant human bcl-6 protein fragment (around aa256-389) (exact	Preservative	0.05% Sodium Azide			
PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID604Gene SymbolBCL6SpeciesHumanMarkerFollicular Lymphoma MarkerSpecificity/SensitivityThe specificity of this monoclonal antibody to its intended target was validated by HuProtTM Array, containing more than 19,000, full-length human proteins. Recognizes a protein of 95KDa, which is identified as Bcl-6. Antibody to bcl-6 is helpful in a number of diagnostic settings: (1) In the differential diagnosis of small B-cell lymphoma. Follicular lymphoma are usually negative. (2) Bcl-6 is an important prognostic marker in diffuse large B-cell lymphoma (NLPHL). The Reed-Sternberg cells of classical Hodgkin lymphoma (NLPHL). The Reed-Sternberg cells of NLPHL are bcl-6 positive. In contrast, anti-Bcl-6 rarely stains mantle-cell lymphoma and MALT lymphoma.ImmunogenRecombinant human bcl-6 protein fragment (around aa256-389) (exact	Isotype	IgG1 Kappa			
Buffer 50mM Sodium Borate   Product Description Host   Host Mouse   Gene ID 604   Gene Symbol BCL6   Species Human   Marker Follicular Lymphoma Marker   Specificity/Sensitivity The specificity of this monoclonal antibody to its intended target was validated by HuProtTM Array, containing more than 19,000, full-length human proteins. Recognizes a protein of 95kDa, which is identified as Bcl-6. Antibody to bcl-6 is helpful in a number of diagnostic settings: (1) In the differential diagnosis of small B-cell lymphoma. Follicular lymphoma will show bcl-6 (and CD10) positivity whereas other small B-cell lymphoma are usually negative. (2) Bcl-6 is an important prognostic marker in diffuse large B-cell lymphomas (DLBCL), where CD10, bcl-6 and MUM1/IRF4 are used to identify germinal center and activated B-cell phenotypes. (3) Bcl-6 can be valuable in distinguishing classical Hodgkin lymphoma from nodular lymphocyte predominant Hodgkin lymphoma (NLPHL). The Reed-Stemberg cells of Classical Hodgkin lymphoma. In contrast, anti-Bcl-6 rarely stains mantle-cell lymphoma and MALT lymphoma.   Immunogen Recombinant human bcl-6 protein fragment (around aa256-389) (exact	Conjugate	Alexa Fluor 700			
Product Description   Host Mouse   Gene ID 604   Gene Symbol BCL6   Species Human   Marker Follicular Lymphoma Marker   Specificity/Sensitivity The specificity of this monoclonal antibody to its intended target was validated by HuProtTM Array, containing more than 19,000, full-length human proteins. Recognizes a protein of 95kDa, which is identified as Bcl-6. Antibody to bcl-6 is helpful in a number of diagnostic settings: (1) In the differential diagnosis of small B-cell lymphoma. Follicular lymphoma are usually negative. (2) Bcl-6 is an important prognostic marker in diffuse large B-cell lymphomas (DLBCL), where CD10, bcl-6 and MUM1/IRF4 are used to identify germinal center and activated B-cell phenotypes. (3) Bcl-6 can be valuable in distinguishing classical Hodgkin lymphoma from nodular lymphocyte predominant Hodgkin lymphoma are bcl-6 negative whereas the large ('LH') cells of NLPHL are bcl-6 positive. In contrast, anti-Bcl-6 rarely stains mantle-cell lymphoma and MALT lymphoma.   Immunogen Recombinant human bcl-6 protein fragment (around aa256-389) (exact	Purity	Protein A or G purified			
HostMouseGene ID604Gene SymbolBCL6SpeciesHumanMarkerFollicular Lymphoma MarkerSpecificity/SensitivityThe specificity of this monoclonal antibody to its intended target was validated by HuProtTM Array, containing more than 19,000, full-length human proteins. Recognizes a protein of 95kDa, which is identified as Bcl-6. Antibody to bcl-6 is helpful in a number of diagnostic settings: (1) In the differential diagnosis of small B-cell lymphoma. Follicular lymphomas are usually negative. (2) Bcl-6 is an important prognostic marker in diffuse large B-cell lymphomas (DLBCL), where CD10, bcl-6 and MUM1/IRF4 are used to identify germinal center and activated B-cell phenotypes. (3) Bcl-6 can be valuable in distinguishing classical Hodgkin lymphoma from nodular lymphocyte predominant Hodgkin lymphoma (NLPHL). The Reed-Sternberg cells of classical Hodgkin lymphoma are bcl-6 negative whereas the large ('LH') cells of NLPHL are bcl-6 positive. In contrast, anti-Bcl-6 rarely stains mantle-cell lymphoma and MALT lymphoma.ImmunogenRecombinant human bcl-6 protein fragment (around aa256-389) (exact	Buffer	50mM Sodium Borate			
Gene ID604Gene SymbolBCL6SpeciesHumanMarkerFollicular Lymphoma MarkerSpecificity/SensitivityThe specificity of this monoclonal antibody to its intended target was validated by HuProtTM Array, containing more than 19,000, full-length human proteins. Recognizes a protein of 95kDa, which is identified as Bcl-6. Antibody to bcl-6 is helpful in a number of diagnostic settings: (1) In the differential diagnosis of small B-cell lymphoma. Follicular lymphoma are usually negative. (2) Bcl-6 is an important prognostic marker in diffuse large B-cell Iymphomas (DLBCL), where CD10, bcl-6 and MUM1/IRF4 are used to identify germinal center and activated B-cell phenotypes. (3) Bcl-6 can be valuable in distinguishing classical Hodgkin lymphoma from nodular lymphocyte predominant Hodgkin lymphoma (NLPHL). The Reed-Sternberg cells of classical Hodgkin lymphoma. ImmunogenImmunogenRecombinant human bcl-6 protein fragment (around aa256-389) (exact	Product Description				
Gene SymbolBCL6SpeciesHumanMarkerFollicular Lymphoma MarkerSpecificity/SensitivityThe specificity of this monoclonal antibody to its intended target was validated by HuProtTM Array, containing more than 19,000, full-length human proteins. Recognizes a protein of 95kDa, which is identified as Bcl-6. Antibody to bcl-6 is helpful in a number of diagnostic settings: (1) In the differential diagnosis of small B-cell lymphoma. Follicular lymphoma will show bcl-6 (and CD10) positivity whereas other small B-cell lymphomas are usually negative. (2) Bcl-6 is an important prognostic marker in diffuse large B-cell lymphomas (DLBCL), where CD10, bcl-6 and MUM1/IRF4 are used to identify germinal center and activated B-cell phenotypes. (3) Bcl-6 can be valuable in distinguishing classical Hodgkin lymphoma from nodular lymphocyte predominant Hodgkin lymphoma (NLPHL). The Reed-Sternberg cells of classical Hodgkin lymphoma are bcl-6 negative whereas the large ('LH') cells of NLPHL are bcl-6 positive. In contrast, anti-Bcl-6 rarely stains mantle-cell lymphoma and MALT lymphoma.ImmunogenRecombinant human bcl-6 protein fragment (around aa256-389) (exact	Host	Mouse			
SpeciesHumanMarkerFollicular Lymphoma MarkerSpecificity/SensitivityThe specificity of this monoclonal antibody to its intended target was validated by HuProtTM Array, containing more than 19,000, full-length human proteins. Recognizes a protein of 95kDa, which is identified as Bcl-6. Antibody to bcl-6 is helpful in a number of diagnostic settings: (1) In the differential diagnosis of small B-cell lymphoma. Follicular lymphoma will show bcl-6 (and CD10) positivity whereas other small B-cell lymphomas are usually negative. (2) Bcl-6 is an important prognostic marker in diffuse large B-cell lymphomas (DLBCL), where CD10, bcl-6 and MUM1/IRF4 are used to identify germinal center and activated B-cell phenotypes. (3) Bcl-6 can be valuable in distinguishing classical Hodgkin lymphoma from nodular lymphocyte predominant Hodgkin lymphoma (NLPHL). The Reed-Sternberg cells of classical Hodgkin lymphoma are bcl-6 negative whereas the large ('LH') cells of NLPHL are bcl-6 positive. In contrast, anti-Bcl-6 rarely stains mantle-cell lymphoma and MALT lymphoma.ImmunogenRecombinant human bcl-6 protein fragment (around aa256-389) (exact	Gene ID	604			
MarkerFollicular Lymphoma MarkerSpecificity/SensitivityThe specificity of this monoclonal antibody to its intended target was validated by HuProtTM Array, containing more than 19,000, full-length human proteins. Recognizes a protein of 95kDa, which is identified as Bcl-6. Antibody to bcl-6 is helpful in a number of diagnostic settings: (1) In the differential diagnosis of small B-cell lymphoma. Follicular lymphoma will show bcl-6 (and CD10) positivity whereas other small B-cell lymphomas are usually negative. (2) Bcl-6 is an important prognostic marker in diffuse large B-cell lymphomas (DLBCL), where CD10, bcl-6 and MUM1/IRF4 are used to identify germinal center and activated B-cell phenotypes. (3) Bcl-6 can be valuable in distinguishing classical Hodgkin lymphoma from nodular lymphocyte predominant Hodgkin lymphoma (NLPHL). The Reed-Sternberg cells of classical Hodgkin lymphoma are bcl-6 negative whereas the large ('LH') cells of NLPHL are bcl-6 positive. In contrast, anti-Bcl-6 rarely stains mantle-cell lymphoma and MALT lymphoma.ImmunogenRecombinant human bcl-6 protein fragment (around aa256-389) (exact	Gene Symbol	BCL6			
Specificity/SensitivityThe specificity of this monoclonal antibody to its intended target was validated by HuProtTM Array, containing more than 19,000, full-length human proteins. Recognizes a protein of 95kDa, which is identified as Bcl-6. Antibody to bcl-6 is helpful in a number of diagnostic settings: (1) In the differential diagnosis of small B-cell lymphoma. Follicular lymphoma will show bcl-6 (and CD10) positivity whereas other small B-cell lymphomas are usually negative. (2) Bcl-6 is an important prognostic marker in diffuse large B-cell lymphomas (DLBCL), where CD10, bcl-6 and MUM1/IRF4 are used to identify germinal center and activated B-cell phenotypes. (3) Bcl-6 can be valuable in distinguishing classical Hodgkin lymphoma from nodular lymphocyte predominant Hodgkin lymphoma (NLPHL). The Reed-Sternberg cells of classical Hodgkin lymphoma are bcl-6 negative whereas the large ('LH') cells of NLPHL are bcl-6 positive. In contrast, anti-Bcl-6 rarely stains mantle-cell lymphoma and MALT lymphoma.ImmunogenRecombinant human bcl-6 protein fragment (around aa256-389) (exact	Species	Human			
HuProtTM Array, containing more than 19,000, full-length human proteins. Recognizes a protein of 95kDa, which is identified as Bcl-6. Antibody to bcl-6 is helpful in a number of diagnostic settings: (1) In the differential diagnosis of small B-cell lymphoma. Follicular lymphoma will show bcl-6 (and CD10) positivity whereas other small B-cell lymphomas are usually negative. (2) Bcl-6 is an important prognostic marker in diffuse large B-cell lymphomas (DLBCL), where CD10, bcl-6 and MUM1/IRF4 are used to identify germinal center and activated B-cell phenotypes. (3) Bcl-6 can be valuable in distinguishing classical Hodgkin lymphoma from nodular lymphocyte predominant Hodgkin lymphoma (NLPHL). The Reed-Sternberg cells of classical Hodgkin lymphoma are bcl-6 negative whereas the large ('LH') cells of NLPHL are bcl-6 positive. In contrast, anti-Bcl-6 rarely stains mantle-cell lymphoma and MALT lymphoma.ImmunogenRecombinant human bcl-6 protein fragment (around aa256-389) (exact	Marker	Follicular Lymphoma Marker			
	Specificity/Sensitivity	HuProtTM Array, containing more than 19,000, full-length human proteins. Recognizes a protein of 95kDa, which is identified as Bcl-6. Antibody to bcl-6 is helpful in a number of diagnostic settings: (1) In the differential diagnosis of small B-cell lymphoma. Follicular lymphoma will show bcl-6 (and CD10) positivity whereas other small B-cell lymphomas are usually negative. (2) Bcl-6 is an important prognostic marker in diffuse large B-cell lymphomas (DLBCL), where CD10, bcl-6 and MUM1/IRF4 are used to identify germinal center and activated B-cell phenotypes. (3) Bcl-6 can be valuable in distinguishing classical Hodgkin lymphoma from nodular lymphocyte predominant Hodgkin lymphoma (NLPHL). The Reed-Sternberg cells of classical Hodgkin lymphoma are bcl-6 negative whereas the large ('LH') cells of NLPHL are bcl-6 positive. In contrast, anti-Bcl-6			
	Immunogen				



Notes	Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.
Product Application Details	
Applications	Western Blot, Flow Cytometry, Protein Array, CyTOF-ready

Applications	vvestern Blot, Flow Cytometry, Protein Array, CyTOF-ready
<b>Recommended Dilutions</b>	Western Blot, Flow Cytometry, Protein Array, CyTOF-ready
Application Notes	Optimal dilution of this antibody should be experimentally determined.

#### Images

Bcl-6 Antibody (BCL6/1475) [Alexa Fluor® 700] [NBP2-59597AF700] - Vial of Alexa Fluor 700 conjugated antibody. Alexa Fluor 700 is optimally excited at 696 nm by the Red laser (633 or 640 nm) and has an emission maximum of 720 nm.				
		AlexaFluo	r® 700	
	Alexa Fluor® 700	Red (633,640		
		EXCITATION MAX (nm)	EMISSION MAX (nm) 720	

www.novusbio.com



technical@novusbio.com



#### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

#### Products Related to NBP2-59597AF700

IC002N H00000604-Q01-10ug	Mouse IgG1 Isotype Control (11711) [Alexa Fluor® 700] Recombinant Human Bcl-6 GST (N-Term) Protein
7268-CT-100	CTLA-4 [Unconjugated]
NBL1-07951	Bcl-6 Overexpression Lysate

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-59597AF700

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

