Product Datasheet

Enolase 2/Neuron-specific Enolase Antibody (ENO2/1462) [Alexa Fluor® 750] NBP2-59603AF750

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-59603AF750

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-59603AF750



NBP2-59603AF750

Enolase 2/Neuron-specific Enolase Antibody (ENO2/1462) [Alexa Fluor® 750]

Product Information Unit Size 0.1 ml Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark. Clonality Monoclonal Clone ENO2/1462 Preservative 0.05% Sodium Azide Isotype IgG2b Kappa Conjgate Alexa Fluor 750 Purity Protein A or G purified Buffer SomM Sodium Borate Product Description 2026 Gene ID 2026 Gene Symbol ENO2 Species Human, Mouse, Rat Marker Neuroendocrine Marker Specificity/Sensitivity The specificity of this monoclonal antibody to its intended target was validated by HuProtTM Array, containing more than 19,000, full-length human proteins. Recognizes a protein of about 50KDa, which is identified a gamma-enolase. Three isoenzymes of enolases are identified, alpha, beta and gamma. Alpha-isoform is expressed in mores tissues, whereas beta-form is expressed predominantly in muscle tissue whereas gamma-enolase. Three isoenzymes of enolases are identified, alpha, beta and gamma-enolase. Three isoenzymes of enolases are identified, alpha, beta and gamma-enolase. Three isoenzymes of enolases are identified, alpha, beta and gamma-enolase. Three isoenzymes of enolases are iden		
ConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneENO2/1462Preservative0.05% Sodium AzideIsotypeIgG2b KappaConjugateAlexa Fluor 750PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description2026Gene ID2026Gene SymbolENO2SpeciesHuman, Mouse, RatMarkerNeuroendocrine MarkerSpecificity/SensitivityThe specificity of this monoclonal antibody to its intended target was validated by HuProtTIM Array, containing more than 19,000, full-length human proteins. Recognizes a protein of about 50KDa, which is identified as gamma-enolase. Three isoenzymes of enolases are identified, alpha, beta and gamma. Alpha- isoform is expressed in most tissue, whereas beta-form is expressed predominantly in muscle tissue whereas gamma-enolase. It it be ylay a role in converting phosphoglyceric acid to phosphenolpyruvic acid in the glycolytic pathway. NSE-gamma is a useful marker to identify peripheral nervous sits as both homodimers and heterodimers, and they play a role in converting phosphoglyceric acid to phosphenolpyruvic acid in the glycolytic pathway. NSE-gamma is a useful marker to identify peripheral nervos sits as both homodimers. And heterodimers, and they play a role in converting phosphoglyceric acid to phosphenolpyruvic acid in the glycolytic pathway. NSE-gamma is a useful marker to identify peripheral nervos sits as both homodimers. It it be usually employed in combination with other markers such as Synaptophysin, Chromogranin A, and Neurofilame	Product Information	
services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneENO2/1462Preservative0.05% Sodium AzideIsotypeIgG2b KappaConjugateAlexa Fluor 750PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description2026HostMouseGene ID2026Specificity/SensitivityThe specificity of this monoclonal antibody to its intended target was validated by HuProtTM Array, containing more than 19.000, full-length human proteins. Recognizes a protein of about 50kDa, which is identified as gamma-enolase. Three iscorrymes of enolases agient in use sties whereas gamma-enolase. Three iscorrymes of enolases agient in the glycolytic pathway. NSE-gamma is a useful marker to identify peripheral nervous tissue. These isoforms exist as both homodimers and heterodimers, and the ylay a role in converting phosphoglyceic acid to phosphonolpyruvic acid in nervous tissue. These isoforms exist as both homodimers and heterodimers, and the ylay a role in converting phosphoglyceic acid to phosphonolpyruvic acid in nervous tissue. These isoforms exist as both homodimers and heterodimers, and the ylay a role in converting phosphoglyceic acid to phosphonolpyruvic acid in nervous tissue. These isoforms exist as both homodimers and heterodimers, and the ylay a role in converting phosphoglyceic acid to phosphonolpyruvic acid in nervous tissue. These isoforms exist as both homodimers and heterodimers, and they ylay a role in converting phosphoglyceic acid to phosphonolpyruvic acid in nervoes and tumors of neuro-endocrine origins, such as pheochromocytomas. It it be usually employed in combination with other markers such as Synaptophysin, Chromogranin	Unit Size	0.1 ml
ClonalityMonoclonalCloneENO2/1462Preservative0.05% Sodium AzideIsotypeIgG2b KappaConjugateAlexa Fluor 750PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description2026Gene ID2026Gene SymbolENO2SpeciesHuman, Mouse, RatMarkerNeuroendocrine MarkerSpecificity/SensitivityThe specificity of this monoclonal antibody to its intended target was validated by HuProTTM Array, containing more than 19,000, full-length human proteins. Recognizes a protein of about SOkDa, which is is es form only in nervous tissue. These isoforms exist as both homodimers and heterodimers, and the yplay a role in converting phosphenglyceric acid to phosphenolpyruic acid in the yplay a role in converting phosphoglyceric acid to phosphenolpyruic acid in the yplay a role in converting phosphoglyceric acid to phosphenolpyruic acid in the yplay a role in converting phosphoglyceric acid to phosphenolpyruic acid in the yplay a role in converting phosphoglyceric acid to phosphenolpyruic acid in the yplay a role in converting phosphoglyceric acid to phosphenolpyruic acid in the yplay a role in converting phosphoglyceric acid to phosphenolpyruic acid in the yplay a role in converting phosphoglyceric acid to synaptophysin, Chromogranin A, and Neurofilament.ImmunogenA synthetic peptide of human Enolase 2/Neuron-specific Enolase (around aa416-	Concentration	
CloneENO2/1462Preservative0.05% Sodium AzideIsotypeIgG2b KappaConjugateAlexa Fluor 750PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description2026HostMouseGene ID2026Gene SymbolENO2SpeciesHuman, Mouse, RatMarkerNeuroendocrine MarkerSpecificity/SensitivityThe specificity of this monoclonal antibody to its intended target was validated by HuProtTM Array, containing more than 19,000, full-length human proteins. Recognizes a protein of about 50kDa, which is identified as gamma-enolase. Three isoenzymes of enolases are identified, alpha, beta and gamma. Alpha- isoform is expressed in most itsues, whereas beta-form is expressed predominantly in muscle tissues whereas gamma-enolase is found only in nervous tissue. These isoforms exist as both homodimers and heterodimers, and they play a role in converting phosphoglyceric acid to phosphenolpyruvic acid in the yplay a role in converting phosphoglyceris such as Synaptophysin, Chromogranin A, and Neurofilament.ImmunogenA synthetic peptide of human Enolase 2/Neuron-specific Enolase (around aa416-	Storage	Store at 4C in the dark.
Preservative0.05% Sodium AzideIsotypeIgG2b KappaConjugateAlexa Fluor 750PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID2026Gene SymbolENO2SpeciesHuman, Mouse, RatMarkerNeuroendocrine MarkerSpecificity/SensitivityThe specificity of this monoclonal antibody to its intended target was validated by HuProtTM Array, containing more than 19,000, full-length human proteins. Recognizes a protein of about 50kDa, which is identified as gamma-enolase. Three isoenzymes of enolases are identified, alpha, beta and gamma. Alpha- isoform is expressed in most tissues, whereas beta-form is expressed ipredominantly in muscle tissue whereas gamma-enolase is found only in nervous tissue. These isoforms exist as both homodimers and heterodimers, and hey play a role in converting phosphoglyceric acid to phosphenolpyruvic acid in the glycolytic pathway. NSE-gamma is a useful marker to identify peripheral nervoes and tumors of neuro-endocrine origins, such as pheochromocytomas. It it be usually employed in combination with other markers such as Synaptophysin, Chromogranin A, and Neurofilament.ImmunogenA synthetic peptide of human Enolase 2/Neuron-specific Enolase (around aa416-	Clonality	Monoclonal
IsotypeIgG2b KappaConjugateAlexa Fluor 750PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID2026Gene SymbolENO2SpeciesHuman, Mouse, RatMarkerNeuroendocrine MarkerSpecificity/SensitivityThe specificity of this monoclonal antibody to its intended target was validated by HuProtTM Array, containing more than 19,000, full-length human proteins. Recognizes a protein of about 50kDa, which is identified as gamma-enolase. Three isoenzymes of enolases are identified, alpha, beta and gamma. Alpha- isoform is expressed in most tissues, whereas gamma-enolase is found only in nervous tissue. These isoforms exist as both homodimers, and they glay a role in converting phosphoglyceric acid to phosphenolpyruvic acid in the glycolytic pathway. NSE-gamma is a useful marker to identify peripheral nerves and tumors of neuro-endocrine origins, such as pheochromocytomas. It it be usually employed in combination with other markers such as Synaptophysin, Chromogranin A, and Neurofilament.ImmunogenA synthetic peptide of human Enolase 2/Neuron-specific Enolase (around aa416-	Clone	ENO2/1462
ConjugateAlexa Fluor 750PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID2026Gene SymbolENO2SpeciesHuman, Mouse, RatMarkerNeuroendocrine MarkerSpecificity/SensitivityThe specificity of this monoclonal antibody to its intended target was validated by HuProtTM Array, containing more than 19,000, full-length human proteins. Recognizes a protein of about 50kDa, which is identified as gamma-enolase. Three isoenzymes of enolases are identified, alpha, beta and gamma. Alpha- isoform is expressed in most tissues, whereas beta-form is expressed predominantly in muscle tissue whereas gamma-enolase is found only in nervous tissue. These isoforms exist as both homodimers and heterodimers, and they play a role in converting phosphoglyceric acid to phosphenolpyruvic acid in the glycolytic pathway. NSE-gamma is a useful marker to identify peripheral nerves and tumors of neuro-endocrine origins, such as Spnaptophysin, Chromogranin A, and Neurofilament.ImmunogenA synthetic peptide of human Enolase 2/Neuron-specific Enolase (around aa416-	Preservative	0.05% Sodium Azide
PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID2026Gene SymbolENO2SpeciesHuman, Mouse, RatMarkerNeuroendocrine MarkerSpecificity/SensitivityThe specificity of this monoclonal antibody to its intended target was validated by HuProtTM Array, containing more than 19,000, full-length human proteins. Recognizes a protein of about 50kDa, which is identified as gamma-enolase. Three isoenzymes of enolases are identified, alpha, beta and gamma. Alpha- isoform is expressed in most tissues, whereas beta-form is expressed predominantly in muscle tissue whereas gamma-enolase is found only in nervous tissue. These isoforms exist as both homodimers and heterodimers, and they play a role in converting phosphoglyceric acid to phosphenolpyruvic acid in the glycolytic pathway. NSE-gamma is a useful marker to identify peripheral nerves and tumors of neuro-endocrine origins, such as Synaptophysin, Chromogranin A, and Neurofilament.ImmunogenA synthetic peptide of human Enolase 2/Neuron-specific Enolase (around aa416-	Isotype	IgG2b Kappa
Buffer50mM Sodium BorateProduct DescriptionHostMouseGene ID2026Gene SymbolENO2SpeciesHuman, Mouse, RatMarkerNeuroendocrine MarkerSpecificity/SensitivityThe specificity of this monoclonal antibody to its intended target was validated by HuProtTM Array, containing more than 19,000, full-length human proteins. Recognizes a protein of about 50kDa, which is identified as gamma-enolase. Three isoenzymes of enolases are identified, alpha, beta and gamma. Alpha- isoform is expressed in most tissues, whereas beta-form is expressed predominantly in muscle tissue whereas gamma-enolase is found only in nervous tissue. These isoforms exist as both homodimers and heterodimers, and they play a role in converting phosphoglyceric acid to phosphenolpyruvic acid in the glycolytic pathway. NSE-gamma is a useful marker to identify peripheral nerves and tumors of neuro-endocrine origins, such as pheochromocytomas. It it be usually employed in combination with other markers such as Synaptophysin, Chromogranin A, and Neurofilament.ImmunogenA synthetic peptide of human Enolase 2/Neuron-specific Enolase (around aa416-	Conjugate	Alexa Fluor 750
Product DescriptionHostMouseGene ID2026Gene SymbolENO2SpeciesHuman, Mouse, RatMarkerNeuroendocrine MarkerSpecificity/SensitivityThe specificity of this monoclonal antibody to its intended target was validated by HuProtTM Array, containing more than 19,000, full-length human proteins. Recognizes a protein of about 50kDa, which is identified as gamma-enolase. Three isoenzymes of enolases are identified, alpha, beta and gamma. Alpha- isoform is expressed in most tissues, whereas beta-form is expressed predominantly in muscle tissue whereas gamma-enolase is found only in nervous tissue. These isoforms exist as both homodimers and heterodimers, and they glycolytic pathway. NSE-gamma is a useful marker to identify peripheral nerves and tumors of neuro-endocrine origins, such as pheochromocytomas. It it be usually employed in combination with other markers such as Synaptophysin, Chromogranin A, and Neurofilament.ImmunogenA synthetic peptide of human Enolase 2/Neuron-specific Enolase (around aa416-	Purity	Protein A or G purified
HostMouseGene ID2026Gene SymbolENO2SpeciesHuman, Mouse, RatMarkerNeuroendocrine MarkerSpecificity/SensitivityThe specificity of this monoclonal antibody to its intended target was validated by HuProtTM Array, containing more than 19,000, full-length human proteins. Recognizes a protein of about 50kDa, which is identified as gamma-enolase. Three isoenzymes of enolases are identified, alpha, beta and gamma. Alpha- isoform is expressed in most tissues, whereas beta-form is expressed predominantly in muscle tissue whereas gamma-enolase is found only in nervous tissue. These isoforms exist as both homodimers and heterodimers, and the glycolytic pathway. NSE-gamma is a useful marker to identify peripheral nerves and tumors of neuro-endocrine origins, such as pheochromocytomas. It it be usually employed in combination with other markers such as Synaptophysin, Chromogranin A, and Neurofilament.ImmunogenA synthetic peptide of human Enolase 2/Neuron-specific Enolase (around aa416-	Buffer	50mM Sodium Borate
Gene ID2026Gene SymbolENO2SpeciesHuman, Mouse, RatMarkerNeuroendocrine MarkerSpecificity/SensitivityThe specificity of this monoclonal antibody to its intended target was validated by HuProtTM Array, containing more than 19,000, full-length human proteins. Recognizes a protein of about 50kDa, which is identified as gamma-enolase. Three isoenzymes of enolases are identified, alpha, beta and gamma. Alpha- isoform is expressed in most tissues, whereas beta-form is expressed predominantly in muscle tissue whereas gamma-enolase is found only in nervous tissue. These isoforms exist as both homodimers and heterodimers, and they play a role in converting phosphoglyceric acid to phosphenolpyruvic acid in the glycolytic pathway. NSE-gamma is a useful marker to identify peripheral nerves and tumors of neuro-endocrine origins, such as pheochromocytomas. It it be usually employed in combination with other markers such as Synaptophysin, Chromogranin A, and Neurofilament.ImmunogenA synthetic peptide of human Enolase 2/Neuron-specific Enolase (around aa416-	Product Description	
Gene SymbolENO2SpeciesHuman, Mouse, RatMarkerNeuroendocrine MarkerSpecificity/SensitivityThe specificity of this monoclonal antibody to its intended target was validated by HuProtTM Array, containing more than 19,000, full-length human proteins. Recognizes a protein of about 50kDa, which is identified as gamma-enolase. Three isoenzymes of enolases are identified, alpha, beta and gamma. Alpha- isoform is expressed in most tissues, whereas beta-form is expressed predominantly in muscle tissue whereas gamma-enolase is found only in nervous tissue. These isoforms exist as both homodimers and heterodimers, and they play a role in converting phosphoglyceric acid to phosphenolpyruvic acid in the glycolytic pathway. NSE-gamma is a useful marker to identify peripheral nerves and tumors of neuro-endocrine origins, such as pheochromocytomas. It it be usually employed in combination with other markers such as Synaptophysin, Chromogranin A, and Neurofilament.ImmunogenA synthetic peptide of human Enolase 2/Neuron-specific Enolase (around aa416-	Host	Mouse
SpeciesHuman, Mouse, RatMarkerNeuroendocrine MarkerSpecificity/SensitivityThe specificity of this monoclonal antibody to its intended target was validated by HuProtTM Array, containing more than 19,000, full-length human proteins. Recognizes a protein of about 50kDa, which is identified as gamma-enolase. Three isoenzymes of enolases are identified, alpha, beta and gamma. Alpha- isoform is expressed in most tissues, whereas beta-form is expressed predominantly in muscle tissue whereas gamma-enolase is found only in nervous tissue. These isoforms exist as both homodimers and heterodimers, and they play a role in converting phosphoglyceric acid to phosphenolpyruvic acid in the glycolytic pathway. NSE-gamma is a useful marker to identify peripheral nerves and tumors of neuro-endocrine origins, such as pheochromocytomas. It it be usually employed in combination with other markers such as Synaptophysin, Chromogranin A, and Neurofilament.ImmunogenA synthetic peptide of human Enolase 2/Neuron-specific Enolase (around aa416-	Gene ID	2026
MarkerNeuroendocrine MarkerSpecificity/SensitivityThe specificity of this monoclonal antibody to its intended target was validated by HuProtTM Array, containing more than 19,000, full-length human proteins. Recognizes a protein of about 50kDa, which is identified as gamma-enolase. Three isoenzymes of enolases are identified, alpha, beta and gamma. Alpha- isoform is expressed in most tissues, whereas beta-form is expressed predominantly in muscle tissue whereas gamma-enolase is found only in nervous tissue. These isoforms exist as both homodimers and heterodimers, and they play a role in converting phosphoglyceric acid to phosphenolpyruvic acid in the glycolytic pathway. NSE-gamma is a useful marker to identify peripheral nerves and tumors of neuro-endocrine origins, such as pheochromocytomas. It it be usually employed in combination with other markers such as Synaptophysin, Chromogranin A, and Neurofilament.ImmunogenA synthetic peptide of human Enolase 2/Neuron-specific Enolase (around aa416-	Gene Symbol	ENO2
Specificity/SensitivityThe specificity of this monoclonal antibody to its intended target was validated by HuProtTM Array, containing more than 19,000, full-length human proteins. Recognizes a protein of about 50kDa, which is identified as gamma-enolase. Three isoenzymes of enolases are identified, alpha, beta and gamma. Alpha- isoform is expressed in most tissues, whereas beta-form is expressed predominantly in muscle tissue whereas gamma-enolase is found only in nervous tissue. These isoforms exist as both homodimers and heterodimers, and they play a role in converting phosphoglyceric acid to phosphenolpyruvic acid in the glycolytic pathway. NSE-gamma is a useful marker to identify peripheral nerves and tumors of neuro-endocrine origins, such as pheochromocytomas. It it be usually employed in combination with other markers such as Synaptophysin, Chromogranin A, and Neurofilament.ImmunogenA synthetic peptide of human Enolase 2/Neuron-specific Enolase (around aa416-	Species	Human, Mouse, Rat
 HuProtTM Array, containing more than 19,000, full-length human proteins. Recognizes a protein of about 50kDa, which is identified as gamma-enolase. Three isoenzymes of enolases are identified, alpha, beta and gamma. Alpha- isoform is expressed in most tissues, whereas beta-form is expressed predominantly in muscle tissue whereas gamma-enolase is found only in nervous tissue. These isoforms exist as both homodimers and heterodimers, and they play a role in converting phosphoglyceric acid to phosphenolpyruvic acid in the glycolytic pathway. NSE-gamma is a useful marker to identify peripheral nerves and tumors of neuro-endocrine origins, such as pheochromocytomas. It it be usually employed in combination with other markers such as Synaptophysin, Chromogranin A, and Neurofilament. Immunogen 	Marker	Neuroendocrine Marker
	Specificity/Sensitivity	HuProtTM Array, containing more than 19,000, full-length human proteins. Recognizes a protein of about 50kDa, which is identified as gamma-enolase. Three isoenzymes of enolases are identified, alpha, beta and gamma. Alpha- isoform is expressed in most tissues, whereas beta-form is expressed predominantly in muscle tissue whereas gamma-enolase is found only in nervous tissue. These isoforms exist as both homodimers and heterodimers, and they play a role in converting phosphoglyceric acid to phosphenolpyruvic acid in the glycolytic pathway. NSE-gamma is a useful marker to identify peripheral nerves and tumors of neuro-endocrine origins, such as pheochromocytomas. It it be usually employed in combination with other markers such as Synaptophysin, Chromogranin A, and Neurofilament.
	Immunogen	

www.novusbio.com



	Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.
Product Application Details	
Applications	Immunohistochemistry, Immunohistochemistry-Paraffin, Protein Array, CyTOF- ready
Recommended Dilutions	Immunohistochemistry, Immunohistochemistry-Paraffin, Protein Array, CyTOF- ready
Application Notes	Optimal dilution of this antibody should be experimentally determined.

Notes





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-59603AF750

NBP1-43317AF750	Mouse IgG2b Kappa Light Chain Isotype Control (MG2b) [Alexa Fluor® 750]
NBP2-61382-1mg	Recombinant Human Enolase 2/Neuron-specific Enolase Protein
7954-GM-010/CF	GM-CSF [Unconjugated]
DENL20	Enolase 2/Neuron-specific Enolase [HRP]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-59603AF750

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

