

Product Datasheet

LC3B Antibody (1251D) - BSA Free NBP2-59800

Unit Size: 0.1 mg

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 1

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-59800

Updated 10/23/2024 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-59800



NBP2-59800

LC3B Antibody (1251D) - BSA Free


Product Information	
Unit Size	0.1 mg
Concentration	1.0 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	1251D
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Protein A or G purified
Buffer	PBS
Target Molecular Weight	14.688 kDa

Product Description	
Host	Rabbit
Gene ID	81631
Gene Symbol	MAP1LC3B
Species	Human, Mouse, Rat
Immunogen	Recombinant monoclonal LC3B Antibody (1251D) was made to a synthetic peptide made to an N-terminal portion of the human LC3B protein sequence (between residues 1-100). [Uniprot: Q9GZQ8].


Product Application Details	
Applications	Western Blot, Simple Western, Flow (Intracellular), Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Knockout Validated
Recommended Dilutions	Western Blot 1 - 2 ug/ml, Simple Western 5 ug/ml, Immunohistochemistry 2 - 5ug/ml, Immunocytochemistry/ Immunofluorescence 5 ug/ml, Immunohistochemistry-Paraffin 2 - 5ug/ml, Flow (Intracellular) 1 ug/ml, Knockout Validated
Application Notes	Western blot bands are seen at ~19 kDa, representing LC3-I, and ~17 kDa, representing LC3-II.

Images


Knockout Validated: LC3B Antibody (1251D) [NBP2-59800] - HeLa human cervical epithelial carcinoma parental cell line and LC3B knockout HeLa cell line (KO) untreated (-) or treated (+) with 50 uM Chloroquine for 18 hrs. PVDF membrane was probed with 0.1 ug/mL of Rabbit Anti-LC3B Monoclonal Antibody (1251D) [Catalog# NBP2-59800] followed by HRP-conjugated Anti-Rabbit IgG Secondary Antibody (Catalog# HAF008). A specific band was detected for LC3B at approximately 15 kDa (as indicated) in the parental HeLa cell line, but is not detectable in the knockout HeLa cell line. GAPDH (Catalog # AF5718) is shown as a loading control. This experiment was conducted under reducing conditions.




Western Blot: LC3B Antibody (1251D) [NBP2-59800] - Total protein from HeLa and Neuro2A cells treated with or without 50 μ M chloroquine for 24 hrs was separated on a 4-15% gel by SDS-PAGE, transferred to 0.2 μ m PVDF (Polyvinylidene difluoride) membrane and blocked in 5% non-fat milk in TBST (Tris-buffered saline, 0.1% Tween 20). The membrane was probed with 2.0 μ g/ml anti-LC3B Antibody (1251d) [Catalog # NBP2-59800] in blocking buffer and detected with an anti-rabbit HRP secondary antibody using chemiluminescence. Note the detection of LC3I and LC3II at a molecular weight of approximately 15 kDa.




Immunocytochemistry/Immunofluorescence: LC3B Antibody (1251D) [NBP2-59800] - HeLa cells were treated with 50 μ M CQ (Chloroquine) overnight, fixed for 10 minutes using 10% formalin and then permeabilized for 5 minutes using 1X TBS + 0.5% Triton X-100. The cells were incubated with anti-LC3B (1251d) [Catalog # NBP2-59800] at 5 μ g/ml overnight at 4C and detected with an anti-mouse DyLight 488 (Green) at a 1:500 dilution. Actin was detected with Phalloidin 568 (Red) at a 1:200 dilution. Nuclei were counterstained with DAPI (Blue). Cells were imaged using a 40X objective.




Immunohistochemistry-Paraffin: LC3B Antibody (1251D) [NBP2-59800] - Analysis of a FFPE (Formalin-Fixed Paraffin-Embedded) tissue section of mouse brain using anti-LC3B antibody (clone 1251D) [Catalog # NBP2-59800] at 5 μ g/ml concentration (1:200 dilution). The primary antibody binding to LC3 in cells was detected using HRP conjugated anti-Rabbit secondary antibody with DAB (3,3'Diaminobenzidine) reagent, and the sections were further counterstained with hematoxylin for labeling cellular nuclei. This LC3 antibody generated a diffused staining in all cell types (except the endothelial cells of blood vessels) and the signal was strongest in a subset of neuronal cells.



Flow (Intracellular): LC3B Antibody (1251D) [NBP2-59800] - HeLa cells were either (A) untreated or (B) treated with 50 μ M chloroquine for 24 hrs. An intracellular stain was performed with anti-LC3B (1251D) antibody [Catalog # NBP2-59800] (blue) and a matched isotype control [Catalog # MAB1050] (orange). Cells were fixed with 4% paraformaldehyde, following fixation, cells were permeabilized with 0.1% saponin. Cells were incubated in an antibody dilution of 1 μ g/mL for 30 minutes at room temperature, followed by rabbit IgG APC-conjugated secondary antibody (F0111, R&D Systems).



Simple Western: LC3B Antibody (1251D) [NBP2-59800] - Lane view shows lysates of NIH-3T3 mouse embryonic fibroblast cell line untreated (-) or treated (+) with 50 uM Chloroquine for 18 hrs, loaded at 0.2 mg/ml. A specific band was detected for LC3B at approximately 17 kDa (as indicated) using 5 ug/mL of Rabbit Anti-Human LC3B Monoclonal Antibody (1251D) [Catalog# NBP2-59800]. This experiment was conducted under reducing conditions and using the 12-230 kDa separation system.



Publications

Klionsky DJ, Abdel-Aziz AK, Abdelfatah S et al. Guidelines for the use and interpretation of assays for monitoring autophagy (4th edition)1 Autophagy 2021-01-01 [PMID: 33634751] (WB)

Procedures

Western Blot protocol for LC3B Antibody (NBP2-59800)

LC3B Antibody (1251D):

Protocol: Inhibition of Autophagy and LC3 Antibody (NBP2-59800) Western Blot

Materials

Chloroquine diphosphate (CQ) (10 mM) in dH₂O

1X PBS

Sample buffer, 2X Laemmli buffer: 4% SDS, 5% 2-mercaptoethanol (BME), 20% glycerol, 0.004% bromophenol blue, 0.125 M Tris HCl, pH 6.8

RIPA buffer: 150 mM NaCl, 1% NP-40 or Triton X-100, 0.5% sodium deoxycholate, 0.1% SDS, 50 mM Tris-HCl, pH 8.0, 20 mM Tris-HCl, pH 7.5

1X Running Buffer: 25 mM Tris-base, 192 mM glycine, 0.1% SDS. Adjust to pH 8.3

1X Transfer buffer (wet): 25 mM Tris-base, 192 mM glycine, 20% methanol, Adjust to pH 8.3

TBS

TBST, TBS and 0.1% Tween

Blocking solution: TBST, 5% non-fat dry milk

rabbit anti-LC3 primary antibody (NBP2-59800) in blocking buffer (~2 ug/mL)

Methods

Tip: For more information on Western Blotting, see our Western Blot handbook.

1. Grow cells (e.g. HeLa or Neuro2A) in vitro to semi-confluency (70-75%).
2. Add CQ to culture dishes to a final concentration of 50 uM and incubate overnight (16 hours). Remember to include an untreated sample as a negative control.
Note: Validated autophagy inducers should be included as positive controls.
3. Rinse cells with ice-cold 1X PBS and lyse cells with sample buffer.
Note: LC3-I and LC3-II are sensitive to degradation, although LC3-I is more labile. These proteins are sensitive to freeze-thaw cycles and SDS sample buffers. Fresh samples should be analyzed quickly to prevent protein degradation.
4. Sonicate and incubate cells for 5 minutes at 95oC.
Tip: Cells are lysed directly in sample buffer or may be lysed in RIPA buffer.
5. Load samples of Chloroquine-treated and -untreated cell lysates 40 ug/lane on a 4-20% polyacrylamide gradient gel (SDS-PAGE).
Tip: For detection of LC3 it is particularly important to monitor the progress of the gel as this protein is relatively small (~14kDa).
Tip: Alternatively, for non-gradient gels, use a 20% polyacrylamide gel.
6. Transfer proteins to a 0.2 um PVDF membrane for 30 minutes at 100V.
7. After transfer, rinse the membrane with dH₂O and stain with Ponceau S for 1-2 minutes to confirm efficiency of protein transfer.
8. Rinse the membrane in dH₂O to remove excess stain and mark the loaded lanes and molecular weight markers using a pencil.
9. Block the membrane using blocking buffer solution (5% non-fat dry milk in TBST) for 1 hour at room temperature.
10. Rinse the membrane with TBST for 5 minutes.
11. Dilute the rabbit anti-LC3 primary antibody (NBP2-59800) (~2 ug/mL) in blocking buffer and incubate the

membrane for 1 hour at room temperature.

12. Rinse the membrane with dH₂O.

13. Rinse the membrane with TBST, 3 times for 10 minutes each.

14. Incubate the membrane with diluted secondary antibody, according with product's specifications, (e.g. anti-rabbit-IgG HRP-conjugated) in blocking buffer for 1 hour at room temperature.

Note: Tween-20 may be added to the blocking or antibody dilution buffer at a final concentration of 0.05-0.2%, provided it does not interfere with antibody-antigen binding.

15. Rinse the membrane with TBST, 3 times for 10 minutes each.

16. Apply the detection reagent of choice (e.g. BioFX Super Plus ECL) in accordance with the manufacturer's instructions.

17. Image the blot.

Tip: LC3-I and its lipidated form LC3-II have different electrophoretic mobility properties, with the lipidated form moving faster in an SDS-PAGE gel, albeit its larger molecular weight. LC3-II runs at 14-16 kDa while LC3-I runs at 16-18kDa.

Note: This assay measures the difference in the LC3-II signal in the presence and absence of inhibitors (e.g., lysosomotropic agents). When autophagic flux is present or induced in a system an increase in the LC3-II signal should be observed with the inhibitor.



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-59800

HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control
NB100-2220PEP	LC3B Antibody Blocking Peptide

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-59800

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

