Product Datasheet

Syndecan-1/CD138 Antibody (JM11-21) NBP2-67174

Unit Size: 100 ul

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 1

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-67174

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-67174



NBP2-67174


Syndecan-1/CD138 Antibody (JM11-21)


Product Information		
Unit Size	100 ul	
Concentration	1 mg/ml	
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.	
Clonality	Monoclonal	
Clone	JM11-21	
Preservative	0.05% Sodium Azide	
Isotype	IgG	
Purity	Protein A purified	
Buffer	TBS (pH7.4), 0.05% BSA, 40% Glycerol	
Product Description		
Host	Rabbit	
Gene ID	6382	
Gene Symbol	SDC1	
Species	Human, Mouse, Rat	
Immunogen	Recombinant protein within Human Syndecan-1/CD138 aa 13-274 / 310. (SwissProt: P18827 Human; SwissProt: P18828 Mouse; SwissProt: P26260 Rat)	
Product Application Details		
Applications	Western Blot, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin	
Recommended Dilutions	Western Blot 1:500-1:1000, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1:100-1:500, Immunohistochemistry-Paraffin 1:50-1:200	
Images		
Western Blot: Syndecan-1/CD138 Antibody (JM11-21) [NBP2-67174] - Analysis of Syndecan 1 on NIH/3T3 cell lysates. Proteins were transferred to a PVDF membrane and blocked with 5% BSA in PBS for 1 hour at room temperature. The primary antibody (1/500) was used in 5% BSA at room temperature for 2 hours. Goat Anti-Rabbit IgG - HRP Secondary Antibody at 1:5,000 dilution was used for 1 hour at room temperature70 -55 -40 -35 		



-15


Immunocytochemistry/Immunofluorescence: Syndecan-1/CD138 Antibody (JM11-21) [NBP2-67174] - Staining Syndecan 1 in HepG2 cells (red). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.






Immunocytochemistry/Immunofluorescence: Syndecan-1/CD138 Antibody (JM11-21) [NBP2-67174] - Staining Syndecan 1 in A431 cells (red). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.

[NBP2-67174] - Analysis of paraffin-embedded mouse prostate tissue using anti-Syndecan 1 antibody. Counter stained with hematoxylin.



Immunocytochemistry/Immunofluorescence: Syndecan-1/CD138 Antibody (JM11-21) [NBP2-67174] - Staining Syndecan 1 in Hela cells (red). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.





Immunohistochemistry-Paraffin: Syndecan-1/CD138 Antibody (JM11-21) [NBP2-67174] - Analysis of paraffin-embedded human kidney tissue using anti-Syndecan 1 antibody. Counter stained with hematoxylin.

Immunohistochemistry-Paraffin: Syndecan-1/CD138 Antibody (JM11-21) [NBP2-67174] - Analysis of paraffin-embedded human lung tissue using anti-Syndecan 1 antibody. Counter stained with hematoxylin.

Immunohistochemistry-Paraffin: Syndecan-1/CD138 Antibody (JM11-21) [NBP2-67174] - Analysis of paraffin-embedded human tonsil tissue using anti-Syndecan 1 antibody. Counter stained with hematoxylin.


Immunohistochemistry-Paraffin: Syndecan-1/CD138 Antibody (JM11-21) [NBP2-67174] - Analysis of paraffin-embedded mouse colon tissue using anti-Syndecan 1 antibody. Counter stained with hematoxylin.


Publications

Abdullah S, Ghio M, Cotton-Betteridge A et al. Succinate metabolism and membrane reorganization drives the endotheliopathy and coagulopathy of traumatic hemorrhage Science advances 2023-06-16 [PMID: 37315138] (IP, Human)

www.novusbio.com









Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-67174

HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control
NBP1-88135PEP	Syndecan-1/CD138 Recombinant Protein Antigen

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-67174

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

