# **Product Datasheet**

# Cytokeratin 1 Antibody (SN72-08) NBP2-67681

Unit Size: 100 ul

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-67681

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP2-67681



# NBP2-67681

Cytokeratin 1 Antibody (SN72-08)


-,		
Product Information		
Unit Size	100 ul	
Concentration	1 mg/ml	
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.	
Clonality	Monoclonal	
Clone	SN72-08	
Preservative	0.05% Sodium Azide	
Isotype	IgG	
Purity	Protein A purified	
Buffer	TBS (pH7.4), 0.05% BSA, 40% Glycerol	
Product Description		
Host	Rabbit	
Gene ID	3848	

Gene Symbol	KRT1
Species	Human, Mouse, Rat
Immunogen	Synthetic peptide within mouse Cytokeratin 1 aa 586-637 / 637. (SwissProt: P04264 Human; SwissProt: P04104 Mouse; SwissProt: Q6IMF3 Rat)
<b>Product Application Details</b>	
Applications	Western Blot, Immunocytochemistry/ Immunofluorescence,


Applications	Western Blot, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin
Recommended Dilutions	Western Blot 1:500-1:1000, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1:200, Immunohistochemistry-Paraffin 1:100-1:500

Images	
Western Blot: Cytokeratin 1 Antibody (SN72-08) [NBP2-67681] - Western blot analysis of Cytokeratin 1 on different lysates. Proteins were transferred to a PVDF membrane and blocked with 5% BSA in PBS for 1 hour at room temperature. The primary antibody (1/500) was used in 5% BSA at room temperature for 2 hours. Goat	1 2 kDa -250 -150 -100 -75
	-37


Immunohistochemistry-Paraffin: Cytokeratin 1 Antibody (SN72-08) [NBP2-67681] - Analysis of paraffin-embedded rat skin tissue with Rabbit anti-Cytokeratin 1 antibody washed with ddH2O and PBS, and then probed with the primary antibody at 1/400 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.




Western Blot: Cytokeratin 1 Antibody (SN72-08) [NBP2-67681] - Analysis of Cytokeratin 1 on different lysates using anti-Cytokeratin 1 antibody at 1/1,000 dilution. Positive control: Lane 1: 293T Lane 2: Hela




Immunohistochemistry-Paraffin: Cytokeratin 1 Antibody (SN72-08) [NBP2-67681] - Analysis of paraffin-embedded mouse skin tissue using anti-Cytokeratin 1 antibody. Counter stained with hematoxylin.



Immunohistochemistry-Paraffin: Cytokeratin 1 Antibody (SN72-08) [NBP2-67681] - Immunofluorescence analysis of paraffin-embedded human skin tissue labeling Cytokeratin 1 (washed with PBS. And then probed with the primary antibodies Cytokeratin 1 (red) at 1/200 dilution and Vimentin (green) at 1/200 dilution at +4 overnight, washed with PBS. Goat Anti-Rabbit IgG H&L (iFluor(TM) 594) and Goat Anti-Mouse IgG H&L (iFluor(TM) 488) were used as the secondary antibodies at 1/1,000 dilution. Nuclei were counterstained with DAPI (blue).



Immunohistochemistry-Paraffin: Cytokeratin 1 Antibody (SN72-08) [NBP2-67681] - Analysis of paraffin-embedded human skin tissue using anti-Cytokeratin 1 antibody. The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 5% BSA for 30 minutes at room temperature, washed with ddH2O and PBS, and then probed with the primary antibody (1/400) for 30 minutes at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.





## Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

### **Products Related to NBP2-67681**

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

H00003848-Q01-10ug Recombinant Human Cytokeratin 1 GST (N-Term) Protein

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-67681

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

