# **Product Datasheet**

# MTAP Antibody (MTAP/1813) [Alexa Fluor® 488] NBP2-75731AF488

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-75731AF488

Updated 10/26/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP2-75731AF488



# NBP2-75731AF488

MTAP Antibody (MTAP/1813) [Alexa Fluor® 488]

Unit Size 0.1 ml  Concentration Please see the vial label for concentration. If unlisted please contact technical services.  Storage Store at 4C in the dark.  Clonality Monoclonal  Clone MTAP/1813  Preservative 0.05% Sodium Azide  Isotype IgG2b Kappa  Conjugate Alexa Fluor 488  Purity Protein A or G purified  Buffer 50mM Sodium Borate  Product Description  Host Mouse  Gene ID 4507  Gene Symbol MTAP  Species Human  Marker Tumor Suppressor Marker  Specificity/Sensitivity Recognizes a protein of 31kDa, which is identified as MTAP (5'-deoxy-5'-methylthioadenosine phosphorplase). It catalyzes the reversible phosphorolysis of methythioadenosine phosphorplase). It catalyzes the reversible phosphorolysis of methythioadenosine phosphorplase). It catalyzes the reversible phosphorolysis or methylthioadenosine phosphorplase). It catalyzes the reversible phosphorolysis or methythioadenosine phosphorplase). It catalyzes the reversible phosphorolysis or methythioadenosine phosphorplase). It catalyzes the reversible phosphorolysis or methythioadenosine phosphorolyaes. It catalyzes the reversible phosphorolysis or methythioadenosine phosphorolyaes. It catalyzes the reversible phosphorolysis or methythioadenosine phosphorolyaes. It catalyzes the reversible phosphorolysis or enterprise phosphoroly and the more particular and methionine. The gene encoding MTAP is linked to the tumor suppressor gene, p16INK4A, Deficient levels of MTAP can occur in cancers primarily through co-deletion of the MTAP gene and the p16INK4A gene. Cells expressing MTAP and possessing adenine salvage pathway activity may be less susceptible to malignancy due to growth-inhibitory actions of agents (e.g. antifolates), whose mechanism of action, in part, involves this de novo purine pathway.  Immunogen	WITAF Altibody (WITAF/1013) [Alexa Fluot® 400]	
Concentration  Please see the vial label for concentration. If unlisted please contact technical services.  Storage  Store at 4C in the dark.  Clonality  Monoclonal  Clone  MTAP/1813  Preservative  0.05% Sodium Azide  Isotype  IgG2b Kappa  Conjugate  Alexa Fluor 488  Purity  Protein A or G purified  Buffer  50mM Sodium Borate  Product Description  Host  Mouse  Gene ID  4507  Gene Symbol  MTAP  Species  Human  Marker  Tumor Suppressor Marker  Specificity/Sensitivity  Recognizes a protein of 31kDa, which is identified as MTAP (5'-deoxy-5'-methylthioadenosine, which is important in polyamine metabolism and for the salvage of adenine and methionine. The gene encoding MTAP is linked to the tumor suppressor gene, p16INK4A. Deficient levels of MTAP can occur in cancers primarily through co-deletion of the MTAP gene and the p16INK4A gene. Cells expressing MTAP and possessing adenine salvage pathway activity may be less susceptible to malignancy due to growth-inhibitory actions of agents (e.g. antifolates), whose mechanism of action, in part, involves this de novo purine pathway.  Immunogen	Product Information	
Storage Store at 4C in the dark.  Clonality Monoclonal  Clone MTAP/1813  Preservative 0.05% Sodium Azide  Isotype IgG2b Kappa  Conjugate Alexa Fluor 488  Purity Protein A or G purified  Buffer 50mM Sodium Borate  Product Description  Host Mouse  Gene ID 4507  Gene Symbol MTAP  Species Human  Marker Tumor Suppressor Marker  Specificity/Sensitivity Recognizes a protein of 31kDa, which is identified as MTAP (5'-deoxy-5'-methylthioadenosine, which is important in polyamine metabolism and for the salvage of adenine and methionine. The gene encoding MTAP is linked to the tumor suppressor gene, p16INK4A. Deficient levels of MTAP can occur in cancers primarily through co-deletion of the MTAP gene and the p16INK4A gene. Cells expressing MTAP and possessing adenine salvage pathway.  Immunogen Recombinant human MTAP protein fragment (aa97-196) (exact sequence is	Unit Size	0.1 ml
Clone MTAP/1813  Preservative 0.05% Sodium Azide  Isotype IgG2b Kappa Conjugate Alexa Fluor 488  Purity Protein A or G purified  Buffer 50mM Sodium Borate  Product Description  Host Mouse Gene ID 4507  Gene Symbol MTAP  Species Human  Marker Tumor Suppressor Marker  Specificity/Sensitivity Recognizes a protein of 31kDa, which is identified as MTAP (5"-deoxy-5"-methylthioadenosine, which is important in polyamine metabolism and for the salvage of adenine and methionine. The gene encoding MTAP can occur in cancers primarily through co-deletion of the MTAP gene and the p16lNK4A gene. Cells expressing MTAP and possessing adenine salvage pathway activity may be less susceptible to malignancy due to growth-inhibitory actions of agents (e.g. antifolates), whose mechanism of action, in part, involves this de novo purine pathway.  Immunogen Recombinant human MTAP protein fragment (aa97-196) (exact sequence is	Concentration	
Clone MTAP/1813  Preservative 0.05% Sodium Azide  Isotype IgG2b Kappa  Conjugate Alexa Fluor 488  Purity Protein A or G purified  Buffer 50mM Sodium Borate  Product Description  Host Mouse  Gene ID 4507  Gene Symbol MTAP  Species Human  Marker Tumor Suppressor Marker  Specificity/Sensitivity Recognizes a protein of 31kDa, which is identified as MTAP (5'-deoxy-5'-methylthioadenosine, which is important in polyamine metabolism and for the salvage of adenine and methionine. The gene encoding MTAP is linked to the tumor suppressor gene, p16INK4A. Deficient levels of MTAP can occur in cancers primarily through co-deletion of the MTAP gene and the p16INK4A gene. Cells expressing MTAP and possessing adenine salvage pathway activity may be less susceptible to malignancy due to growth-inhibitory actions of agents (e.g. antifolates), whose mechanism of action, in part, involves this de novo purine pathway.  Immunogen Recombinant human MTAP protein fragment (aa97-196) (exact sequence is	Storage	Store at 4C in the dark.
Preservative  IgG2b Kappa  Conjugate  Alexa Fluor 488  Purity  Protein A or G purified  Buffer  50mM Sodium Borate  Product Description  Host  Gene ID  4507  Gene Symbol  MTAP  Species  Human  Marker  Tumor Suppressor Marker  Specificity/Sensitivity  Recognizes a protein of 31kDa, which is identified as MTAP (5'-deoxy-5'-methylthioadenosine, which is important in polyamine metabolism and for the salvage of adenine and methionine. The gene encoding MTAP is linked to the tumor suppressor gene, p16lNk4A. Deficient levels of MTAP can occur in cancers primarily through co-deletion of the MTAP gene and the p16lNk4A gene. Cells expressing MTAP and possessing adenine salvage pathway activity may be less susceptible to malignancy due to growth-inhibitory actions of agents (e.g. antifolates), whose mechanism of action, in part, involves this de novo purine pathway.  Immunogen  Recombinant human MTAP protein fragment (aa97-196) (exact sequence is	Clonality	Monoclonal
Isotype	Clone	MTAP/1813
Conjugate  Alexa Fluor 488  Purity  Protein A or G purified  Buffer  50mM Sodium Borate  Product Description  Host  Mouse  Gene ID  4507  Gene Symbol  MTAP  Species  Human  Marker  Tumor Suppressor Marker  Specificity/Sensitivity  Recognizes a protein of 31kDa, which is identified as MTAP (5'-deoxy-5'-methylthioadenosine phosphorylase). It catalyzes the reversible phosphorolysis of methylthioadenosine, which is important in polyamine metabolism and for the salvage of adenine and methionine. The gene encoding MTAP is linked to the tumor suppressor gene, p16INK4A. Deficient levels of MTAP can occur in cancers primarily through co-deletion of the MTAP gene and the p16INK4A gene. Cells expressing MTAP and possessing adenine salvage pathway activity may be less susceptible to malignancy due to growth-inhibitory actions of agents (e.g. antifolates), whose mechanism of action, in part, involves this de novo purine pathway.  Immunogen  Recombinant human MTAP protein fragment (aa97-196) (exact sequence is	Preservative	0.05% Sodium Azide
Purity Protein A or G purified  Buffer 50mM Sodium Borate  Product Description  Host Mouse  Gene ID 4507  Gene Symbol MTAP  Species Human  Marker Tumor Suppressor Marker  Specificity/Sensitivity Recognizes a protein of 31kDa, which is identified as MTAP (5'-deoxy-5'-methylthioadenosine phosphorylase). It catalyzes the reversible phosphorolysis of methylthioadenosine, which is important in polyamine metabolism and for the salvage of adenine and methionine. The gene encoding MTAP is linked to the tumor suppressor gene, p16lNK4A. Deficient levels of MTAP can occur in cancers primarily through co-deletion of the MTAP gene and the p16lNK4A gene. Cells expressing MTAP and possessing adenine salvage pathway activity may be less susceptible to malignancy due to growth-inhibitory actions of agents (e.g. antifolates), whose mechanism of action, in part, involves this de novo purine pathway.  Immunogen Recombinant human MTAP protein fragment (aa97-196) (exact sequence is	Isotype	IgG2b Kappa
Buffer 50mM Sodium Borate  Product Description  Host Mouse  Gene ID 4507  Gene Symbol MTAP  Species Human  Marker Tumor Suppressor Marker  Specificity/Sensitivity Recognizes a protein of 31kDa, which is identified as MTAP (5'-deoxy-5'-methylthioadenosine phosphorylase). It catalyzes the reversible phosphorolysis of methylthioadenosine, which is important in polyamine metabolism and for the salvage of adenine and methionine. The gene encoding MTAP is linked to the tumor suppressor gene, p16lNK4A. Deficient levels of MTAP can occur in cancers primarily through co-deletion of the MTAP gene and the p16lNK4A gene. Cells expressing MTAP and possessing adenine salvage pathway activity may be less susceptible to malignancy due to growth-inhibitory actions of agents (e.g. antifolates), whose mechanism of action, in part, involves this de novo purine pathway.  Immunogen Recombinant human MTAP protein fragment (aa97-196) (exact sequence is	Conjugate	Alexa Fluor 488
Product Description  Host Mouse  Gene ID 4507  Gene Symbol MTAP  Species Human  Marker Tumor Suppressor Marker  Specificity/Sensitivity Recognizes a protein of 31kDa, which is identified as MTAP (5'-deoxy-5'-methylthioadenosine phosphorylase). It catalyzes the reversible phosphorolysis of methylthioadenosine, which is important in polyamine metabolism and for the salvage of adenine and methionine. The gene encoding MTAP is linked to the tumor suppressor gene, p16lNK4A. Deficient levels of MTAP can occur in cancers primarily through co-deletion of the MTAP gene and the p16lNK4A gene. Cells expressing MTAP and possessing adenine salvage pathway activity may be less susceptible to malignancy due to growth-inhibitory actions of agents (e.g. antifolates), whose mechanism of action, in part, involves this de novo purine pathway.  Immunogen Recombinant human MTAP protein fragment (aa97-196) (exact sequence is	Purity	Protein A or G purified
Host Gene ID  4507  Gene Symbol  MTAP  Species  Human  Tumor Suppressor Marker  Specificity/Sensitivity  Recognizes a protein of 31kDa, which is identified as MTAP (5'-deoxy-5'-methylthioadenosine phosphorylase). It catalyzes the reversible phosphorolysis of methylthioadenosine, which is important in polyamine metabolism and for the salvage of adenine and methionine. The gene encoding MTAP is linked to the tumor suppressor gene, p16INK4A. Deficient levels of MTAP can occur in cancers primarily through co-deletion of the MTAP gene and the p16INK4A gene. Cells expressing MTAP and possessing adenine salvage pathway activity may be less susceptible to malignancy due to growth-inhibitory actions of agents (e.g. antifolates), whose mechanism of action, in part, involves this de novo purine pathway.  Immunogen  Recombinant human MTAP protein fragment (aa97-196) (exact sequence is	Buffer	50mM Sodium Borate
Gene Symbol  MTAP  Species  Human  Marker  Tumor Suppressor Marker  Specificity/Sensitivity  Recognizes a protein of 31kDa, which is identified as MTAP (5'-deoxy-5'-methylthioadenosine phosphorylase). It catalyzes the reversible phosphorolysis of methylthioadenosine, which is important in polyamine metabolism and for the salvage of adenine and methionine. The gene encoding MTAP is linked to the tumor suppressor gene, p16INK4A. Deficient levels of MTAP can occur in cancers primarily through co-deletion of the MTAP gene and the p16INK4A gene. Cells expressing MTAP and possessing adenine salvage pathway activity may be less susceptible to malignancy due to growth-inhibitory actions of agents (e.g. antifolates), whose mechanism of action, in part, involves this de novo purine pathway.  Immunogen  Recombinant human MTAP protein fragment (aa97-196) (exact sequence is	Product Description	
Gene Symbol  MTAP  Species  Human  Tumor Suppressor Marker  Specificity/Sensitivity  Recognizes a protein of 31kDa, which is identified as MTAP (5'-deoxy-5'-methylthioadenosine phosphorylase). It catalyzes the reversible phosphorolysis of methylthioadenosine, which is important in polyamine metabolism and for the salvage of adenine and methionine. The gene encoding MTAP is linked to the tumor suppressor gene, p16INK4A. Deficient levels of MTAP can occur in cancers primarily through co-deletion of the MTAP gene and the p16INK4A gene. Cells expressing MTAP and possessing adenine salvage pathway activity may be less susceptible to malignancy due to growth-inhibitory actions of agents (e.g. antifolates), whose mechanism of action, in part, involves this de novo purine pathway.  Immunogen  Recombinant human MTAP protein fragment (aa97-196) (exact sequence is	Host	Mouse
Specificity/Sensitivity   Tumor Suppressor Marker	Gene ID	4507
Tumor Suppressor Marker  Recognizes a protein of 31kDa, which is identified as MTAP (5'-deoxy-5'-methylthioadenosine phosphorylase). It catalyzes the reversible phosphorolysis of methylthioadenosine, which is important in polyamine metabolism and for the salvage of adenine and methionine. The gene encoding MTAP is linked to the tumor suppressor gene, p16INK4A. Deficient levels of MTAP can occur in cancers primarily through co-deletion of the MTAP gene and the p16INK4A gene. Cells expressing MTAP and possessing adenine salvage pathway activity may be less susceptible to malignancy due to growth-inhibitory actions of agents (e.g. antifolates), whose mechanism of action, in part, involves this de novo purine pathway.  Immunogen  Recombinant human MTAP protein fragment (aa97-196) (exact sequence is	Gene Symbol	MTAP
Recognizes a protein of 31kDa, which is identified as MTAP (5'-deoxy-5'-methylthioadenosine phosphorylase). It catalyzes the reversible phosphorolysis of methylthioadenosine, which is important in polyamine metabolism and for the salvage of adenine and methionine. The gene encoding MTAP is linked to the tumor suppressor gene, p16INK4A. Deficient levels of MTAP can occur in cancers primarily through co-deletion of the MTAP gene and the p16INK4A gene. Cells expressing MTAP and possessing adenine salvage pathway activity may be less susceptible to malignancy due to growth-inhibitory actions of agents (e.g. antifolates), whose mechanism of action, in part, involves this de novo purine pathway.  Immunogen  Recombinant human MTAP protein fragment (aa97-196) (exact sequence is	Species	Human
methylthioadenosine phosphorylase). It catalyzes the reversible phosphorolysis of methylthioadenosine, which is important in polyamine metabolism and for the salvage of adenine and methionine. The gene encoding MTAP is linked to the tumor suppressor gene, p16lNK4A. Deficient levels of MTAP can occur in cancers primarily through co-deletion of the MTAP gene and the p16lNK4A gene. Cells expressing MTAP and possessing adenine salvage pathway activity may be less susceptible to malignancy due to growth-inhibitory actions of agents (e.g. antifolates), whose mechanism of action, in part, involves this de novo purine pathway.  Immunogen  Recombinant human MTAP protein fragment (aa97-196) (exact sequence is	Marker	Tumor Suppressor Marker
		methylthioadenosine phosphorylase). It catalyzes the reversible phosphorolysis of methylthioadenosine, which is important in polyamine metabolism and for the salvage of adenine and methionine. The gene encoding MTAP is linked to the tumor suppressor gene, p16INK4A. Deficient levels of MTAP can occur in cancers primarily through co-deletion of the MTAP gene and the p16INK4A gene. Cells expressing MTAP and possessing adenine salvage pathway activity may be less susceptible to malignancy due to growth-inhibitory actions of agents (e.g. antifolates), whose mechanism of action, in part, involves this de novo purine pathway.
	Immunogen	



Page 2 of 3 v.20.1 Updated 10/26/2023	
Notes	Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.
Product Application Details	
Applications	Western Blot, ELISA, Immunohistochemistry, Immunohistochemistry-Paraffin, Protein Array
Recommended Dilutions	Western Blot, ELISA, Immunohistochemistry, Immunohistochemistry-Paraffin,

Optimal dilution of this antibody should be experimentally determined.

Protein Array

**Application Notes** 



### **Novus Biologicals USA**

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

## **Products Related to NBP2-75731AF488**

NBP1-43317AF488 Mouse IgG2b Kappa Light Chain Isotype Control (MG2b) [Alexa Fluor®

488]

NBP2-56796PEP MTAP Recombinant Protein Antigen

8499-IF-010/CF IFN-beta [Unconjugated] 10379-MT-050 MTAP [Unconjugated]

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-75731AF488

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

