Product Datasheet

Histone H1.2 Antibody (JU43-48) NBP2-75932

Unit Size: 100 ul

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 1

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-75932

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.


Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-75932




NBP2-75932

Histone H1.2 Antibody (JU43-48)


Thistoric TTT.2 / Willbody (0040 4	
Product Information	
Unit Size	100 ul
Concentration	1 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	JU43-48
Preservative	0.05% Sodium Azide
Isotype	IgG
Purity	Protein A purified
Buffer	TBS (pH7.4), 0.05% BSA, 40% Glycerol
Target Molecular Weight	21 kDa
Product Description	
Host	Rabbit
Gene ID	3006
Gene Symbol	H1-2
Species	Human, Mouse, Rat
lmmunogen	Synthetic peptide within Human Histone H1.2 aa 1-50 / 213. (SwissProt: P16403 Human; SwissProt: P15864 Mouse; SwissProt: P15865 Rat)
Product Application Details	
Applications	Western Blot, Chromatin Immunoprecipitation, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin
Recommended Dilutions	Western Blot 1:500-1:2000, Chromatin Immunoprecipitation 1:10-1:500, Immunohistochemistry 1:10 - 1:500, Immunocytochemistry/ Immunofluorescence 1:50-1:100, Immunohistochemistry-Paraffin 1:50-1:200
Images	
	tibody (JU43-48) [NBP2-75932] - analysis d Mouse lung tissue lysates using anti-




Immunocytochemistry/Immunofluorescence: Histone H1.2 Antibody (JU43-48) [NBP2-75932] - staining Histone H1.2 in SK-Br-3 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.




Immunohistochemistry-Paraffin: Histone H1.2 Antibody (JU43-48) [NBP2 -75932] - Immunohistochemical analysis of paraffin-embedded human lung carcinoma tissue using anti-Histone H1.2 antibody. The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 8.0-8.4) for 20 minutes. The tissues were blocked in 5% BSA for 30 minutes at room temperature, washed with ddH2O and PBS, and then probed with the primary antibody (1/50) for 30 minutes at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.




Western Blot: Histone H1.2 Antibody (JU43-48) [NBP2-75932] - analysis of Histone H1.2 on different cell lysates using anti-Histone H1.2 antibody at 1/500 dilution. Positive control: Lane 1: Hela Lane 2: 293 Lane 3: MCF-7




Immunocytochemistry/Immunofluorescence: Histone H1.2 Antibody (JU43-48) [NBP2-75932] - staining Histone H1.2 in Hela cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.




Immunocytochemistry/Immunofluorescence: Histone H1.2 Antibody (JU43-48) [NBP2-75932] - staining Histone H1.2 in PC-3M cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.




Immunohistochemistry-Paraffin: Histone H1.2 Antibody (JU43-48) [NBP2 -75932] - analysis of paraffin-embedded mouse colon tissue using anti-Histone H1.2 antibody. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6) for 20 mins. Counter stained with hematoxylin.



Immunohistochemistry-Paraffin: Histone H1.2 Antibody (JU43-48) [NBP2 -75932] - analysis of paraffin-embedded human kidney tissue using anti-Histone H1.2 antibody. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6) for 20 mins. Counter stained with hematoxylin.



Immunohistochemistry-Paraffin: Histone H1.2 Antibody (JU43-48) [NBP2-75932] - analysis of paraffin-embedded mouse colon tissue using anti-Histone H1.2 antibody. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6) for 20 mins. Counter stained with hematoxylin.



Publications

Wenliang Pan, Maria G Tsokos, Marc Scherlinger, Wei Li, George C Tsokos The PP2A regulatory subunit PPP2R2A controls NAD + biosynthesis to regulate T cell subset differentiation in systemic autoimmunity. Cell reports 2024-06-17 [PMID: 38889006]





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP2-75932

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

NBP2-49202PEP Histone H1.2 Recombinant Protein Antigen

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-75932

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

