Product Datasheet

Cytokeratin, pan Antibody (PAN-CK (Cocktail)) [Alexa Fluor® 488] NBP2-76425AF488

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-76425AF488

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP2-76425AF488



NBP2-76425AF488

Cytokeratin, pan Antibody (PAN-CK (Cocktail)) [Alexa Fluor® 488]

Unit Size 0.1 ml Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark. Clonality Monoclonal Clone PAN-CK (Cocktail) Preservative 0.05% Sodium Azide Isotype IgG Kappa Conjugate Alexa Fluor 488 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 3848 Gene Bymbol KRT1 Species Human, Mouse, Rat, Bovine, Canine, Chicken, Feline, Monkey, Rabbit Reactivity Notes Use in Mouse reported in scientific literature (PMID:34289354) Marker Epithelial Marker Specificity/Sensitivity Twenty human keratins are resolved with two-dimensional gel electrophoresis into acidic (p1 6:0) subfamilies. This antibody cocktail, which (CK1); 58ADa (CK4); 58ADa (CK5); 56KDa (CK7); 52KDa (CK7); 52KDa (CK7); 64KDa (CK1); 64KDa (CK1)	Cytokeratin, pan Antibody (PAN-0	CK (Cocktail)) [Alexa Fluor® 488]
Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark. Clonal Clone PAN-CK (Cocktail) Preservative 0.05% Sodium Azide Isotype IgG Kappa Conjugate Alexa Fluor 488 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 3848 Gene ID 3848 Gene Symbol KRT1 Species Human, Mouse, Rat, Bovine, Canine, Chicken, Feline, Monkey, Rabbit Reactivity Notes Use in Mouse reported in scientific literature (PMID:34289354) Marker Specificity/Sensitivity Twenty human keratins are resolved with two-dimensional gel electrophoresis into acidic (pl 6.0) subfamilies. This antibody cocktail recognizes acidic (Type I or LMW) and basic (Type II or HMW) cytokeratins, with 67kDa (Ck1); 64kDa (Ck3); 59kDa (Ck1); 58kDa (CK1); 58kDa (CK1); 58kDa (CK1); 58kDa (CK1); 48kDa	Product Information	
Storage Store at 4C in the dark. Clonality Monoclonal Clone PAN-CK (Cocktail) Preservative 0.05% Sodium Azide Isotype IgG Kappa Conjugate Alexa Fluor 488 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 3848 Gene Symbol KRT1 Species Human, Mouse, Rat, Bovine, Canine, Chicken, Feline, Monkey, Rabbit Reactivity Notes Use in Mouse reported in scientific literature (PMID:34289354) Marker Epithelial Marker Specificity/Sensitivity Twenty human keratins are resolved with two-dimensional gel electrophoresis into acidic (pl 6.0) subfamilies. This antibody cocktail recognizes acidic (Type I or LMW) and basic (Type II or HMW) cytokeratins, with 67kba (CK1); 64kba (CK3); 58kba (CK4); 58kba (CK6); 55kba (CK7); 52kba (CK3); 65kba (CK19); 46kba (CK19); 46kba (CK17); 45kba (CK18) and 40kba (CK19). Many studies have shown the usefulness of keratins as markers in cancer research and tumor diagnosis. KRT-PAN is a broad-spectrum anti pan-cytokeratin antibody cocktail, which differentiates epithelial tumors from non-epithelial mors e.g. syuamous vs. adenocarcinoma of the lung, liver carcinoma, breast cancer, and esophageal cancer. It is useful in characterizing the source of various neoplasms and to study the distribution of cytokeratin containing cells in epithelia during normal development and during the development of epithelial neoplasms. This antibody stains cytokeratins present in normal and abnormal human tissues eand shows high sensitivity in the recognition of epithelial cells and carcinomas.	Unit Size	0.1 ml
Clone PAN-CK (Cocktail) Preservative 0.05% Sodium Azide Isotype IgG Kappa Conjugate Alexa Fluor 488 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 3848 Gene Symbol KRT1 Species Human, Mouse, Rat, Bovine, Canine, Chicken, Feline, Monkey, Rabbit Reactivity Notes Use in Mouse reported in scientific literature (PMID:34289354) Marker Epithelial Marker Specificity/Sensitivity Twenty human keratins are resolved with two-dimensional gel electrophoresis into acidic (pl 6.0) subfamilies. This antibody cocktail recognizes acidic (Type I or LMW) and basic (Type I or HMW) cytokeratins, with 67kDa (CK1); 64kDa (CK3); 59kDa (CK4); 55kDa (CK7); 55kDa (CK7); 55kDa (CK1); 46kDa (CK10); 45kDa (CK11); 45kDa	Concentration	•
Clone PAN-CK (Cocktail) Preservative 0.05% Sodium Azide Isotype IgG Kappa Conjugate Alexa Fluor 488 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 3848 Gene Symbol KRT1 Species Human, Mouse, Rat, Bovine, Canine, Chicken, Feline, Monkey, Rabbit Reactivity Notes Use in Mouse reported in scientific literature (PMID:34289354) Marker Epithelial Marker Specificity/Sensitivity Twenty human keratins are resolved with two-dimensional gel electrophoresis into acidic (pl 6.0) subfamilies. This antibody cocktail recognizes acidic (Type I or LMW) and basic (Type II or HMW) cytokeratins, with 67kDa (CK1); 64kDa (CK1); 58kDa (CK1); 58kDa (CK1); 54kDa (CK1); 46kDa (CK17); 45kDa (CK18); 36bSa (CK19); 58kDa (CK19); Mary Studies have shown the usefulness of keratins as markers in cancer research and tumor diagnosis. KRT-PAN is a broad-spectrum anti pan-cytokeratin antibody cocktail, which differentiates epithelial tumors from non-epithelial tumors e.g. squamous vs. adenocarcinoma of the lung, liver carcinoma, breast cancer, and esophageal cancer. It is useful in characterizing the source of various neoplasms and to study the distribution of cytokeratins containing cells in epithelia during normal development and during the development of epithelial neoplasms. This antibody stains cytokeratins present in normal and abnormal human tissues and shows high sensitivity in the recognition of epithelial cells and carcinomas.	Storage	Store at 4C in the dark.
Preservative 0.05% Sodium Azide Isotype	Clonality	Monoclonal
IgG Kappa	Clone	PAN-CK (Cocktail)
Conjugate Alexa Fluor 488 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 3848 Gene Symbol KRT1 Species Human, Mouse, Rat, Bovine, Canine, Chicken, Feline, Monkey, Rabbit Reactivity Notes Use in Mouse reported in scientific literature (PMID:34289354) Marker Specificity/Sensitivity Twenty human keratins are resolved with two-dimensional gel electrophoresis into acidic (p1 6.0) subfamilies. This antibody cocktail recognizes acidic (Type I or LMW) and basic (Type II or HMW) cytokeratins, with 67kDa (CK1); 64kDa (CK3); 59kDa (CK4); 58kDa (CK13); 50kDa (CK13); 50kDa (CK13); 48kDa (CK16); 46kDa (CK17); 45kDa (CK13); 50kDa (CK13); 48kDa (CK16); 46kDa (CK17); 45kDa (CK18); 46kDa	Preservative	0.05% Sodium Azide
Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 3848 Gene Symbol KRT1 Species Human, Mouse, Rat, Bovine, Canine, Chicken, Feline, Monkey, Rabbit Reactivity Notes Use in Mouse reported in scientific literature (PMID:34289354) Marker Epithelial Marker Specificity/Sensitivity Twenty human keratins are resolved with two-dimensional gel electrophoresis into acidic (p1 6.0) subfamilies. This antibody cocktail recognizes acidic (Type I or LMW) cytokeratins, with 67kDa (CK1); 64kDa (CK3); 59kDa (CK4); 58kDa (CK13); 50kDa (CK14); 50kDa (CK15); 48kDa (CK16); 48kDa (CK16); 53kDa (CK17); 45kDa (CK18) and 40kDa (CK19). Many studies have shown the usefulness of keratins as markers in cancer research and tumor diagnosis. KRT-PAN is a broad-spectrum anti pan-cytokeratin antibody cocktail, which differentiates epithelial tumors from non-epithelial tumors e.g. squamous vs. adenocarcinoma of the lung, liver carcinoma, breast cancer, and esophageal cancer. It is useful in characterizing the source of various neoplasms and to study the distribution of cytokeratin containing cells in epithelia during normal development and during the development of epithelial neoplasms. This antibody stains cytokeratins present in normal and abnormal human tissues and shows high sensitivity in the recognition of epithelial cells and carcinomas.	Isotype	IgG Kappa
Buffer 50mM Sodium Borate	Conjugate	Alexa Fluor 488
Product Description Host Mouse Gene ID 3848 Gene Symbol KRT1 Species Human, Mouse, Rat, Bovine, Canine, Chicken, Feline, Monkey, Rabbit Reactivity Notes Use in Mouse reported in scientific literature (PMID:34289354) Marker Epithelial Marker Specificity/Sensitivity Twenty human keratins are resolved with two-dimensional gel electrophoresis into acidic (pl 6.0) subfamilies. This antibody cocktail recognizes acidic (Type I or LMW) and basic (Type II or HMW) cytokeratins, with 67kDa (CK1); 64kDa (CK3); 59kDa (CK4); 58kDa (CK5); 56kDa (CK6); 55kDa (CK7); 52kDa (CK8); 56.5kDa (CK10); 53kDa (CK13); 50kDa (CK14); 50kDa (CK15); 48kDa (CK16); 46kDa (CK17); 45kDa (CK18) and 40kDa (CK19). Many studies have shown the usefulness of keratins as markers in cancer research and tumor diagnosis. KRT-PAN is a broad-spectrum anti pan-cytokeratin antibody cocktail, which differentiates epithelial tumors from non-epithelial tumors e.g. squamous vs. adenocarcinoma of the lung, liver carcinoma, breast cancer, and esophageal cancer. It is useful in characterizing the source of various neoplasms and to study the distribution of cytokeratin containing cells in epithelia during normal development and during the development of epithelial neoplasms. This antibody stains cytokeratins present in normal and abnormal human tissues and shows high sensitivity in the recognition of epithelial cells and carcinomas.	Purity	Protein A or G purified
Mouse	Buffer	50mM Sodium Borate
Gene Symbol KRT1 Species Human, Mouse, Rat, Bovine, Canine, Chicken, Feline, Monkey, Rabbit Reactivity Notes Use in Mouse reported in scientific literature (PMID:34289354) Marker Epithelial Marker Specificity/Sensitivity Twenty human keratins are resolved with two-dimensional gel electrophoresis into acidic (pl 6.0) subfamilies. This antibody cocktail recognizes acidic (Type I or LMW) and basic (Type II or HMW) cytokeratins, with 67kDa (CK1); 64kDa (CK3); 59kDa (CK4); 58kDa (CK5); 56kDa (CK6); 55kDa (CK7); 52kDa (CK8); 56.5kDa (CK10); 53kDa (CK10); 53kDa (CK13); 50kDa (CK15); 48kDa (CK16); 46kDa (CK17); 45kDa (CK18) and 40kDa (CK19). Many studies have shown the usefulness of keratins as markers in cancer research and tumor diagnosis. KRT-PAN is a broad-spectrum anti pan-cytokeratin antibody cocktail, which differentiates epithelial tumors from non-epithelial tumors e.g. squamous vs. adenocarcinoma of the lung, liver carcinoma, breast cancer, and esophageal cancer. It is useful in characterizing the source of various neoplasms and to study the distribution of cytokeratin containing cells in epithelia during normal development and during the development of epithelial neoplasms. This antibody stains cytokeratins present in normal and abnormal human tissues and shows high sensitivity in the recognition of epithelial cells and carcinomas.	Product Description	
Gene Symbol KRT1 Species Human, Mouse, Rat, Bovine, Canine, Chicken, Feline, Monkey, Rabbit Reactivity Notes Use in Mouse reported in scientific literature (PMID:34289354) Marker Epithelial Marker Twenty human keratins are resolved with two-dimensional gel electrophoresis into acidic (pl 6.0) subfamilies. This antibody cocktail recognizes acidic (Type I or LMW) and basic (Type II or HMW) cytokeratins, with 67kDa (CK1); 64kDa (CK3); 59kDa (CK4); 58kDa (CK5); 56kDa (CK6); 55kDa (CK7); 52kDa (CK10); 53kDa (CK13); 50kDa (CK14); 50kDa (CK15); 48kDa (CK16); 46kDa (CK17); 45kDa (CK18) and 40kDa (CK19). Many studies have shown the usefulness of keratins as markers in cancer research and tumor diagnosis. KRT-PAN is a broad-spectrum anti pan-cytokeratin antibody cocktail, which differentiates epithelial tumors from non-epithelial tumors e.g. squamous vs. adenocarcinoma of the lung, liver carcinoma, breast cancer, and esophageal cancer. It is useful in characterizing the source of various neoplasms and to study the distribution of cytokeratin containing cells in epithelia during normal development and during the development of epithelial neoplasms. This antibody stains cytokeratins present in normal and abnormal human tissues and shows high sensitivity in the recognition of epithelial cells and carcinomas.	Host	Mouse
Species	Gene ID	3848
Reactivity Notes Use in Mouse reported in scientific literature (PMID:34289354)	Gene Symbol	KRT1
Marker Specificity/Sensitivity Twenty human keratins are resolved with two-dimensional gel electrophoresis into acidic (pl 6.0) subfamilies. This antibody cocktail recognizes acidic (Type I or LMW) and basic (Type II or HMW) cytokeratins, with 67kDa (CK1); 64kDa (CK3); 59kDa (CK4); 58kDa (CK5); 55kDa (CK6); 55kDa (CK7); 52kDa (CK8); 56.5kDa (CK10); 53kDa (CK13); 50kDa (CK14); 50kDa (CK15); 48kDa (CK16); 46kDa (CK17); 45kDa (CK18) and 40kDa (CK19). Many studies have shown the usefulness of keratins as markers in cancer research and tumor diagnosis. KRT-PAN is a broad-spectrum anti pan-cytokeratin antibody cocktail, which differentiates epithelial tumors from non-epithelial tumors e.g. squamous vs. adenocarcinoma of the lung, liver carcinoma, breast cancer, and esophageal cancer. It is useful in characterizing the source of various neoplasms and to study the distribution of cytokeratin containing cells in epithelia during normal development and during the development of epithelial neoplasms. This antibody stains cytokeratins present in normal and abnormal human tissues and shows high sensitivity in the recognition of epithelial cells and carcinomas.	Species	Human, Mouse, Rat, Bovine, Canine, Chicken, Feline, Monkey, Rabbit
Twenty human keratins are resolved with two-dimensional gel electrophoresis into acidic (pl 6.0) subfamilies. This antibody cocktail recognizes acidic (Type I or LMW) and basic (Type II or HMW) cytokeratins, with 67kDa (CK1); 64kDa (CK3); 59kDa (CK4); 58kDa (CK5); 56kDa (CK6); 55kDa (CK7); 52kDa (CK8); 56.5kDa (CK10); 53kDa (CK13); 50kDa (CK14); 50kDa (CK15); 48kDa (CK16); 46kDa (CK17); 45kDa (CK18) and 40kDa (CK19). Many studies have shown the usefulness of keratins as markers in cancer research and tumor diagnosis. KRT-PAN is a broad-spectrum anti pan-cytokeratin antibody cocktail, which differentiates epithelial tumors from non-epithelial tumors e.g. squamous vs. adenocarcinoma of the lung, liver carcinoma, breast cancer, and esophageal cancer. It is useful in characterizing the source of various neoplasms and to study the distribution of cytokeratin containing cells in epithelia during normal development and during the development of epithelial neoplasms. This antibody stains cytokeratins present in normal and abnormal human tissues and shows high sensitivity in the recognition of epithelial cells and carcinomas.	Reactivity Notes	Use in Mouse reported in scientific literature (PMID:34289354)
into acidic (pl 6.0) subfamilies. This antibody cocktail recognizes acidic (Type I or LMW) and basic (Type II or HMW) cytokeratins, with 67kDa (CK1); 64kDa (CK3); 59kDa (CK4); 58kDa (CK5); 56kDa (CK6); 55kDa (CK7); 52kDa (CK8); 56.5kDa (CK10); 53kDa (CK13); 50kDa (CK14); 50kDa (CK15); 48kDa (CK16); 46kDa (CK17); 45kDa (CK18) and 40kDa (CK19). Many studies have shown the usefulness of keratins as markers in cancer research and tumor diagnosis. KRT-PAN is a broad-spectrum anti pan-cytokeratin antibody cocktail, which differentiates epithelial tumors from non-epithelial tumors e.g. squamous vs. adenocarcinoma of the lung, liver carcinoma, breast cancer, and esophageal cancer. It is useful in characterizing the source of various neoplasms and to study the distribution of cytokeratin containing cells in epithelia during normal development and during the development of epithelial neoplasms. This antibody stains cytokeratins present in normal and abnormal human tissues and shows high sensitivity in the recognition of epithelial cells and carcinomas.	Marker	Epithelial Marker
Immunogen Human epidermal keratins	Specificity/Sensitivity	into acidic (pl 6.0) subfamilies. This antibody cocktail recognizes acidic (Type I or LMW) and basic (Type II or HMW) cytokeratins, with 67kDa (CK1); 64kDa (CK3); 59kDa (CK4); 58kDa (CK5); 56kDa (CK6); 55kDa (CK7); 52kDa (CK8); 56.5kDa (CK10); 53kDa (CK13); 50kDa (CK14); 50kDa (CK15); 48kDa (CK16); 46kDa (CK17); 45kDa (CK18) and 40kDa (CK19). Many studies have shown the usefulness of keratins as markers in cancer research and tumor diagnosis. KRT-PAN is a broad-spectrum anti pan-cytokeratin antibody cocktail, which differentiates epithelial tumors from non-epithelial tumors e.g. squamous vs. adenocarcinoma of the lung, liver carcinoma, breast cancer, and esophageal cancer. It is useful in characterizing the source of various neoplasms and to study the distribution of cytokeratin containing cells in epithelia during normal development and during the development of epithelial neoplasms. This antibody stains cytokeratins present in normal and abnormal human tissues and shows
	Immunogen	Human epidermal keratins




Notes

Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.

Product Application Details	
Applications	Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Frozen, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence
Recommended Dilutions	Western Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, Immunohistochemistry- Frozen, Immunofluorescence, CyTOF-ready
Application Notes	Optimal dilution of this antibody should be experimentally determined.

Images

Immunohistochemistry-Frozen: Mouse Monoclonal Cytokeratin, pan Antibody (PAN-CK (Cocktail)) [Alexa Fluor® 488] [NBP2-76425AF488] - MDA-MB-231 tumor cells (human breast tumor cell line) have been injected intracranially in immunodeficient mouse pups to form tumor. Sections have been stained with Pan-CK-A488 (green) (Catalog # NBP2 -76425AF488) and Hoechst (blue). Image from a verified customer review.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP2-76425AF488

DNST0 Endostatin [HRP]

MAB1455 Albumin Antibody (188835) [Unconjugated] - Serum

NB100-687 Cytokeratin 19 Antibody - BSA Free NBP2-16094 Cytokeratin 8 Antibody - BSA Free

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-76425AF488

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

