

# Product Datasheet

## Peripheral Node Addressin Antibody (MECA-79R) - BSA Free NBP2-78792

Unit Size: 0.5 mg

Store at -20 °C.

[www.novusbio.com](http://www.novusbio.com)



[technical@novusbio.com](mailto:technical@novusbio.com)

**Publications: 26**

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:  
[www.novusbio.com/NBP2-78792](http://www.novusbio.com/NBP2-78792)

Updated 1/29/2025 v.20.1

Earn rewards for product  
reviews and publications.

Submit a publication at [www.novusbio.com/publications](http://www.novusbio.com/publications)

Submit a review at [www.novusbio.com/reviews/destination/NBP2-78792](http://www.novusbio.com/reviews/destination/NBP2-78792)



**NBP2-78792**

Peripheral Node Addressin Antibody (MECA-79R) - BSA Free

**Product Information**

<b>Unit Size</b>	0.5 mg
<b>Concentration</b>	1.0 mg/ml
<b>Storage</b>	Store at -20 °C.
<b>Clonality</b>	Monoclonal
<b>Clone</b>	MECA-79R
<b>Preservative</b>	0.02% Sodium Azide
<b>Isotype</b>	IgG1
<b>Purity</b>	Protein A or G purified
<b>Buffer</b>	PBS

**Product Description**


<b>Description</b>	Clone MECA-79R (rat IgG1) is a recombinant version of the original clone MECA-79 (rat IgM).
<b>Host</b>	Rat
<b>Species</b>	Human, Mouse
<b>Immunogen</b>	Mouse lymph node stromal cells

**Product Application Details**


<b>Applications</b>	Western Blot, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Frozen, Immunohistochemistry-Paraffin, In vivo assay, Multiplex Immunofluorescence
<b>Recommended Dilutions</b>	Western Blot 1:100-1:2000, Immunohistochemistry 1:100-1:500, Immunocytochemistry/ Immunofluorescence reported in scientific literature (PMID 31278331), Immunohistochemistry-Paraffin 1:100-1:500, Immunohistochemistry-Frozen 1:100-1:500, In vivo assay reported in scientific literature (PMID 30277476), Multiplex Immunofluorescence 1:50
<b>Application Notes</b>	Additional reported application of in vitro and in vivo blocking of cell adhesion.

**Images**


Peripheral Node Addressin was detected in immersion fixed paraffin-embedded sections of human Tonsil using Rat Anti-Human Peripheral Node Addressin (MECA-79R), Monoclonal Antibody (Catalog #NBP2-78792) at 1:50 dilution at 37° Celsius for 4 minutes. Before incubation with the primary antibody, tissue underwent an all-in-one dewaxing and antigen retrieval preprocessing using PreTreatment Module (PT Module) and Dewax and HIER Buffer H (pH 9; Eprelia Catalog # TA-999-DHBH). Tissue was stained using the Alexa Fluor™ 647 Goat anti-Rat IgG Secondary Antibody at 1:200 at 37 ° Celsius for 2 minutes. (Yellow; Lunaphore Catalog # [DR647RT](#)) and counterstained with DAPI (blue; Lunaphore Catalog # [DR100](#)). Specific staining was localized to endothelial cells. Protocol available in [COMET™ Panel Builder](#).




**Western Blot: Peripheral Node Addressin Antibody (MECA-79R) [NBP2-78792]** - Total protein from human Tonsil, Lymph node, Spleen and mouse Spleen was separated on a 7.5% gel by SDS-PAGE, transferred to PVDF membrane and blocked in 5% non-fat milk in TBST. The membrane was probed with 2.0 ug/ml anti-PNAd in blocking buffer and detected with an anti-rat HRP secondary antibody using West Pico PLUS chemiluminescence detection reagent.




**Immunohistochemistry-Paraffin: Peripheral Node Addressin Antibody (MECA-79R) [NBP2-78792]** - Analysis of FFPE mouse adipose tissue section (with lymph node areas) using Peripheral Node Addressin antibody (clone MECA-79) at 1:100. The staining was developed with HRP-DAB detection method and the counterstaining was performed using hematoxylin. This Peripheral Node Addressin antibody generated a strong and specific staining of MECA-79 antigen in the the cytoplasm and the membranes of high endothelial venules (HEVs) aka peripheral lymph node addressin (PNAd) in lymph node areas of tested adipose tissue section.



**Immunohistochemistry-Paraffin: Peripheral Node Addressin Antibody (MECA-79R) [NBP2-78792]** - Analysis of FFPE human tonsil tissue section using Peripheral Node Addressin antibody (clone MECA-79R) at 1:100. The staining was developed with HRP-DAB detection method and the counterstaining was performed using hematoxylin.



**Immunohistochemistry: Peripheral Node Addressin Antibody (MECA-79R) [NBP2-78792]** - Local release of anti-IL-6 from IMB suppresses LN fibrosis. The skin allografts were harvested at 7 days post-transplantation. Lymphatic vessel expansion (Lyve-1+) and HEV elongation (MECA79+) were similar between all groups. Dense staining of collagen I and PDPN was seen in DLNs harvested from control and GelMA group compared to those from GelMA/anti-IL-6 group. (representative images from 4 different mice per group). Image collected and cropped by CiteAb from the following publication (<https://www.nature.com/articles/s41598-019-42349-w>), licensed under a CC-BY license.



## Publications

Zhao J, Jung S, Li X et al. Delivery of costimulatory blockade to lymph nodes promotes transplant acceptance in mice Journal of Clinical Investigation 2022-12-15 [PMID: 36519543]

Gavish A, Tyler M, Greenwald AC et al. Hallmarks of transcriptional intratumour heterogeneity across a thousand tumours Nature 2023-06-01 [PMID: 37258682]

Park JA, Espinosa-Cotton M, Guo HF et al. Targeting tumor vasculature to improve antitumor activity of T cells armed ex vivo with T cell engaging bispecific antibody Journal for immunotherapy of cancer 2023-03-01 [PMID: 36990507] (IHC, Mouse)

Quigley LT, Pang L, Tavanche E et al. Protocol for investigating tertiary lymphoid structures in human and murine fixed tissue sections using Opal<sup>®</sup>-TSA multiplex immunohistochemistry STAR protocols 2023-01-10 [PMID: 36633948] (IHC-P, Human)

### Details:

Dilution used in IHC-P 1:100

Smith KJ, Minns D, McHugh BJ Et al. The antimicrobial peptide cathelicidin drives development of experimental autoimmune encephalomyelitis in mice by affecting Th17 differentiation PLoS Biol 2022-08-26 [PMID: 36026478]

### Details:

Citation using the Non-Recombinant Monoclonal version of this antibody.

Li Y, Amaladas N, O'Mahony M et al. Treatment with a VEGFR-2 antibody results in intra-tumor immune modulation and enhances anti-tumor efficacy of PD-L1 blockade in syngeneic murine tumor models PloS one 2022-07-18 [PMID: 35849586] (IHC-P, Mouse)

Pinkard, H, Baghdassarian, H Et al. Learned adaptive multiphoton illumination microscopy for large-scale immune response imaging. Nat Commun 2021-03-26 [PMID: 33772022]

Li X, Zhao J, Kasinath V et al. Lymph node fibroblastic reticular cells deposit fibrosis-associated collagen following organ transplantation J. Clin. Invest. 2020-06-29 [PMID: 32597832] (IHC-Fr, Human)

### Details:

Citation using the Non-Recombinant Monoclonal format of this antibody.

Helmink BA, Reddy SM, Gao J et al. B cells and tertiary lymphoid structures promote immunotherapy response Nature. [PMID: 31942075] (MI, Human)

He B, Johansson-Percival A, Backhouse J et al. Remodeling of Metastatic Vasculature Reduces Lung Colonization and Sensitizes Overt Metastases to Immunotherapy Cell Rep. [PMID: 31968248] (ICC/IF, Mouse, Human)

Matsubara S, Seki M, Suzuki S et al. Tertiary lymphoid organs in the inflammatory myopathy associated with PD-1 inhibitors J Immunother Cancer. 2019-09-18 [PMID: 31533865] (IF/IHC, Human)

### Details:

Citation used the Non-Recombinant Monoclonal format of this antibody.

Berg EL, Robinson MK, Warnock RA et al The human peripheral lymph node vascular addressin is a ligand for LECAM-1, the peripheral lymph node homing receptor. J Cell Biol. 1991-07-01 [PMID: 1712790]

### Details:

Citation using the Non-Recombinant Monoclonal version of this antibody.

More publications at <http://www.novusbio.com/NBP2-78792>





### **Novus Biologicals USA**

10730 E. Briarwood Avenue  
Centennial, CO 80112  
USA  
Phone: 303.730.1950  
Toll Free: 1.888.506.6887  
Fax: 303.730.1966  
nb-customerservice@bio-techne.com

### **Bio-Techne Canada**

21 Canmotor Ave  
Toronto, ON M8Z 4E6  
Canada  
Phone: 905.827.6400  
Toll Free: 855.668.8722  
Fax: 905.827.6402  
canada.inquires@bio-techne.com

### **Bio-Techne Ltd**

19 Barton Lane  
Abingdon Science Park  
Abingdon, OX14 3NB, United Kingdom  
Phone: (44) (0) 1235 529449  
Free Phone: 0800 37 34 15  
Fax: (44) (0) 1235 533420  
info.EMEA@bio-techne.com

### **General Contact Information**

www.novusbio.com  
Technical Support: nb-technical@bio-  
techne.com  
Orders: nb-customerservice@bio-techne.com  
General: novus@novusbio.com

### **Products Related to NBP2-78792**

---

HAF005	Goat anti-Rat IgG Secondary Antibody [HRP]
F0105B	Goat anti-Rat IgG Secondary Antibody [Phycoerythrin]
DDXCR01	Rat IgG1 Isotype Control

---

### **Limitations**

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit [www.novusbio.com/guarantee](http://www.novusbio.com/guarantee)

Earn gift cards/discounts by submitting a review: [www.novusbio.com/reviews/submit/NBP2-78792](http://www.novusbio.com/reviews/submit/NBP2-78792)

Earn gift cards/discounts by submitting a publication using this product:  
[www.novusbio.com/publications](http://www.novusbio.com/publications)

