Product Datasheet

IL-3R alpha/CD123 Antibody (IL3RA/2947R) NBP2-79717-100ug

Unit Size: 100 ug

Store at 4C.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-79717

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-79717



NBP2-79717-100ug

IL-3R alpha/CD123 Antibody (IL3RA/2947R)


Product Information Unit Size 100 ug Concentration 0.2 mg/ml Storage Store at 4C. Clonality Monoclonal Clone IL3RA/2947R Preservative 0.05% Sodium Azide Isotype IgG Purity Protein A or G purified Buffer 10 mM PBS with 0.05% BSA Target Molecular Weight 70 kDa Product Description 200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP2-79851) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Rabbit Gene ID Gene ID 3563 Geneesymbol IL3RA Species Human Marker Acute Myeloid Leukemia Marker Acute Myeloid Leukemia Marker Immunogen ELISA, Immunohistochemistry, Immunohistochemistry-Paraffin, Protein Array Recombinant fragment of human IL-3R alpha/CD123 protein (around aa 26-171) (b.25-0.5 ug/ml, Immunohistochemistry-Paraffin 0.25-0.5			
Concentration 0.2 mg/ml Storage Stor at 4C. Clonality Monoclonal Clone IL3RA/2947R Preservative 0.05% Sodium Azide Isotype IgG Purity Protein A or G purified Buffer 10 mM PBS with 0.05% BSA Target Molecular Weight 70 kDa Product Description 200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP2-79851) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Rabbit Gene ID 3563 Gene Symbol IL3RA Species Human Marker Acute Myeloid Leukemia Marker Immunogen Recombinant fragment of human IL-3R alpha/CD123 protein (around aa 26-171) (exact sequence is proprietary) (Uniprot: P26951) Product Application Details Applications ELISA, Immunohistochemistry 0.25-0.5 ug/ml for 30 minutes at RT. Staining of formalin-fixed it issues is enhanced by heating tissue sections in 10mM Trs with 1mM EDTA, PI 9.0 for 45 min at 95C followed by cooling at RT or 20 minutes.	Product Information		
Storage Store at 4C. Clonality Monoclonal Clone IL3RA/2947R Preservative 0.05% Sodium Azide Isotype IgG Purity Protein A or G purified Buffer 10 mM PBS with 0.05% BSA Target Molecular Weight 70 kDa Product Description 200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP2-79851) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Rabbit Gene ID 3563 Gene Symbol IL3RA Species Human Marker Acute Myeloid Leukemia Marker Immunogen Recombinant fragment of human IL-3R alpha/CD123 protein (around aa 26-171) (exact sequence is proprietary) (Uniprot: P26951) Product Application Details ELISA, Immunohistochemistry. Jamunohistochemistry-Paraffin, Protein Array Application Notes ELISA. For coating, order antibody without BSA). Immunohistochemistry-Paraffin 0.25-0.5 ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues is enhanced by heating tissue sections in 10mM Tis with 1mM EDTA, PI 9.0 for 45 min at 95C followed by cooling at RT for 20 minutes.	Unit Size	100 ug	
Clonality Monoclonal Clone IL3RA/2947R Preservative 0.05% Sodium Azide Isotype IgG Purity Protein A or G purified Buffer 10 mM PBS with 0.05% BSA Target Molecular Weight 70 kDa Product Description 200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP2-79851) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Rabbit Gene ID 3563 Gene Symbol IL3RA Species Human Marker Acute Myeloid Leukemia Marker Immunogen ELISA, Immunohistochemistry, Immunohistochemistry-Paraffin, Protein Array Recombinant fragment of human IL-3R alpha/CD123 protein (around aa 26-171) (exact sequence is proprietary) (Uniprot: P26951) Product Application Details ELISA, Immunohistochemistry, O.25-0.5 ug/ml, Immunohistochemistry-Paraffin 0.25-0.5 ug/ml, Protein Array Application Notes ELISA, Immunohistochemistry 0.25-0.5 ug/ml, for 30 minutes at RT. Staining of formalin-fixed tissues is enhanced by heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0 for 45 min at 95C followed by cooli	Concentration	0.2 mg/ml	
Clone IL3RA/2947R Preservative 0.05% Sodium Azide Isotype IgG Purity Protein A or G purified Buffer 10 mM PBS with 0.05% BSA Target Molecular Weight 70 kDa Product Description 200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP2-79851) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Rabbit Gene ID 3563 Gene Symbol IL3RA Species Human Marker Acute Myeloid Leukemia Marker Immunogen ELISA, Immunohistochemistry. Immunohistochemistry-Paraffin, Protein Array Recommended Dilutions ELISA, Immunohistochemistry 0.25-0.5 ug/ml, Immunohistochemistry-Paraffin Application Notes ELISA, For coating, order antibody without BSA). Immunohistochemistry.Paraffin for 30 minutes at RT. Staining of formalin-fixed issues is enhanced by heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0 for 45 min at 95C followed by cooling at RT for 20 minutes.	Storage	Store at 4C.	
Preservative 0.05% Sodium Azide Isotype IgG Purity Protein A or G purified Buffer 10 mM PBS with 0.05% BSA Target Molecular Weight 70 kDa Product Description 200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP2-79851) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Rabbit Gene ID 3563 Gene Symbol IL3RA Species Human Marker Acute Myeloid Leukemia Marker Immunogen ELISA, Immunohistochemistry. (Uniprot: P26951) Product Application Details ELISA, Immunohistochemistry 0.25-0.5 ug/ml, Immunohistochemistry-Paraffin, 0.25-0.5 ug/ml, Immunohistochemistry.Paraffin Application Notes ELISA: For coating, order artibody without BSA). Immunohistochemistry (Formalin-fixed): 0.25-0.5 ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues is enhanced by heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0 tor 45 min at 95C followed by cooling at RT for 20 minutes.	Clonality	Monoclonal	
tsotype IgG Purity Protein A or G purified Buffer 10 mM PBS with 0.05% BSA Target Molecular Weight 70 kDa Product Description 200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP2-79851) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Rabbit Gene ID 3563 Gene Symbol IL3RA Species Human Marker Acute Myeloid Leukemia Marker Immunogen ELISA, Immunohistochemistry (Uniprot: P26951) Product Application Details ELISA, Immunohistochemistry 0.25-0.5 ug/ml, Immunohistochemistry-Paraffin 0.25-0.5 ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues is enhanced by heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0 for 45 min at 95C followed by cooling at RT for 20 minutes.	Clone	IL3RA/2947R	
Purity Protein A or G purified Buffer 10 mM PBS with 0.05% BSA Target Molecular Weight 70 kDa Product Description 200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP2-79851) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Rabbit Gene ID 3563 Gene Symbol IL3RA Species Human Marker Acute Myeloid Leukemia Marker Immunogen Recombinant fragment of human IL-3R alpha/CD123 protein (around aa 26-171) (exact sequence is proprietary) (Uniprot: P26951) Product Application Details ELISA, Immunohistochemistry, Immunohistochemistry-Paraffin, Protein Array Recommended Dilutions ELISA, Immunohistochemistry 0.25-0.5 ug/ml, Immunohistochemistry-Paraffin 0.25-0.5 ug/ml, Protein Array Application Notes ELISA: For coating, order antibody without BSA). Immunohistochemistry (Formalin-fixed): 0.25-0.5 ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues is enhanced by heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0 for 45 min at 95C followed by cooling at RT for 20 minutes.	Preservative	0.05% Sodium Azide	
Buffer 10 mM PBS with 0.05% BSA Target Molecular Weight 70 kDa Product Description 200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP2-79851) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Rabbit Gene ID 3563 Gene Symbol IL3RA Species Human Marker Acute Myeloid Leukemia Marker Immunogen Recombinant fragment of human IL-3R alpha/CD123 protein (around aa 26-171) (exact sequence is proprietary) (Uniprot: P26951) Product Application Details ELISA, Immunohistochemistry. Immunohistochemistry-Paraffin, Protein Array Recommended Dilutions ELISA, Immunohistochemistry 0.25-0.5 ug/ml, Immunohistochemistry-Paraffin 0.25-0.5 ug/ml, Protein Array Application Notes ELISA: For coating, order antibody without BSA). Immunohistochemistry (Formalin-fixed tissues is enhanced by heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0 for 45 min at 95C followed by cooling at RT for 20 minutes.	Isotype	IgG	
Target Molecular Weight 70 kDa Product Description 200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP2-79851) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Rabbit Gene ID 3563 Gene Symbol IL3RA Species Human Marker Acute Myeloid Leukemia Marker Immunogen Recombinant fragment of human IL-3R alpha/CD123 protein (around aa 26-171) (exact sequence is proprietary) (Uniprot: P26951) Product Application Details ELISA, Immunohistochemistry, Immunohistochemistry-Paraffin, Protein Array Recommended Dilutions ELISA, Immunohistochemistry 0.25-0.5 ug/ml, Immunohistochemistry-Paraffin 0.25-0.5 ug/ml, Protein Array ELISA: For coating, order antibody without BSA). Immunohistochemistry (Formalin-fixed): 0.25-0.5 ug/ml for 30 minutes at RT. Staining of formalin-fixed itsues is enhanced by heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0 for 45 min at 95C followed by cooling at RT for 20 minutes.	Purity	Protein A or G purified	
Product Description 200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP2-79851) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. -80C. Host Rabbit Gene ID 3563 Gene Symbol IL3RA Species Human Marker Acute Myeloid Leukemia Marker Immunogen Recombinant fragment of human IL-3R alpha/CD123 protein (around aa 26-171) (exact sequence is proprietary) (Uniprot: P26951) Product Application Details ELISA, Immunohistochemistry, Immunohistochemistry-Paraffin, Protein Array Recommended Dilutions ELISA, Immunohistochemistry 0.25-0.5 ug/ml, Immunohistochemistry-Paraffin 0.25-0.5 ug/ml, Protein Array Application Notes ELISA: For coating, order antibody without BSA). Immunohistochemistry (Formalin-fixed): 0.25-0.5ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues is enhanced by heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0 for 45 min at 95C followed by cooling at RT for 20 minutes.	Buffer	10 mM PBS with 0.05% BSA	
Description200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP2-79851) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C.HostRabbitGene ID3563Gene SymbolIL3RASpeciesHumanMarkerAcute Myeloid Leukemia MarkerImmunogenRecombinant fragment of human IL-3R alpha/CD123 protein (around aa 26-171) (exact sequence is proprietary) (Uniprot: P26951)Product Application DetailsELISA, Immunohistochemistry 0.25-0.5 ug/ml, Immunohistochemistry-Paraffin 0.25-0.5 ug/ml, Protein ArrayRecommended DilutionsELISA, Icr coating, order antibody without BSA). Immunohistochemistry (Formalin-fixed): 0.25-0.5 ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues is enhanced by heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0 for 45 min at 95C followed by cooling at RT for 20 minutes.	Target Molecular Weight	70 kDa	
Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP2-79851) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C.HostRabbitGene ID3563Gene SymbolIL3RASpeciesHumanMarkerAcute Myeloid Leukemia MarkerImmunogenRecombinant fragment of human IL-3R alpha/CD123 protein (around aa 26-171) (exact sequence is proprietary) (Uniprot: P26951)Product Application DetailsELISA, Immunohistochemistry 0.25-0.5 ug/ml, Protein ArrayApplication NotesELISA, Immunohistochemistry 0.25-0.5 ug/ml, Immunohistochemistry-Paraffin 0.25-0.5 ug/ml, Protein ArrayApplication NotesELISA: For coating, order antibody without BSA). Immunohistochemistry (Formalin-fixed): 0.25-0.5 ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues is enhanced by heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0 for 45 min at 95C followed by cooling at RT for 20 minutes.	Product Description		
HostRabbitGene ID3563Gene SymbolIL3RASpeciesHumanMarkerAcute Myeloid Leukemia MarkerImmunogenRecombinant fragment of human IL-3R alpha/CD123 protein (around aa 26-171) (exact sequence is proprietary) (Uniprot: P26951)Product Application DetailsApplicationsELISA, Immunohistochemistry, Immunohistochemistry-Paraffin, Protein ArrayRecommended DilutionsELISA, Immunohistochemistry 0.25-0.5 ug/ml, Immunohistochemistry-Paraffin 0.25-0.5 ug/ml, Protein ArrayApplication NotesELISA: For coating, order antibody without BSA). Immunohistochemistry (Formalin-fixed): 0.25-0.5ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues is enhanced by heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0 for 45 min at 95C followed by cooling at RT for 20 minutes.	Description	Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP2-79851) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to	
Gene SymbolIL3RASpeciesHumanMarkerAcute Myeloid Leukemia MarkerImmunogenRecombinant fragment of human IL-3R alpha/CD123 protein (around aa 26-171) (exact sequence is proprietary) (Uniprot: P26951)Product Application DetailsELISA, Immunohistochemistry, Immunohistochemistry-Paraffin, Protein ArrayRecommended DilutionsELISA, Immunohistochemistry 0.25-0.5 ug/ml, Immunohistochemistry-Paraffin 0.25-0.5 ug/ml, Protein ArrayApplication NotesELISA: For coating, order antibody without BSA). Immunohistochemistry (Formalin-fixed): 0.25-0.5ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues is enhanced by heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0 for 45 min at 95C followed by cooling at RT for 20 minutes.	Host		
SpeciesHumanMarkerAcute Myeloid Leukemia MarkerImmunogenRecombinant fragment of human IL-3R alpha/CD123 protein (around aa 26-171) (exact sequence is proprietary) (Uniprot: P26951)Product Application DetailsELISA, Immunohistochemistry, Immunohistochemistry-Paraffin, Protein ArrayRecommended DilutionsELISA, Immunohistochemistry 0.25-0.5 ug/ml, Immunohistochemistry-Paraffin 0.25-0.5 ug/ml, Protein ArrayApplication NotesELISA: For coating, order antibody without BSA). Immunohistochemistry (Formalin-fixed): 0.25-0.5ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues is enhanced by heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0 for 45 min at 95C followed by cooling at RT for 20 minutes.	Gene ID	3563	
MarkerAcute Myeloid Leukemia MarkerImmunogenRecombinant fragment of human IL-3R alpha/CD123 protein (around aa 26-171) (exact sequence is proprietary) (Uniprot: P26951)Product Application DetailsELISA, Immunohistochemistry, Immunohistochemistry-Paraffin, Protein ArrayApplicationsELISA, Immunohistochemistry 0.25-0.5 ug/ml, Immunohistochemistry-Paraffin 0.25-0.5 ug/ml, Protein ArrayApplication NotesELISA: For coating, order antibody without BSA). Immunohistochemistry (Formalin-fixed): 0.25-0.5ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues is enhanced by heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0 for 45 min at 95C followed by cooling at RT for 20 minutes.	Gene Symbol	IL3RA	
ImmunogenRecombinant fragment of human IL-3R alpha/CD123 protein (around aa 26-171) (exact sequence is proprietary) (Uniprot: P26951)Product Application DetailsApplicationsELISA, Immunohistochemistry, Immunohistochemistry-Paraffin, Protein ArrayRecommended DilutionsELISA, Immunohistochemistry 0.25-0.5 ug/ml, Immunohistochemistry-Paraffin 0.25-0.5 ug/ml, Protein ArrayApplication NotesELISA: For coating, order antibody without BSA). Immunohistochemistry (Formalin-fixed): 0.25-0.5ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues is enhanced by heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0 for 45 min at 95C followed by cooling at RT for 20 minutes.	Species	Human	
(exact sequence is proprietary) (Uniprot: P26951)Product Application DetailsApplicationsELISA, Immunohistochemistry, Immunohistochemistry-Paraffin, Protein ArrayRecommended DilutionsELISA, Immunohistochemistry 0.25-0.5 ug/ml, Immunohistochemistry-Paraffin 0.25-0.5 ug/ml, Protein ArrayApplication NotesELISA: For coating, order antibody without BSA). Immunohistochemistry (Formalin-fixed): 0.25-0.5ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues is enhanced by heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0 for 45 min at 95C followed by cooling at RT for 20 minutes.	Marker	Acute Myeloid Leukemia Marker	
ApplicationsELISA, Immunohistochemistry, Immunohistochemistry-Paraffin, Protein ArrayRecommended DilutionsELISA, Immunohistochemistry 0.25-0.5 ug/ml, Immunohistochemistry-Paraffin 0.25-0.5 ug/ml, Protein ArrayApplication NotesELISA: For coating, order antibody without BSA). Immunohistochemistry (Formalin-fixed): 0.25-0.5ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues is enhanced by heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0 for 45 min at 95C followed by cooling at RT for 20 minutes.	Immunogen		
Recommended DilutionsELISA, Immunohistochemistry 0.25-0.5 ug/ml, Immunohistochemistry-Paraffin 0.25-0.5 ug/ml, Protein ArrayApplication NotesELISA: For coating, order antibody without BSA). Immunohistochemistry (Formalin-fixed): 0.25-0.5ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues is enhanced by heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0 for 45 min at 95C followed by cooling at RT for 20 minutes.	Product Application Details		
0.25-0.5 ug/ml, Protein Array Application Notes ELISA: For coating, order antibody without BSA). Immunohistochemistry (Formalin-fixed): 0.25-0.5ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues is enhanced by heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0 for 45 min at 95C followed by cooling at RT for 20 minutes.	Applications	ELISA, Immunohistochemistry, Immunohistochemistry-Paraffin, Protein Array	
Immunohistochemistry (Formalin-fixed): 0.25-0.5ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues is enhanced by heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0 for 45 min at 95C followed by cooling at RT for 20 minutes.	Recommended Dilutions		
	Application Notes	Immunohistochemistry (Formalin-fixed): 0.25-0.5ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues is enhanced by heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0 for 45 min at 95C followed by cooling at RT for 20 minutes.	





Images

Immunohistochemistry-Paraffin: IL-3R alpha/CD123 Antibody (IL3RA/2947R) [NBP2-79717] - Formalin-fixed, paraffin-embedded human Tonsil stained with IL-3R alpha/CD123 Antibody (IL3RA/2947R).



Protein Array: IL-3R alpha/CD123 Antibody (IL3RA/2947R) [NBP2-79717] - Analysis of Protein Array containing more than 19,000 fulllength human proteins using IL-3R alpha/CD123 Antibody (IL3RA/2947R). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (Monoclonal Antibody) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt (TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a Monoclonal Antibody to its intended target. A Monoclonal Antibody is considered to specific to its intended target, if the Monoclonal Antibody has an S-score of at least 2.5





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-79717-100ug

HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control
NBP3-18359-10ug	Recombinant Mouse IL-3R alpha/CD123 hIgG-His Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-79717

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

