

Product Datasheet

beta-Catenin Antibody (CTNNB1/2030R) - Azide and BSA Free NBP2-79888

Unit Size: 100 ug

Store at -20 to -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-79888

Updated 10/23/2024 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-79888



NBP2-79888


beta-Catenin Antibody (CTNNB1/2030R) - Azide and BSA Free

Product Information	
Unit Size	100 ug
Concentration	1 mg/ml
Storage	Store at -20 to -80C. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	CTNNB1/2030R
Preservative	No Preservative
Isotype	IgG Kappa
Purity	Protein A purified
Buffer	10 mM PBS
Target Molecular Weight	92 kDa
Product Description	
Description	1.0 mg/ml of antibody purified by Protein A Column. Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WITH BSA & azide (NBP2-79753). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80 C.
Host	Rabbit
Gene ID	1499
Gene Symbol	CTNNB1
Species	Human
Immunogen	Recombinant full-length human beta-Catenin protein (Uniprot: P35222)
Product Application Details	
Applications	Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Protein Array, CyTOF-ready
Recommended Dilutions	Western Blot 1-2 ug/ml, Flow Cytometry 1-2 ug/million cells, Immunohistochemistry 1-2 ug/ml, Immunocytochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry-Paraffin 1-2 ug/ml, Protein Array, CyTOF-ready
Application Notes	Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes. Optimal dilution for a specific application should be determined.




Images


Immunocytochemistry/Immunofluorescence: beta-Catenin Antibody (CTNNB1/2030R) - Azide and BSA Free [NBP2-79888] - Immunofluorescence Analysis of PFA fixed HeLa cells labeling Beta-Catenin with Beta-Catenin Recombinant Rabbit Monoclonal Ab (CTNNB1/2030R followed by Goat anti-Mouse IgG-CF488 (Green). The nuclear counterstain is Reddot (Red).



Immunohistochemistry-Paraffin: beta-Catenin Antibody (CTNNB1/2030R) - Azide and BSA Free [NBP2-79888] - Formalin-fixed, paraffin-embedded human Cervical Carcinoma stained with Beta-Catenin Recombinant Rabbit Monoclonal Antibody (CTNNB1/2030R).




Flow Cytometry: beta-Catenin Antibody (CTNNB1/2030R) - Azide and BSA Free [NBP2-79888] - Flow Cytometric Analysis of PFA fixed HeLa cells using Beta-Catenin Recombinant Rabbit Monoclonal Ab (CTNNB1/2030R) followed by Goat anti-Rabbit IgG-CF488 (Blue); Isotype Control (Red).



Immunohistochemistry-Paraffin: beta-Catenin Antibody (CTNNB1/2030R) - Azide and BSA Free [NBP2-79888] - Formalin-fixed, paraffin-embedded human Tonsil stained with Beta-Catenin Recombinant Rabbit Monoclonal Antibody (CTNNB1/2030R).



Protein Array: beta-Catenin Antibody (CTNNB1/2030R) - Azide and BSA Free [NBP2-79888] - Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-79888

HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-61628PEP	beta-Catenin Recombinant Protein Antigen
DYC1329-2	beta-Catenin [Biotin]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-79888

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

