

Product Datasheet

Hepatic Sinusoidal Endothelial Cells Antibody (SE-1) - Azide and BSA Free NBP2-80759

Unit Size: 0.1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-80759

Updated 8/21/2023 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-80759




NBP2-80759**Hepatic Sinusoidal Endothelial Cells Antibody (SE-1) - Azide and BSA Free**


Product Information	
Unit Size	0.1 ml
Concentration	1 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	SE-1
Preservative	No Preservative
Isotype	IgG2a Kappa
Purity	Protein A purified
Buffer	Tris-Glycine, 0.15 M NaCl
Product Description	
Host	Mouse
Species	Human, Mouse, Rat
Immunogen	Rat Hepatic Sinusoidal Endothelial Cells
Product Application Details	
Applications	Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Frozen, Immunohistochemistry-Paraffin
Recommended Dilutions	Western Blot 1-5 ug/ml, Flow Cytometry reported in scientific literature (Motoyama et al; PMID: 9428229), Immunohistochemistry 1-5 ug/ml, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin 1-5 ug/ml, Immunohistochemistry-Frozen 1-5 ug/ml
Application Notes	In WB, Hepatic Sinusoidal Endothelial Cells antibody (clone SE-1) generates a specific band around 45-50 kDa molecular weight position. For IHC use neutral buffered formalin fixated (perfusion fixation is recommended) paraffin embedded tissues after proteinase K treatment or acetone fixed frozen sections.

Images


Western Blot: Hepatic Sinusoidal Endothelial Cells Antibody (SE-1) - Azide and BSA Free [NBP2-80759] - Total protein from human, mouse and rat liver was separated on a 12% gel by SDS-PAGE, transferred to PVDF membrane and blocked in 5% non-fat milk in TBST. The membrane was probed with 2.0 ug/ml anti-H.S.E.C in 1% non-fat milk in TBST and detected with an anti-mouse HRP secondary antibody using chemiluminescence. Image from the standard format of this antibody.




Immunocytochemistry/Immunofluorescence: Hepatic Sinusoidal Endothelial Cells Antibody (SE-1) - Azide and BSA Free [NBP2-80759] - Release of structural liver cells into the perfusate after cold ischemia. Representative images of surface marker expression of hepatocytes (top), LSEC (middle), and stellate cells (below). Scale bars: 5 μ m. Image collected and cropped by CiteAb from the following publication (<https://www.nature.com/articles/s41598-020-57589-4>), licensed under a CC-BY license.




Immunohistochemistry-Frozen: Hepatic Sinusoidal Endothelial Cells Antibody (SE-1) - Azide and BSA Free [NBP2-80759] - Frozen rat liver tissue sections. DAPI (blue), SE1 (green), cytochrome P450 (red). Tissue sections were acetone fixed. SE1 antibody at 1:500, incubated at 4C overnight. Multiplexed with cytochrome P450 from another vendor. IHC-Fr image submitted by a veri



Flow Cytometry: Hepatic Sinusoidal Endothelial Cells Antibody (SE-1) - Azide and BSA Free [NBP2-80759] - Using the FITC direct conjugate Flow Cytometry: Surface staining of Rat Liver sinusoidal endothelial cells with Mouse anti-Rat Hepatic Sinusoidal Endothelial Cells [FITC] [NB110-68095F] and negative control. Total viable cells were used for analysis. Image courtesy of customer.



Immunohistochemistry: Hepatic Sinusoidal Endothelial Cells Antibody (SE-1) - Azide and BSA Free [NBP2-80759] - Analysis of frozen normal rat liver tissue sections using anti-Hepatic Sinusoidal Endothelial Cells antibody (green). Nuclei were counterstained with DAPI (blue). Image from verified customer review. Image from the standard format of this antibody.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-80759

HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB720-B	Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]
NBP1-96981-0.5mg	Mouse IgG2a Kappa Isotype Control (M2AK)

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-80759

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

