

Product Datasheet

MMP-9 Antibody (4A3) - Azide and BSA Free NBP2-80855

Unit Size: 0.1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Reviews: 1 Publications: 2


Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-80855

Updated 8/21/2023 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-80855



NBP2-80855

MMP-9 Antibody (4A3) - Azide and BSA Free


Product Information	
Unit Size	0.1 ml
Concentration	1 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	4A3
Preservative	No Preservative
Isotype	IgG1 Kappa
Purity	Protein G purified
Buffer	Tris-Glycine (pH 7.5), 0.15 M NaCl

Product Description	
Host	Mouse
Gene ID	4318
Gene Symbol	MMP9
Species	Human, Mouse, Rat
Reactivity Notes	Mouse reactivity reported in scientific literature (PMID: 30772382).
Immunogen	Synthetic peptide from the N-terminus of human MMP9 [Swiss-Prot# P14780]


Product Application Details	
Applications	Western Blot, ELISA, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready
Recommended Dilutions	Western Blot 1:500-1:1000, ELISA 1:100 - 1:2000, Immunohistochemistry 1:100-1:200, Immunohistochemistry-Paraffin 1:100-1:200, CyTOF-ready reported by customer review
Application Notes	In Western blot a band can be seen at 82 or 63 kDa, representing the active forms of MMP9. In IHC, cytoplasmic and extracellular staining was observed in HeLa cells. Prior to immunostaining paraffin tissues, antigen retrieval with sodium citrate buffer (pH 6.0) is recommended.

Images


Western Blot: MMP-9 Antibody (4A3) - Azide and BSA Free [NBP2-80855] - Identification of gelatinases by Western blot analysis. A pool of sera from 10 patients with inactive RR-MS was used. For 2-DZ, pooled sera were concentrated 10-fold and 40 ug of total proteins were applied (left panels). For Western blot analysis pooled sera were concentrated 40-fold and 160 ug of total proteins were subjected to 2-DE (right panels). 2-DE gels were cut and the corresponding nitrocellulose membranes were treated with anti-NGAL Ab (A'), anti-MMP-2 Ab (B') and Anti-MMP-9 (4A3) Ab, recognizing the 65 kD and the 82 kD activated forms (C'); Gel C show the region with isoelectric point between pH 4-5.5 and molecular mass between 85 and 55 kD, analysed by Western blot in C', respectively. Image collected and cropped by CiteAb from the following publication (null), licensed under a CC-BY license. Image from the standard format of this antibody.




Immunohistochemistry-Paraffin: MMP-9 Antibody (4A3) - Azide and BSA Free [NBP2-80855] - Analysis of a FFPE section of human esophageal cancer using MMP9 antibody (clone 4A3) at 1:200. The staining was developed with HRP-labelled secondary antibody and DAB reagent followed by hematoxylin counterstaining. This antibody clone generated an extra




Western Blot: MMP-9 Antibody (4A3) - Azide and BSA Free [NBP2-80855] - 1.5 ul of serum from healthy control (HC) or from a patient with inactive RR-MS loaded on a 7.5% polyacrylamide gel copolymerized with 0.1% (w/v) gelatin (1-DZ). 2 ul of fivefold concentrated (Vivaspin 500, MWCO 50,000; GE Healthcare) serum samples (the same as in 1-DZ) separated on a 7.5% polyacrylamide gel without gelatin (1-DE), transferred on nitrocellulose membranes & incubated with the following primary antibodies: Anti-NGAL (5G5) at 2.0 ug/ml; Anti-MMP-9 (Ab-8) (IA5) at 2.66 ug/ml; Anti-MMP-2 (Ab-4) (75-7F7) at 1.0 ug/ml & Anti-MMP-9 Ab (4A3) at 2.0 ug/ml. Image collected & cropped by CiteAb from the following publication (null), licensed under a CC-BY license. Image from the standard format of this antibody.




Western Blot: MMP-9 Antibody (4A3) - Azide and BSA Free [NBP2-80855] - Analysis of activated MMP9 expression in MMP9 proenzyme incubated with trypsin for various times. Image from the standard format of this antibody.




Immunohistochemistry: MMP-9 Antibody (4A3) - Azide and BSA Free [NBP2-80855] - MMP-9 (green) was detected in murine brain (cortex) using MMP-9-DyLight 488 (4A3) with a concentration of 1:20 in PBS for 2 hours. MMP9-Signals are overlapped with the corresponding phase contrast image. Image from verified customer review. Image using the



Immunohistochemistry-Paraffin: MMP-9 Antibody (4A3) - Azide and BSA Free [NBP2-80855] - Analysis of a FFPE tissue section of human esophageal cancer using MMP9 antibody (clone 4A3) at 1:200. The staining was developed with HRP-labelled secondary antibody and DAB reagent followed by hematoxylin counterstaining. This antibody generated a speci



CyTOF-ready: MMP-9 Antibody (4A3) - Azide and BSA Free [NBP2-80855] - Analysis of MMP-9 antibody on human tissue. MMP9 (red), nuclei (blue). Scale, 50um.



Publications

Radnaa E, Richardson L, Goldman B Et al. Stress signaler p38 mitogen-activated kinase activation: a cause for concern? Clin Sci (Lond) 2022-10-17 [PMID: 36250628] (CyTOF-ready, Human)

Details:

Citation using the Azide and BSA Free version of this antibody.

Gama Sosa MA, De Gasperi R, Pryor D et al. Low-level blast exposure induces chronic vascular remodeling, perivascular astrocytic degeneration and vascular-associated neuroinflammation Acta Neuropathol Commun 2021-10-15 [PMID: 34654480] (IF/IHC, Rat)

Details:

Citation using the Azide and BSA Free format of this antibody.



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-80855

HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB720-B	Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]
NBP1-43319-0.5mg	Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1)
NBP1-57940PEP	MMP-9 Antibody Blocking Peptide

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-80855

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

