

Product Datasheet

Fc gamma RI/CD64 Antibody (H22) [Biotin] NBP2-81098B

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-81098B

Updated 7/11/2023 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-81098B



NBP2-81098B

Fc gamma RI/CD64 Antibody (H22) [Biotin]

Product Information	
Unit Size	0.1 ml
Concentration	Please see the vial label for concentration. If unlisted please contact technical services.
Storage	Store at 4C in the dark.
Clonality	Monoclonal
Clone	H22
Preservative	0.05% Sodium Azide
Isotype	IgG Kappa
Conjugate	Biotin
Purity	Protein A purified
Buffer	PBS
Product Description	
Host	Rabbit
Gene ID	2209
Gene Symbol	FCGR1A
Species	Human
Specificity/Sensitivity	This immunotoxin is specific for human Fc gamma RI/CD64, and specifically an epitope outside the Fc binding domain. The binding activity of the VH/VL antibody format was unaffected by fusion of the m22(scFv) coding regions to the truncated ETA coding sequences
Immunogen	This recombinant antibody was prepared by cloning the single chain fragment (scFv) derived from the murine anti-human Fc gamma RI/CD64 mAb m22 into the bacterial expression vector pBM1.1, and fusing it to a deletion mutant of Pseudomonas exotoxin A.
Product Application Details	
Applications	Flow Cytometry, Immunocytochemistry/Immunofluorescence, Competitive ELISA
Recommended Dilutions	Flow Cytometry, Immunocytochemistry/Immunofluorescence, Competitive ELISA
Application Notes	Optimal dilution of this antibody should be experimentally determined. This antibody binds human CD64 and can be bound to a toxin so that together it displays specific cytotoxicity, and thus can be used to selectively eliminate AML cells; FC analysis indicated that this immunotoxin drove 41% of primary leukemia cells from a patient with CD64-positive AML into early apoptosis (Tur, 2003). Similarly, in an in vivo mouse model, IHC analysis revealed efficient elimination of human CD64+ tumor cells in mouse organs (Tur, 2011).





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-81098B

210-TA-005	TNF-alpha [Unconjugated]
1257-FC-050	Fc gamma RI/CD64 [Unconjugated]
D6050	IL-6 [HRP]
285-IF-100	IFN-gamma [Unconjugated]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-81098B

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

