## **Product Datasheet**

# CRISPR-Cas9 Antibody (6H4) - BSA Free NBP2-81124

Unit Size: 0.1 mg

Store at 4C for up to 3 months. For longer storage, aliquot and store at -20C.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-81124

Updated 9/11/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP2-81124



### NBP2-81124

CRISPR-Cas9 Antibody (6H4) - BSA Free


Product Information	
Unit Size	0.1 mg
Concentration	1.0 mg/ml
Storage	Store at 4C for up to 3 months. For longer storage, aliquot and store at -20C.
Clonality	Monoclonal
Clone	6H4
Preservative	0.02% Sodium Azide
Purity	Protein A or G purified
Buffer	PBS
Target Molecular Weight	158.4 kDa

Mouse
901176
Bacteria
This CRISPR-Cas9 antibody (6H4) was raised against partial recombinant protein made to an N terminal portion of the Staphylococcus aureus CRISPR/Cas9 protein
E

<b>Product Application Details</b>	
Applications	Western Blot, Immunocytochemistry/ Immunofluorescence, Immunoprecipitation
Recommended Dilutions	Western Blot 0.5 - 1.0 ug/ml, Immunocytochemistry/ Immunofluorescence 1:1000, Immunoprecipitation 1 ug / 200 ug lysate

## **Images**

Western Blot: CRISPR-Cas9 Antibody (6H4) [NBP2-81124] - Total protein from mock and Cas9 transfected 293 cells, HeLa and 3T3 cells was separated on a 7.5% gel by SDS-PAGE, transferred to PVDF membrane and blocked in 5% non-fat milk in TBST. The membrane was probed with 0.5 ug/ml anti-Cas9 in block buffer and detected with an anti-mouse HRP secondary antibody using West Pico PLUS chemiluminescence detection reagent. Theoretical molecular weight: 158.4 kDa.



#### **Procedures**

## Western Blot protocol for CRISPR-Cas9 Antibody (NBP2-81124)

CRISPR-Cas9 Antibody (6H4):

Western Blot Protocol

- 1. Perform SDS-PAGE on samples to be analyzed, loading 10-25 ug of total protein per lane.
- 2. Transfer proteins to PVDF membrane according to the instructions provided by the manufacturer of the membrane and transfer apparatus.
- 3. Stain the membrane with Ponceau S (or similar product) to assess transfer success, and mark molecular weight standards where appropriate.
- 4. Rinse the blot TBS -0.05% Tween 20 (TBST).
- 5. Block the membrane in 5% Non-fat milk in TBST (blocking buffer) for at least 1 hour.
- 6. Wash the membrane in TBST three times for 10 minutes each.
- 7. Dilute primary antibody in blocking buffer and incubate overnight at 4C with gentle rocking.
- 8. Wash the membrane in TBST three times for 10 minutes each.
- 9. Incubate the membrane in diluted HRP conjugated secondary antibody in blocking buffer (as per manufacturer's instructions) for 1 hour at room temperature.
- 10. Wash the blot in TBST three times for 10 minutes each (this step can be repeated as required to reduce background).
- 11. Apply the detection reagent of choice in accordance with the manufacturers instructions.





## Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

## **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

## **General Contact Information**

www.novusbio.com Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

## **Products Related to NBP2-81124**

HAF007 Goat anti-Mouse IgG Secondary Antibody [HRP]

NB720-B Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]

NBP2-52986 CRISPR-Cas9 Antibody Pack

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-81124

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

