Product Datasheet

SKIV2L Antibody - BSA Free NBP2-81903

Unit Size: 0.1 mg

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-81903

Updated 10/19/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-81903



NBP2-81903

SKIV2L Antibody - BSA Free

Product Information Unit Size 0.1 mg Concentration 1 mg/ml Storage Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles. Clonality Polyclonal Preservative 0.02% Sodium Azide Isotype IgG Purity Peptide affinity purified Buffer PBS Target Molecular Weight 135 kDa Product Description Kabbit Gene ID 6499 Gene Symbol SKIV2L Species Human, Mouse, Rat Specificity/Sensitivity At least two isoforms of SKIV2L are known to exist; this antibody will detect both isoforms. Immunogen SKIV2L antibody was raised against a 16 amino acid peptide near the amino terminus of human SKIV2L. The immunogen is located within amino acids 60 - 110 of SKIV2L amino Acid Squence: EHSARKWQRKTDWSL Product Application Details Western Blot, ELISA, Immunocytochemistry/ Immunofiluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin Recommended Dilutions Western Blot, 1-2 ug/ml, ELISA, Immunohistochemistry 5 ug/ml, Immunohistochemistry-Paraffin	-		
Concentration 1 mg/ml Storage Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles. Clonality Polyclonal Preservative 0.02% Sodium Azide Isotype IgG Purity Peptide affinity purified Buffer PBS Target Molecular Weight 135 kDa Product Description Host Host Rabbit Gene ID 6499 Gene Symbol SKIV2L Species Human, Mouse, Rat Specificity/Sensitivity At least two isoforms of SKIV2L are known to exist; this antibody will detect both isoforms. Immunogen SKIV2L antibody was raised against a 16 amino acid peptide near the amino terminus of human SKIV2L. The immunogen is located within amino acids 60 - 110 of SKIV2L. Amino Acid Squence: EHSARKWQRKTDPWSL Product Application Details Western Blot, ELISA, Immunocytochemistry/Immunofluorescence, Immunohistochemistry.Paraffin Recommended Dilutions Western Blot, 22 ug/ml, ELISA, Immunohistochemistry 5 ug/ml, Immunohistochemistry.	Product Information		
Storage Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles. Clonality Polyclonal Preservative 0.02% Sodium Azide Isotype IgG Purity Peptide affinity purified Buffer PBS Target Molecular Weight 135 kDa Product Description Host Rabbit Gene ID Gene ID 6499 Gene Symbol SKIV2L Species Human, Mouse, Rat Species Human, Mouse, Rat Specificity/Sensitivity At least two isoforms of SKIV2L are known to exist; this antibody will detect both isoforms. Immunogen SKIV2L. Antibody was raised against a 16 amino acid peptide near the amino terminus of human SKIV2L. The immunogen is located within amino acids 60 - 110 of SKIV2L. Amino Acid Squence: EHSARKWQRKTDPWSL Product Application Details Western Blot, ELISA, Immunocytochemistry/Immunofluorescence, Immunohistochemistry.Paraffin Recommended Dilutions Western Blot, 22 uy/ml, ELISA, Immunohistochemistry 5 ug/ml, Immunohistochemistry.	Unit Size	0.1 mg	
cycles.ClonalityPolyclonalPreservative0.02% Sodium AzideIsotypeIgGPurityPeptide affinity purifiedBufferPBSTarget Molecular Weight135 kDaProduct DescriptionRabbitHostRabbitGene ID6499Gene SymbolSKIV2LSpeciesHuman, Mouse, RatSpecificity/SensitivityAt least two isoforms of SKIV2L are known to exist; this antibody will detect both isoforms.ImmunogenSKIV2L. Amino Acid Squence: EHSARKWQRKTDPWSLProduct Application DetailsWestern Blot, ELISA, Immunocytochemistry-Immunofluorescence, Immunohistochemistry-Immunohistochemistry-Sug/ml, Immunohistochemistry-Sug/ml, Immunohistochemistry-Immunohistochemistry-Sug/ml, Immunohistochemistry-Immunohistochemistry Sug/ml, Immunohistochemistry/Immunohistochemistry-Sug/ml, Immunohistochemistry/Immunohistochemistry Sug/ml, Immunohistochemistry/Immunohistochemistry-Sug/ml, Immunohistochemistry-Sug/ml, Immunohistochemistry-Sug/ml, Immunohistochemistry-Immunohistochemistry-Sug/ml, Immunohistochemistry-Sug/ml, Immunohistochemistry-Immunohistochemistry-Sug/ml, Immunohistochemistry-Immunohistochemistry-Sug/ml, Immunohistochemistry-Immunohistochemistry-Sug/ml, Immunohistochemistry-Immunohistochemistry-Sug/ml, Immunohistochemistry-Immunohistochemistry-Sug/ml, Immunohistochemistry-Immunohistochemistry-Sug/ml, Immunohistochemistry-Immunohistochemistry-Sug/ml, Immunohistochemistry/Immunohistochemistry-Sug/ml, Immunohistochemistry-Immunohistochemistry-Sug/ml, Immunohistochemistry-Immunohistochemistry-Sug/ml, Immunohistochemistry-Immunohistochemistry-Immunohistochemistry-Sug/ml, Immunohistochemistry-Immunohistochemistry-Immunohistochemistry-Immunohistochemistry-Immunohistochemistry-Immunohistochemistry-Immunohist	Concentration	1 mg/ml	
Preservative0.02% Sodium AzideIsotypeIgGPurityPeptide affinity purifiedBufferPBSTarget Molecular Weight135 kDaProduct DescriptionKabbitHostRabbitGene ID6499Gene SymbolSKIV2LSpeciesHuman, Mouse, RatSpecificity/SensitivityAt least two isoforms of SKIV2L are known to exist; this antibody will detect both isoforms.ImmunogenSKIV2L antibody was raised against a 16 amino acid peptide near the amino terminus of human SKIV2L. The immunogen is located within amino acids 60 - 110 of SKIV2L. Amino Acid Squence: EHSARKWQRKTDPWSLProduct Application DetailsWestern Blot, ELISA, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunohistochemistry 5 ug/ml, Immunocytochemistry 5 ug/ml, Immunocytochemistry/Immunofluorescence 20 ug/ml, Immunohistochemistry-Immunofluorescence 20 ug/ml, Immunocytochemistry-Immunofluorescence 20 ug/ml, Immunohistochemistry-Immunofluorescence 20 ug/ml, Immunoflu	Storage		
IsotypeIgGPurityPeptide affinity purifiedBufferPBSTarget Molecular Weight135 kDaProduct DescriptionKabbitHostRabbitGene ID6499Gene SymbolSKIV2LSpeciesHuman, Mouse, RatSpecificity/SensitivityAt least two isoforms of SKIV2L are known to exist; this antibody will detect both isoforms.ImmunogenSKIV2L antibody was raised against a 16 amino acid peptide near the amino terminus of human SKIV2L. The immunogen is located within amino acids 60 - 110 of SKIV2L. Amino Acid Squence: EHSARKWQRKTDPWSLProduct Application DetailsWestern Blot, ELISA, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot 1-2 ug/ml, ELISA, Immunohistochemistry 5 ug/ml, Immunohistochemistry/Immunofluorescence, Immunofluorescence 20 ug/ml, Immunohistochemistry	Clonality	Polyclonal	
PurityPeptide affinity purifiedPurityPeptide affinity purifiedBufferPBSTarget Molecular Weight135 kDaProduct DescriptionIterationHostRabbitGene ID6499Gene SymbolSKIV2LSpeciesHuman, Mouse, RatSpecificity/SensitivityAt least two isoforms of SKIV2L are known to exist; this antibody will detect both isoforms.ImmunogenSKIV2L antibody was raised against a 16 amino acid peptide near the amino terminus of human SKIV2L. The immunogen is located within amino acids 60 - 110 of SKIV2L. Amino Acid Squence: EHSARKWQRKTDPWSLProduct Application DetailsWestern Blot, ELISA, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot 1-2 ug/ml, ELISA, Immunohistochemistry 5 ug/ml, Immunohistochemistry/ Immunohistochemistry 5 ug/ml, Immunohistochemistry/ Immunohistochemistry 5 ug/ml, Immunohistochemistry/ Immunohistochemistry	Preservative	0.02% Sodium Azide	
Buffer PBS Target Molecular Weight 135 kDa Product Description Intervention Host Rabbit Gene ID 6499 Gene Symbol SKIV2L Species Human, Mouse, Rat Specificity/Sensitivity At least two isoforms of SKIV2L are known to exist; this antibody will detect both isoforms. Immunogen SKIV2L antibody was raised against a 16 amino acid peptide near the amino terminus of human SKIV2L. The immunogen is located within amino acids 60 - 110 of SKIV2L. Amino Acid Squence: EHSARKWQRKTDPWSL Product Application Details Western Blot, ELISA, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin Recommended Dilutions Western Blot 1-2 ug/ml, ELISA, Immunohistochemistry 5 ug/ml, Immunohistochemistry/ Immunofluorescence 20 ug/ml, Immunohistochemistry-	Isotype	IgG	
Target Molecular Weight135 kDaProduct DescriptionHostRabbitGene ID6499Gene SymbolSKIV2LSpeciesHuman, Mouse, RatSpecificity/SensitivityAt least two isoforms of SKIV2L are known to exist; this antibody will detect both isoforms.ImmunogenSKIV2L antibody was raised against a 16 amino acid peptide near the amino terminus of human SKIV2L. The immunogen is located within amino acids 60 - 110 of SKIV2L. Amino Acid Squence: EHSARKWQRKTDPWSLProduct Application DetailsWestern Blot, ELISA, Immunocytochemistry/Immunofluorescence, Immunohistochemistry. ParaffinRecommended DilutionsWestern Blot 1-2 ug/ml, ELISA, Immunohistochemistry 5 ug/ml, Immunohistochemistry/Immunofluorescence 20 ug/ml, Immunohistochemistry-	Purity	Peptide affinity purified	
Product DescriptionHostRabbitGene ID6499Gene SymbolSKIV2LSpeciesHuman, Mouse, RatSpecificity/SensitivityAt least two isoforms of SKIV2L are known to exist; this antibody will detect both isoforms.ImmunogenSKIV2L antibody was raised against a 16 amino acid peptide near the amino terminus of human SKIV2L. The immunogen is located within amino acids 60 - 110 of SKIV2L. Amino Acid Squence: EHSARKWQRKTDPWSLProduct Application DetailsWestern Blot, ELISA, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot 1-2 ug/ml, ELISA, Immunohistochemistry 5 ug/ml, Immunohistochemistry/ Immunofluorescence 20 ug/ml, Immunohistochemistry-	Buffer	PBS	
HostRabbitGene ID6499Gene SymbolSKIV2LSpeciesHuman, Mouse, RatSpecificity/SensitivityAt least two isoforms of SKIV2L are known to exist; this antibody will detect both isoforms.ImmunogenSKIV2L antibody was raised against a 16 amino acid peptide near the amino terminus of human SKIV2L. The immunogen is located within amino acids 60 - 110 of SKIV2L. Amino Acid Squence: EHSARKWQRKTDPWSLProduct Application DetailsWestern Blot, ELISA, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot 1-2 ug/ml, ELISA, Immunohistochemistry 5 ug/ml, Immunohistochemistry/ Immunofluorescence 20 ug/ml, Immunohistochemistry-	Target Molecular Weight	135 kDa	
Gene ID6499Gene SymbolSKIV2LSpeciesHuman, Mouse, RatSpecificity/SensitivityAt least two isoforms of SKIV2L are known to exist; this antibody will detect both isoforms.ImmunogenSKIV2L antibody was raised against a 16 amino acid peptide near the amino terminus of human SKIV2L. The immunogen is located within amino acids 60 - 110 of SKIV2L. Amino Acid Squence: EHSARKWQRKTDPWSLProduct Application DetailsWestern Blot, ELISA, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot 1-2 ug/ml, ELISA, Immunohistochemistry 5 ug/ml, Immunocytochemistry/ Immunohistochemistry-Immunohistochemistry-	Product Description		
Gene SymbolSKIV2LSpeciesHuman, Mouse, RatSpecificity/SensitivityAt least two isoforms of SKIV2L are known to exist; this antibody will detect both isoforms.ImmunogenSKIV2L antibody was raised against a 16 amino acid peptide near the amino terminus of human SKIV2L. The immunogen is located within amino acids 60 - 110 of SKIV2L. Amino Acid Squence: EHSARKWQRKTDPWSLProduct Application DetailsWestern Blot, ELISA, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot 1-2 ug/ml, ELISA, Immunohistochemistry 5 ug/ml, Immunocytochemistry/ Immunohistochemistry-Immunohistochemistry-	Host	Rabbit	
SpeciesHuman, Mouse, RatSpecificity/SensitivityAt least two isoforms of SKIV2L are known to exist; this antibody will detect both isoforms.ImmunogenSKIV2L antibody was raised against a 16 amino acid peptide near the amino terminus of human SKIV2L. The immunogen is located within amino acids 60 - 110 of SKIV2L. Amino Acid Squence: EHSARKWQRKTDPWSLProduct Application DetailsWestern Blot, ELISA, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot 1-2 ug/ml, ELISA, Immunohistochemistry 5 ug/ml, Immunohistochemistry/ Immunohistochemistry-	Gene ID	6499	
Specificity/SensitivityAt least two isoforms of SKIV2L are known to exist; this antibody will detect both isoforms.ImmunogenSKIV2L antibody was raised against a 16 amino acid peptide near the amino terminus of human SKIV2L. The immunogen is located within amino acids 60 - 110 of SKIV2L. Amino Acid Squence: EHSARKWQRKTDPWSLProduct Application DetailsWestern Blot, ELISA, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot 1-2 ug/ml, ELISA, Immunohistochemistry 5 ug/ml, Immunohistochemistry/ Immunofluorescence 20 ug/ml, Immunohistochemistry-	Gene Symbol	SKIV2L	
isoforms.ImmunogenSKIV2L antibody was raised against a 16 amino acid peptide near the amino terminus of human SKIV2L. The immunogen is located within amino acids 60 - 110 of SKIV2L. Amino Acid Squence: EHSARKWQRKTDPWSLProduct Application DetailsApplicationsWestern Blot, ELISA, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot 1-2 ug/ml, ELISA, Immunohistochemistry 5 ug/ml, Immunohistochemistry/ Immunofluorescence 20 ug/ml, Immunohistochemistry-	Species	Human, Mouse, Rat	
terminus of human SKIV2L. The immunogen is located within amino acids 60 - 110 of SKIV2L. Amino Acid Squence: EHSARKWQRKTDPWSLProduct Application DetailsApplicationsWestern Blot, ELISA, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot 1-2 ug/ml, ELISA, Immunofluorescence 20 ug/ml, Immunohistochemistry-	Specificity/Sensitivity		
ApplicationsWestern Blot, ELISA, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot 1-2 ug/ml, ELISA, Immunohistochemistry 5 ug/ml, Immunocytochemistry/ Immunofluorescence 20 ug/ml, Immunohistochemistry-	Immunogen	terminus of human SKIV2L. The immunogen is located within amino acids 60 -	
Immunohistochemistry, Immunohistochemistry-Paraffin Recommended Dilutions Western Blot 1-2 ug/ml, ELISA, Immunohistochemistry 5 ug/ml, Immunocytochemistry/ Immunofluorescence 20 ug/ml, Immunohistochemistry-	Product Application Details		
Immunocytochemistry/ Immunofluorescence 20 ug/ml, Immunohistochemistry-	Applications		
	Recommended Dilutions	Immunocytochemistry/ Immunofluorescence 20 ug/ml, Immunohistochemistry-	





images	
Western Blot: SKIV2L Antibody [NBP2-81903] - Analysis of SKIV2L in mouse stomach tissue lysate with SKIV2L antibody at 1 ug/ml.	250-
	130- SKI2W
	100-
Immunocytochemistry/Immunofluorescence: SKIV2L Antibody [NBP2- 81903] - Analysis of SKIV2L in human stomach tissue with SKIV2L antibody at 20 ug/ml.	
Immunohistochemistry: SKIV2L Antibody [NBP2-81903] - Analysis of SKIV2L in human stomach tissue with SKIV2L antibody at 5 ug/ml.	





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-81903

	NBP2-24891	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP] Rabbit IgG Isotype Control
H00006499-Q01-10ug Recombinant Human SKIV2L GST (N-Term) Protein	H00006499-Q01-10ug	Recombinant Human SKIV2L GST (N-Term) Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-81903

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

