Product Datasheet

CD36 Antibody (185-1G2) [Allophycocyanin] NBP2-95271APC

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-95271APC

Updated 10/26/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-95271APC



NBP2-95271APC

CD36 Antibody (185-1G2) [Allophycocyanin]

/ / / /	
Product Information	
Unit Size	0.1 ml
Concentration	Please see the vial label for concentration. If unlisted please contact technical services.
Storage	Store at 4C in the dark.
Clonality	Monoclonal
Clone	185-1G2
Preservative	0.05% Sodium Azide
Isotype	IgG2a Kappa
Conjugate	Allophycocyanin
Purity	Protein A or G purified
Buffer	PBS
Product Description	
Host	Mouse
Gene ID	948
Gene Symbol	CD36
Species	Human
Marker	Platelet & Microvessel Marker
Specificity/Sensitivity	Recognizes a protein of 80kDa-90kDa, identified as CD36 (Workshop IV; Code P-26). Its epitope maps between aa155-183. It is expressed on platelets, monocytes and macrophages, microvascular endothelial cells, erythrocyte precursors, mammary epithelial cells, and some macrophage derived dendritic cells. CD36 acts as a receptor for thrombospondin (TSP), collagen types I, IV and V, P. falciparum malaria-infected erythrocytes, and sickle erythrocytes. It also functions as a scavenger receptor, mediating macrophage uptake of oxidized low-density lipoprotein (LDL) and recognition of apoptotic polymorphonuclear leukocytes (PMN). CD36 plays a role in platelet aggregation, macrophage foam cell development, inflammation, and the tissue ischemia observed in sickle cell disease and cerebral malaria. Note that 1-4% of Japanese and East Asia population lack CD36. This monoclonal antibody blocks adhesion of P. falciparum parasitized red blood cells to CD36 and strongly inhibits collagen-induced platelet aggregation.
Immunogen	This CD36 Antibody (185-1G2) was developed against stimulated human leukocytes
Product Application Details	
Applications	Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunofluorescence
Recommended Dilutions	Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunofluorescence
Application Notes	Optimal dilution of this antibody should be experimentally determined.
4	



Images

CD36 Antibody (185-1G2) [Allophycocyanin] [NBP2-95271APC] - Vial of APC conjugated antibody. APC is optimally excited at 650 nm by the Red laser (633 or 640 nm) and has an emission maximum of 660 nm.







Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-95271APC

NBP1-96981APCMouse IgG2a Kappa Isotype Control (M2AK) [Allophycocyanin]NB400-145PEPCD36 Antibody Blocking Peptide210-TA-005TNF-alpha [Unconjugated]NB400-144PEPCD36 Antibody Blocking Peptide

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-95271APC

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

