Product Datasheet

NDUFC2 Antibody - Azide and BSA Free NBP3-02955-100ul

Unit Size: 100 ul

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-02955

Updated 11/7/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-02955



NBP3-02955-100ul

Recommended Dilutions

NDUFC2 Antibody - Azide and BSA Free


INDOI GZ Allibody - Azide alid ba	SATTEE			
Product Information				
Unit Size	100 ul			
Concentration	Please see the vial label for concentration. If unlisted please contact technical services.			
Storage	Store at -20C. Avoid freeze-thaw cycles.			
Clonality	Polyclonal			
Preservative	0.01% Thimerosal			
Isotype	IgG			
Purity	Affinity purified			
Buffer	PBS with 50% glycerol, pH7.3.			
Product Description				
Host	Rabbit			
Gene ID	4718			
Gene Symbol	NDUFC2			
Species	Human, Mouse, Rat			
Immunogen	A synthetic peptide corresponding to a sequence within amino acids 1-100 of human NDUFC2 (NP_004540.1). MIARRNPEPLRFLPDEARSLPPPKLTDPRLLYIGFLGYCSGLIDNLIRRRPIATAG LHRQLLYITAFFFAGYYLVKREDYLYAVRDREMFGYMKLHPEDF			
Product Application Details				
Applications	Western Blot, Knockout Validated			

Western Blot 1:500 - 1:2000, Knockout Validated




Images

Western Blot: NDUFC2 Antibody [NBP3-02955] - Western blot analysis of extracts of various cell lines, using NDUFC2 antibody (NBP3-02955) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 5s.



Knockout Validated: NDUFC2 Antibody [NBP3-02955] - Analysis of extracts from normal (control) and NDUFC2 knockout (KO) HeLa cells, using NDUFC2 antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit . Exposure time: 3min.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP3-02955-100ul

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

NBP2-46831PEP NDUFC2 Recombinant Protein Antigen

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-02955

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

